

ICPAM-VAN 2018

11-13 SEPTEMBER 2018,
VAN YÜZÜNCÜ YIL UNIVERSITY
VAN-TURKEY



BOOK OF ABSTRACTS AND PROGRAMME

**2ND INTERNATIONAL CONFERENCE ON
PURE AND APPLIED MATHEMATICS**



icpam.yyu.edu.tr



icpam2018@yyu.edu.tr
icpam2018.van@gmail.com



Van Yüzüncü Yıl University
Abdullah Gül University



Edited by Zeynep Kayar



Designed by Fatih Kutlu

Van Yüzüncü Yıl University

**2nd International Conference on Pure and
Applied Mathematics (ICPAM-VAN 2018)**

BOOK OF ABSTRACTS

Van, TURKEY
September 11 - 13, 2018

Sponsors:

Van Yüzüncü Yıl University
Abdullah Gül University

Scientific Committee:

DIFFERENTIAL EQUATIONS

Michal Fečkan Komenský (Comenius) University of Bratislava, SLOVAKIA
Martin Bohner, Missouri University of Science and Technology, Rolla, USA
Vitali I. Slynko, National Academy of Sciences of Ukraine, Kiev, UKRAINE
Samir H. Saker, Mansoura University Mansoura, EGYPT

FUZZY TOPOLOGY

Oscar Castillo, Instituto Tecnológico de Tijuana Tijuana BC, MEXICO
Krassimir Todorov Atanassov, Bulgarian Academy of Sciences Sofia, BULGARIA
Said Melliani, Sultan Moulay Slimane University Beni Mellal, MOROCCO

ANALYSIS

Heybetkulu S. Mustafayev-Seferoglu, Yznc Yil niversitesi (Centennial University) Van, TURKEY
Muhammad Aslam Noor, COMSATS Institute of Information Technology, Islamabad, PAKISTAN
Naim L. Braha, University of Prishtina REPUBLIC OF KOSOVA

NUMERICAL ANALYSIS

Abdon Atangana, University of the Orange Free State Bloemfontein, SOUTH AFRICA

APPLIED MATHEMATICS

Soon-Mo Jung, Hong-Ik University, Jochiwon Campus Chochiwon, REPUBLIC OF KOREA
Abdel-Shafy Fahmy Obada, Al-Azhar University Nasr City, Cairo, EGYPT
Aliakbar M. Haghghi, Prairie View A&M University Prairie View TX, USA

ALGEBRA

Richard M. Low, San Jose State University San Jose CA, USA

Editor:

Zeynep Kayar (Van Yüzüncü Yıl University)

Local Organizing Committee:

Peyami BATTAL (Honorary Board - President of Van Yuzuncu Yil University)

Cemil TUNÇ (Chair - Dean of Faculty of Science)

Nagehan Alsoy Akgün (Scientific Program)

İsmail Hakkı Denizler (Finance Organizer)

Sultan Erdur (Local Organization and Secretary)

Bahar Kalkan (Local Organization and Secretary)

Zeynep Kayar (Scientific Program)

Fatih Kutlu (Web and Technical Support)

Ömer Küsmüş (Local Organization and Secretary)

Murat Luzum (Local Organization and Secretary)

Onur Saldır (Local Organization and Secretary)

Hayri Topal (Local Organization and Secretary)

Ali Hakan Tor (Web and Technical Support)

Osman Tunç (Local Organization and Secretary)

Ramazan Yazgan (Local Organization and Secretary)

Welcome

Dear Participants,

Welcome to 2nd International Conference on Pure and Applied Mathematics, ICPAM - VAN 2018, Van, Turkey. The conference is organized and is to be held at Van Yüzüncü Yıl University from September 11th to September 13th. This is the second ICPAM meeting in Van and the first one was organized in 2015. The reason for our 3-year long break is to be a host of another high involvement conference. We are happy to see you here in Van.

The main aim of this successive conference is to provide participants with an opportunity to exchange the latest information and ideas, and to encourage debate on many issues in international mathematical researches. During the conference you will certainly meet old and new colleagues, exchange ideas, develop new projects. You will also feel and enjoy the special atmosphere of Van and Turkey.

With 9 parallel sessions (totally 36 sessions) and 131 presentations and more than 150 participants from 11 countries, Algeria, China, Egypt, India, Iran, Pakistan, Portugal, Taiwan, Turkey, Ukraine, United Kingdom, as well as people from 37 different university from Turkey, ICPAM - VAN 2018 will provide a stimulating opportunity for a global interchange of ideas on recent advances in mathematics.

I would like to express my deep gratitude to Prof. Dr. Peyami BATTAL, President of Van Yüzüncü Yıl University, for his encouragement and support in all stages of this conference.

I am grateful to all the participants in the International Conference on Pure and Applied Mathematics, particularly the members of the Scientific and Organizing Committees, the referees and the authors for producing such a high standard conference.

The conference is almost entirely from the registration support of participants. Behind this, we are grateful to Rectorate of YYU, Faculty of Sciences of YYU and Abdullah Gül University for their financial support. We would also like to thank to Governor of Van for his generous support. Have a pleasant stay in Van.

Professor Cemil Tunç
Chair Organizer of ICPAM - VAN Organizing Committee

Acknowledgements

Dear Participants,

It is a great pleasure having this opportunity to thank all the people who made 2nd International Conference on Pure and Applied Mathematics (ICPAM-VAN 2018) possible.

We would like to express our gratitude and deep appreciation to Prof. Dr. Cemil Tunc for his precious guidance, continuous encouragement, boundless energy which never seems to end and persuasive support throughout the organization. His great enthusiasm and belief in us make this conference possible.

We extend our sincere thanks to our colleagues who worked with us during the preparation of this conference for their help and support anytime we needed. We are grateful to all the members of Van Yüzüncü Yıl University Mathematics family, academic and administrative, who kindly help us at every stage of this project.

We would also like to thank to the sponsors, Van Yüzüncü Yıl University and Abdullah Gül University for their generous support.

The last but not the least, we would like to give our heartfelt thanks to participants for their understanding, patience and contributions which make our conference more successful. Since the conference is almost entirely from the registration support of participants, we are grateful for their financial support as well. Thank you all again for the unbelievable amount of support and understanding you have shared during this period. It means more than words can express. We were honored and happy to welcome you to in Van. We hope to see you again in Van, the city of the sun.

Zeynep Kayar (Van Yüzüncü Yıl University)

Ali Hakan Tor (Abdullah Gül University)

Contents

Welcome	4
Acknowledgements	5
Scientific programme in details	11
September 11, 2018	11
September 12, 2018	17
September 13, 2018	23
Abstracts of invited speakers	24
DUMITRU BALEANU, Nonlocality and fractional calculus: Finding the best kernel	25
YEN-JEN CHENG, CHIH-WEN WENG, FENG-LEI FAN, A conjecture on the spectral radius of a bipartite graph	26
SNEZHANA HRISTOVA, RAVI AGARWAL, RONALD O'REGAN, Stability of Caputo non-instantaneous impulsive fractional differential equations with delays	27
ALIREZA KHALILI GOLMANKHANEH, Generalized Reimann calculus on fractal sets and curves and application	28
MEHRDAD LAKESTANI, Proximal algorithm for compressive sensing	29
SEDAGHAT SHAHMORAD, YOUNES TALAEI, Review of the Tau method	30
VITALII I. SLYN'KO, CEMİL TUNÇ, Stability of linear switched impulsive systems with unstable subsystems	31
YUSUF YAYLI, Quaternions and their applications	32
Abstracts of participants' talks	33
ASGHAR AHMADKHANLU, VEDAT SUAT ERTURK, Existence of positive solutions for a singular fractional boundary value problem	34
EVRIİM AKALAN, HIDETOSHI MARUBAYASHI, AKIRA UEDA, Strongly graded rings over hereditary Noetherian prime rings	35
IREM AKBULUT, CEMİL TUNÇ, Analysis of behaviors of solutions of a coupled Volterra integro-differential equations	36
TOLGA AKTURK, HASAN BULUT, Modified expansion function method to the nonlinear problem	37
MEHMET AKİF AKYOL, Some properties of conformal generic submersions	38
HAYRİYE ESRA AKYÜZ, Approximate confidence interval based on winsorized mean for the coefficient of variation of positively skewed populations	39
NAGEHAN ALSOY-AKGÜN, Numerical study of nanofluids under DDMC in a lid driven cavity	40
HIFSI ALTINOK, DERYA DENİZ, MİTHAT KASAP, Some properties of sequence class $S^\beta(\Delta, F, f)$ defined by a modulus function	41
HIFSI ALTINOK, MİTHAT KASAP, DERYA DENİZ, Strongly Cesaro summability of order β with respect to a modulus function	42
MEHMET ŞERİF ALDEMİR, ABDALLA KHDİR ABDALLA MANGURI, On eccentricity based indices of generalized Petersen graphs	43
MEHMET ŞERİF ALDEMİR, ABDALLA KHDİR ABDALLA MANGURI, On stratified domination and Zagreb indices	44
DERYA ALTINTAN, HEINZ KOEPPL, Inference algorithms for jump-diffusion approximations of multi-scale processes	45
YENER ALTUN, CEMİL TUNÇ, On the global exponential stability of nonlinear neutral differential equations with time-varying delays	46

YENER ALTUN, CEMİL TUNÇ, ABDULLAH YİĞİT, On the exponential stability in nonlinear neutral differential equations	47
NAZLIM DENİZ ARAL, ZEYNEP SEVİNÇ, On a new type of q -Baskakov-Kantorovich operators	48
DERYA ARSLAN, Second-order difference approximation for nonlocal boundary value problem with boundary layers	49
REŞAT ASLAN, AYDIN İZGİ, Approximation by bivariate Bernstein-Kantorovich operators on a triangular domain	50
ÖZKAN ATAN, FATİH KUTLU, Synchronization control of two chaotic systems via a novel fuzzy control method	51
MERVE ATASEVER, SEZGİN ALTAY DEMİRBAĞ, On the structure of Ricci solitons on gradient Einstein-type manifolds	52
EMINE ATICI ENDES, JONATHAN A. SHERRATT, A non-local model for E and N cadherin-dependent cell-cell adhesion	53
OZGUR AYDOGMUS, Approximating the stochastic evolution via difference equations	54
JAFAR A'ZAMI, MARYAM KHAJEPOUR, Reduction and coreduction of modules	55
SELÇUK BAŞ, TALAT KÖRPİNAR, RIDVAN CEM DEMİRKOL, MUSTAFA YENEROĞLU, Roller coaster surface according to modified orthogonal frame in Euclidean space	56
ŞENAY BAYDAŞ, BÜLENT KARAKAŞ, On mechanisms in three-dimensional Minkowski space	57
HOSHANG BEHRAVESH, MEHDI GHAFFARZADEH, MOHSEN GHASEMI, Groups whose codegree graphs have no triangle	58
MURAT BEKAR, YUSUF YAYLI, Some algebraic properties of elliptic biquaternions	59
ELENA BESPALOVA, NATALIYA YAREMCHENKO, Quasilinearization method in problems of the subcritical deformation of flexible shell systems	60
CANAN BOZKAYA, MHD natural convection flow in a porous cavity	61
LEYLA BUGAY, A general approach to find generating sets of certain finite subsemigroups of symmetric inverse semigroup	62
MURAT CANCAN, SÜLEYMAN EDİZ, MEHMET ŞERİF ALDEMİR, On ve -degrees in Cartesian product of two graphs	63
CANSU CENGİZ, SERKAN ALI DUZCE, Extortion strategies in non-symmetric iterated Prisoner's Dilemma	64
F. AYCA CETINKAYA, ILKNUR AYDIN, Spectral properties of a q -fractional boundary value problem	65
AKRAM CHEHRAZI, TURGUT HANOYMAK, Mathematical aspects of quantum cryptography	66
EMEK DEMİRCİ AKARSU, Random process generated by the short incomplete Gauss sums	67
RIDVAN CEM DEMİRKOL, TALAT KÖRPİNAR, VEDAT ASİL, SELÇUK BAŞ, A new approach to a bending energy of elastica for space curves in De-Sitter space	68
İSMAİL HAKKI DENİZLER, An analogue of the Artin-Rees Lemma for Artinian modules	69
VEDAT DÖRMA, BÜLENT KARAKAŞ, ŞENAY BAYDAŞ, Kinematics of $4R$ and $2RPR$ mechanisms in Clifford algebra	70
UGUR DURAN, MEHMET ACIKGOZ, SERKAN ARACI, A note on q -Fubini polynomials	71
SÜLEYMAN EDİZ, MURAT CANCAN, MEHMET ŞERİF ALDEMİR, On ve -degrees in direct and strong products of two graphs	72
SULTAN ERDUR, CEMİL TUNÇ, On the existence of periodic solutions of third order nonlinear differential equations with multiple delays	73
ESRA ERKAN, SALİM YÜCE, The theory of Bézier curves in \mathbb{E}^4	74
ELİF ERTEM AKBAŞ, MUSTAFA GÖK, Misconceptions regarding representativeness in probability subject of high school students: Van case	75

MELEK GÖZEN, CEMİL TUNÇ, New exponential stability criteria for certain neutral differential equations with interval discrete and distributed time-varying delays	76
EDA GUNAYDIN, YUSUF GUREFE, TOLGA AKTURK, On the kink type and singular solitons solutions to the nonlinear partial differential equation	77
SİNEM GÜLER, Lorentzian homogeneous generalized Ricci solitons of dimension $n \geq 3$. .	78
TUBA GÜLEŞCE TATLI, TURGUT HANOYMAK, ÖMER KÜSMÜŞ, Multi-party key exchange protocol and man in the middle attack	79
HANDE GÜNAY AKDEMİR, Simulation studies for credibility-based multi-objective programming problems with fuzzy parameters	80
HAKKI GÜNGÖR, Regularization of inverse coefficient determination problem in a hyperbolic problem	81
ŞULE YÜKSEL GÜNGÖR, Approximation by summation-integral type operators involving Brenke polynomials	82
KAZEM HAGHNEJAD AZAR, Some notes on the order-to-topology continuous operators	83
TURGUT HANOYMAK, ATILLA BEKTAŞ, On mathematical aspects of blockchain architecture	84
HÜSEYİN IŞIK, A new class of set-valued contractions and related results	85
SEDA İĞRET ARAZ, On numerical solution of an optimal control problem involving hyperbolic equation	86
MOHAMMAD REZA JABBARZADEH, Conditional expectation operators on measurable function spaces	87
BAHAR KALKAN, ŞENAY BAYDAŞ, BÜLENT KARAKAŞ, Inverse kinematics computation for a 6-DOF articulated robot arm using conformal geometric algebra	88
RAMAZAN KAMA, On some vector valued multiplier spaces obtained by Zweir matrix method	89
MAHMUT KARAKUŞ, TUNAY BİLGİN, Λ - matrix as a summability operator and completeness of certain normed spaces via weakly unconditionally Cauchy series	90
YASİN KAYA, The maximal function in Sobolev spaces	91
ZEYNEP KAYAR, A novel Lyapunov type inequality for quasilinear impulsive systems . .	92
ADEL P. KAZEMI, Total dominator coloring of a graph	93
MOHAMMAD BAGHER KAZEMI BALGESHIR, Invariant submanifolds of statistical Kenmotsu manifolds and their curvatures	94
WALEED S. KHEDR, Reduction of Navier-Stokes equation to a linear equation	95
AHMAD KHOJALI, A note on the annihilator of certain local cohomology modules	96
ERDAL KORKMAZ, CEMİL TUNC, Sufficient conditions for global asymptotic stability of neural networks with time-varying delays	97
ZELİHA KÖRPINAR, FATİH COŞKUN, On the cubic nonlinear Shrodinger's equation with repulsive delta potential	98
ZELİHA KÖRPINAR, MUSTAFA İNÇ, On numerical solutions for fractional (1+1)-dimensional Biswas-Milovic equation	99
ZELİHA KÖRPINAR, TALAT KÖRPINAR, SELÇUK BAŞ, M. TALAT SARIAYDIN, On inextensible flow with Schrödinger flow	100
OMER KUSMUS, I. HAKKI DENIZLER, NECAT GORENTAS, Idempotent unit group in commutative group rings of direct products	101
HATİCE KUŞAK SAMANCI, MUHSİN İNCESU, Investigating a quadratic Bezier curve according to N-Bishop frame	102
FATİH KUTLU, ÖZKAN ATAN, Review on fuzzy thermal image processing applications .	103
FATİH KUTLU, FERİDE TUĞRUL, MEHMET ÇİTİL, On temporal intuitionistic fuzzy De Morgan triplets	104
NISAR A. LONE, An interplay between Riemann integrability and weaker forms of continuity	105

ROSELAINÉ NEVES MACHADO, LUIZ GUERREIRO LOPES, Ehrlich-Aberth's type method with King's correction for the simultaneous approximation of polynomial zeros	106
MOHAMMAD REZA MOTALLEBI, Direct sum of neighborhoods in locally convex cones	107
OĞUZ OĞUR, A note on superposition operators in Fibonacci sequence spaces $l_p(F)$	108
SİNEM ONARAN, On contact surgeries and a counterexample	109
RAMAZAN OZARSLAN, ERDAL BAS, AHU ERCAN, Singular eigenvalue problems via Hilfer derivative	110
NURAY ÖKTEM, An efficient TVD-WAF scheme application for the 2D shallow water equations on unstructured meshes	111
ERHAN PIŞKİN, FATMA EKİNCİ, Blow up of solutions for a quasilinear Kirchhoff-type wave equations with degenerate damping terms	112
MURAT POLAT, Semi-tensor bundle and the vertical lift of tensor fields	113
KHALIL UR REHMAN, M. Y. MALIK, New Lie group of transformation for the non-Newtonian fluid flow narrating differential equations	114
MEHMET GIYAS SAKAR, ONUR SALDIR, FEVZİ ERDOĞAN, Reproducing kernel method with Bernstein polynomials for fractional boundary value problems	115
ONUR SALDIR, MEHMET GIYAS SAKAR, FEVZİ ERDOĞAN, A numerical approach for time-fractional Kawahara equation with reproducing kernel method	116
BÜLENT SARAÇ, Rings without a middle class: past and recent	117
UMİT SARP, DAEYOUL KIM, SEBAHATTİN İKİKARDES, Some applications about Mobius function	118
SANEM SEHRIBANOĞLU, Estimation of parameters of Gumbel distribution data	119
GUZİDE SENEL, Some convergences in metric spaces	120
TUGBA SENLİK CERDİK, FULYA YORUK DEREN, NUKET AYKUT HAMAL, Existence of positive solutions for boundary value problems of nonlinear fractional differential equations	121
GÖKHAN SOYDAN, GAMZE SAVAŞ ÇELİK, Elliptic curves containing sequences of consecutive cubes	122
ABDULGANİ ŞAHİN, BÜNYAMİN ŞAHİN, Jones polynomial for graphs of twist knots	123
SERAP ŞAHİNKAYA, TRUONG CONG QUYNH, Kernel stable and uniquely generated modules	124
ERDOĞAN ŞEN, Generalized class of boundary value problems with a constant retarded argument	125
SÜLEYMAN ŞENYURT, ABDUSSAMET ÇALIŞKAN, Dual pole indicatrix curve and surface	126
SÜLEYMAN ŞENYURT, ABDUSSAMET ÇALIŞKAN, Curves and ruled surfaces according to alternative frame in dual space	127
TANFER TANRIVERDİ, Schrödinger equation with potential vanishing exponentially fast	128
HATICE TASKESEN, Blow up of solutions for a stochastic Klein-Gordon equation	129
HATICE TASKESEN, MOHANAD ALALOUSH, On the blow-up of solutions for a stochastic Camassa-Holm equation	130
MUSTAFA TELCİ, Caristi type related fixed point theorems in two metric spaces	131
CESİM TEMEL, Krasnoselskii fixed point theorem for singlevalued operators and multivalued operators	132
SEDAT TEMEL, TUNCAR SAHAN, OSMAN MUCUK, Crossed modules of group-groupoids and double group-groupoids	133
ALİ HAKAN TOR, An optimality condition for non-smooth convex problems via $*$ -subgradient	134
FATİH TUĞRUL, ŞENAY BAYDAŞ, BÜLENT KARAKAŞ, On the orbit surface of two parameter motion	135
CEMİL TUNÇ, Instability in nonlinear functional differential equations of higher order	136

OSMAN TUNÇ, A note on certain qualitative properties of solutions in Volterra integro-differential equations	137
FATMA TUTAR, ŞENAY BAYDAŞ, BÜLENT KARAKAŞ, Galois theory and palindromic polynomials	138
SUMEYRA UCAR, NİHAL YILMAZ ÖZGÜR, Canonical finite Blaschke products and decomposibility	139
OZLEM UGUZ, ALI DEMIRCI, HANZADE HAYKIRI ACMA, SERDAR YAMAN, Applicability of regression analysis on the oxygen enriched combustion of Kutahya-Tuncbilek lignite	140
OZLEM UGUZ, ALI DEMIRCI, HANZADE HAYKIRI ACMA, SERDAR YAMAN, Applicability of regression analysis on the oxygen enriched combustion of Adiyaman-Golbasi lignite	141
RABIA NAGEHAN UREGEN, On uniformly <i>pr</i> -ideals in commutative rings	142
REZVAN VARMAZYAR, On 2-absorbing ideals	143
ZEKİ YALÇINKAYA, ŞENOL KUBİLAY, ALİ SAVRAN, NECLA ÇALIŞKAN, Studying the kinetic parameters and mechanism of the thermal decomposition (dehydration, dehydroxylation and decarbonylation) of some clays using TG traces	144
ZEKİ YALÇINKAYA, ALİ SAVRAN, ŞENOL KUBİLAY, NECLA ÇALIŞKAN, Clay transition (dehydration, dehydroxylation and decarbonylation) kinetics by DTA	145
RAMAZAN YAZGAN, CEMİL TUNÇ, On the weighted pseudo almost periodic solutions of nonlinear functional Nicholson's blowflies equations	146
SERDAL YAZICI, BAYRAM ÇEKİM, A new generalization of Szász operators and its approximation properties	147
TÜLAY YILDIRIM, Characterization of regular morphisms in terms of abelian categories	148
ASUMAN YILMAZ, MAHMUT KARA, Statistical inference for the inverse Weibull distribution	149
FULYA YORUK DEREN, Existence of solutions for a system of coupled fractional boundary value problems	150
AWAIS YOUNUS, HONGWEI LOU, Input distinguishability of linear dynamic control systems	151
Abstracts of posters	152
MOHAMMED SALAH ABDELOUAHAB, SAFA BOURAFA, On Caputo and Riemann-Liouville fractional-order derivatives with fixed memory length	153
ECEM ACAR, AYDIN İZGİ, On approximation by generalized Bernstein-Durrmeyer operators	154
FİLİZ KANBAY, NURTEN VARDAR, A fuzzy methodology on surface representation of greenhouse gas estimation	155
MOHAMMAD BAGHER MOGHIMI, Asymptotic aspect of some functional equations	156
ILKAY YASLAN KARACA, AYCAN SİNANOĞLU ARISOY, Existence of positive solutions for second order impulsive boundary value problems on the half-line	157

Scientific programme in details

September 11, 2018

Amphi 1 Chair: Cemil TUNÇ

10:00-10:30 **Opening Ceremony**

10:30-11:15 **Dumitru BALEANU**

Nonlocality and fractional calculus: Finding the best kernel

11:15-12:00 **Alireza KHALILI GOLMANKHANEH**, Cemil TUNÇ

Generalized Reimann calculus on fractal sets and curves and application

12:00-13:30 Lunch Break

Amphi 1 Chair: Heybetkulu S. MUSTAFAYEV

13:30-14:15 **Snezhana HRISTOVA**, Ravi AGARWAL, Ronald O'REGAN

Stability of Caputo non-instantaneous impulsive fractional differential equations with delays

14:15-15:00 **Vitalii I. SLYN'KO**, Cemil TUNÇ

Stability of linear switched impulsive systems with unstable subsystems

15:00-15:15 Coffee Break

Amphi 2 Chair: Onur SALDIR

15:15-15:35 **MELEK GÖZEN**, Cemil TUNÇ

New exponential stability criteria for certain neutral differential equations with interval discrete and distributed time-varying delays

15:35-15:55 **Sultan ERDUR**, Cemil TUNÇ

On the existence of periodic solutions of third order nonlinear differential equations with multiple delays

15:55-16:15 Hoshang BEHRAVESH, Mehdi GHAFFARZADEH, **Mohsen GHASEMI**

Groups whose codegree graphs have no triangle

16:15-16:35 **Erdal KORKMAZ**, Cemil TUNÇ

Sufficient conditions for global asymptotic stability of neural networks with time-varying delays

16:35-16:50 Coffee Break

Amphi 2 Chair: Bülent SARAÇ

16:50-17:10 **Ramazan YAZGAN**, Cemil TUNÇ

On the weighted pseudo almost periodic solutions of nonlinear functional Nicholson's blowflies equations

17:10-17:30 **Yener ALTUN**, Cemil TUNÇ

On the global exponential stability of nonlinear neutral differential equations with time-varying delays

17:30-17:50 Yener ALTUN, Cemil TUNÇ, **Abdullah YİĞİT**

On the exponential stability in nonlinear neutral differential equations

19:00-20:00 CONFERENCE DINNER

September 11, 2018**Room 101** Chair: Nuray ÖKTEM

- 15:15-15:35 **Canan BOZKAYA**
MHD natural convection flow in a porous cavity
- 15:35-15:55 **Hakkı GÜNGÖR**
Regularization of inverse coefficient determination problem in a hyperbolic problem
- 15:55-16:15 **Emine ATICI ENDES**, Jonathan A. SHERRATT
A non-local model for E and N cadherin-dependent cell-cell adhesion
- 16:35-16:50 Coffee Break

Room 101 Chair: Canan BOZKAYA

- 16:50-17:10 Erhan PIŞKİN, **Fatma EKİNCİ**
Blow up of solutions for a quasilinear Kirchhoff-type wave equations with degenerate damping terms
- 17:10-17:30 **Nuray ÖKTEM**
An efficient TVD-WAF scheme application for the 2D shallow water equations on unstructured meshes
- 17:30-17:50 **Tanfer TANRIVERDI**
Schrödinger equation with potential vanishing exponentially fast
- 19:00-20:00 CONFERENCE DINNER

Room 102 Chair: Sanem SEHRIBANOGLU

- 15:15-15:35 **Derya ALTINTAN**, Heinz KOEPL
Inference algorithms for jump-diffusion approximations of multi-scale processes
- 15:35-15:55 **Hatice TASKESEN**
Blow up of solutions for a stochastic Klein-Gordon equation
- 15:55-16:15 Hatice TASKESEN, **Mohanad ALALOUSH**
On the blow-up of solutions for a stochastic Camassa-Holm equation
- 16:15-16:35 **Hayriye Esra AKYÜZ**
Approximate confidence interval based on winsorized mean for the coefficient of variation of positively skewed populations
- 16:35-16:50 Coffee Break

Room 102 Chair: Derya ALTINTAN

- 16:50-17:10 **Sanem SEHRIBANOGLU**
Estimation of parameters of Gumbel distribution data
- 17:10-17:30 Ozlem UGUZ, **Ali DEMIRCI**, Hanzade HAYKIRI ACMA, Serdar YAMAN
Applicability of regression analysis on the oxygen enriched combustion of Adiyaman-Golbasi lignite
- 17:30-17:50 Ozlem UGUZ, **Ali DEMIRCI**, Hanzade HAYKIRI ACMA, Serdar YAMAN
Applicability of regression analysis on the oxygen enriched combustion of Kutahya-Tuncbilek lignite
- 19:00-20:00 CONFERENCE DINNER

September 11, 2018

Room 103 Chair: Rıdvan Cem DEMİRKOL

15:15-15:35 **Sinem ONARAN**

On contact surgeries and a counterexample

15:35-15:55 **Mehmet Akif AKYOL**

Some properties of conformal generic submersions

15:55-16:15 Zeliha KÖRPINAR, **Talat KÖRPINAR**, Selçuk BAŞ, M. Talat SARIAYDIN

On inextensible flow with Schrödinger flow

16:15-16:35 **Sinem GÜLER**

Lorentzian homogeneous generalized Ricci solitons of dimension $n \geq 3$

16:35-16:50 Coffee Break

Room 103 Chair: Sinem ONARAN

16:50-17:10 **Rıdvan Cem DEMİRKOL**, Talat KÖRPINAR, Vedat ASİL, Selçuk BAŞ

A new approach to a bending energy of elastica for space curves in De-Sitter space

17:10-17:30 **Murat POLAT**

Semi-tensor bundle and the vertical lift of tensor fields

17:30-17:50 **Merve ATASEVER**, Sezgin ALTAY DEMİRBAĞ

On the structure of Ricci solitons on gradient Einstein-type manifolds

19:00-20:00 CONFERENCE DINNER

Room 104 Chair: Turgut HANOYMAK

15:15-15:35 **Şenay BAYDAŞ**, Bülent KARAKAŞ

On mechanisms in three-dimensional Minkowski space

15:35-15:55 **Awais YOUNUS**, Hongwei LOU

Input distinguishability of linear dynamic control systems

15:55-16:15 Süleyman ŞENYURT, **Abdussamet ÇALIŞKAN**

Dual pole indicatrix curve and surface

16:15-16:35 **Abdulgani ŞAHİN**, Bünyamin ŞAHİN

Jones polynomial for graphs of twist knots

16:35-16:50 Coffee Break

Room 104 Chair: Alireza KHALILI GOLMANKHANEH

16:50-17:10 **Osman TUNÇ**

A note on certain qualitative properties of solutions in Volterra integro-differential equations

17:10-17:30 **Irem AKBULUT**, Cemil TUNÇ

Analysis of behaviors of solutions of a coupled Volterra integro-differential equations

17:30-17:50 **Cemil TUNÇ**

Instability in nonlinear functional differential equations of higher order

19:00-20:00 CONFERENCE DINNER

September 11, 2018**Room 105** Chair: Sedat TEMEL

- 15:15-15:35 **Bülent SARAÇ**
Rings without a middle class: past and recent
- 15:35-15:55 **Serap ŞAHİNKAYA**, Truong Cong QUYNH
Kernel stable and uniquely generated modules
- 15:55-16:15 **Evrım AKALAN**, Hidetoshi MARUBAYASHI, Akira UEDA
Strongly graded rings over hereditary Noetherian prime rings
- 16:15-16:35 **Fatih TUĞRUL**, Şenay BAYDAŞ, Bülent KARAKAŞ
On the orbit surface of two parameter motion
- 16:35-16:50 Coffee Break

Room 105 Chair: Ö. AYDOĞMUŞ

- 16:50-17:10 **Leyla BUGAY**
A general approach to find generating sets of certain finite subsemigroups of symmetric inverse semigroup
- 17:10-17:30 **Sedat TEMEL**, Tuncar SAHAN, Osman MUCUK
Crossed modules of group-groupoids and double group-groupoids
- 17:30-17:50 **Mohammad Reza MOTALLEBI**
Direct sum of neighborhoods in locally convex cones
- 19:00-20:00 CONFERENCE DINNER

Room 106 Chair: Özkan ATAN

- 15:15-15:35 **Murat CANSAN**, Süleyman EDİZ, Mehmet Şerif ALDEMİR
On ve -degrees in Cartesian product of two graphs
- 15:35-15:55 **Ramazan KAMA**
On some vector valued multiplier spaces obtained by Zweir matrix method
- 15:55-16:15 **Mohammad Reza JABBARZADEH**
Conditional expectation operators on measurable function spaces
- 16:15-16:35 **Mustafa TELCİ**
Caristi type related fixed point theorems in two metric spaces
- 16:35-16:50 Coffee Break

Room 106 Chair: Mustafa TELCİ

- 16:50-17:10 **Oğuz OĞUR**
A note on superposition operators in Fibonacci sequence spaces $l_p(F)$
- 17:10-17:30 **Özkan ATAN**, Fatih KUTLU
Synchronization control of two chaotic systems via a novel fuzzy control method
- 17:30-17:50 **Khalil Ur REHMAN**, M. Y. MALIK
New Lie group of transformation for the non-Newtonian fluid flow narrating differential equations
- 19:00-20:00 CONFERENCE DINNER

September 11, 2018

Room 107 Chair: Nagehan ALSOY-AKGÜN

15:15-15:35 **Zeynep KAYAR**

A novel Lyapunov type inequality for quasilinear impulsive systems

15:35-15:55 **Serdal YAZICI**, Bayram ÇEKİM

A new generalization of Szász operators and its approximation properties

15:55-16:15 **Şule Yüksel GÜNGÖR**

Approximation by summation-integral type operators involving Brenke polynomials

16:15-16:35 **Reşat ASLAN**, Aydın İZGİ

Approximation by bivariate Bernstein-Kantorovich operators on a triangular domain

16:35-16:50 Coffee Break

Room 107 Chair: Ali Hakan TOR

16:50-17:10 **Tuba GÜLEŞCE TATLI**, Turgut HANOYMAK, Ömer KÜSMÜŞ

Multi-party key exchange protocol and man in the middle attack

17:10-17:30 **Turgut HANOYMAK**, Atilla BEKTAŞ

On mathematical aspects of blockchain architecture

17:30-17:50 **Akram CHEHRAZI**, Turgut HANOYMAK

Mathematical aspects of quantum cryptography

Room 108 Chair: Zeki YALÇINKAYA

15:15-15:35 Nazlım Deniz ARAL, **Zeynep SEVİNÇ**

On a new type of q -Baskakov-Kantorovich operators

15:35-15:55 **Ali Hakan TOR**

An optimality condition for non-smooth convex problems via $*$ -subgradient

15:55-16:15 **Cansu CENGİZ**, Serkan Ali DUZCE

Extortion strategies in non-symmetric iterated Prisoner's Dilemma

16:15-16:35 **Ozgur AYDOGMUS**

Approximating the stochastic evolution via difference equations

16:35-16:50 Coffee Break

Room 108 Chair: Feride TUĞRUL

16:50-17:10 **Waleed S. KHEDR**

Reduction of Navier-Stokes equation to a linear equation

17:10-17:30 **Zeki YALÇINKAYA**, Şenol KUBİLAY, Ali SAVRAN, Necla ÇALIŞKAN

Studying the kinetic parameters and mechanism of the thermal decomposition (dehydration, dehydroxylation and decarbonylation) of some clays using TG traces

17:30-17:50 Zeki YALÇINKAYA, Ali SAVRAN, **Şenol KUBİLAY**, Necla ÇALIŞKAN

Clay transition (dehydration, dehydroxylation and decarbonylation) kinetics by DTA

19:00-20:00 CONFERENCE DINNER

September 11, 2018 Tuesday										
Time	Amphi 1	Amphi 2	Room 101	Room 102	Room 103	Room 104	Room 105	Room 106	Room 107	Room 108
9:00-10:00	Registration									
Chair	C. TUNC									
10:00-10:30	Opening Ceremony									
10:30-11:15	Invited Speaker									
11:15-12:00	Invited Speaker									
12:00-13:30	Lunch Break									
Chair	H. MUSTAFAYEV									
13:30-14:15	Invited Speaker									
14:15-15:00	Invited Speaker									
15:00-15:15	Coffee Break									
Chair	O. SALDIR	N. ÖKTEM	S. ŞEHRİBAN-OĞLU	R. C. DEMİRKOL	T. HANOY-MAK	S. TEMEL	Ö. ATAN	N. ALSOY-AKGÜN	Z. YAL-CINKAYA	
15:15-15:35	M. GÖZEN	C. BOZKAYA	D. ALTIN-TAN	S. ONARAN	Ş. BAYDAŞ	B. SARAÇ	M. CAN-CAN	Z. KAYAR	Z. SEVİNÇ	
15:35-15:55	S. ERDUR	H. GÜNGÖR	H. TAŞKESEN	M. A. AKYOL	A. YOUNUS	S. ŞAHİNKAYA	R. KAMA	S. YAZICI	A. H. TOR	
15:55-16:15	M. GHASEMI	E. ATICI ENDES	M. ALALOUSH	T. KÖRPİNAR	A. ÇALIŞKAN	E. AKALAN	M. R. JAB-BARZADEH	Ş. Y. GÜNGÖR	C. CENGİZ	
16:15-16:35	E. KORKMAZ	E. AKYÜZ	H. E.	S. GÜLER	A. ŞAHİN	F. TUĞRUL	M. TELCİ	R. ASLAN	Ö. AYDOĞMUŞ	
16:35-16:50	Coffee Break									
Chair	B. SARAÇ	C. BOZKAYA	D. ALTIN-TAN	S. ONARAN	A. KHALILI GOLMAN-KHANEH	Ö. AYDOĞ-MUŞ	M. TELCİ	A. H. TOR	F. TUĞRUL	
16:50-17:10	R. YAZGAN	F. EKİNCİ	S. ŞEHRİBAN-OĞLU	R. C. DEMİRKOL	O. TUNÇ	L. BUGAY	O. OĞUR	T. GÜLEŞCE TATLI	W. S. KHEDR	
17:10-17:30	Y. ALTUN	N. ÖKTEM	A. DEMİRCİ	M. POLAT	İ. AKBULUT	S. TEMEL	Ö. ATAN	T. HANOY-MAK	Z. YAL-CINKAYA	
17:30-17:50	A. YİĞİT	T. TAN-RIVERDİ	A. DEMİRCİ	M. ATA-SEVER	C. TUNC	M. R. MO-TALLEBI	K. U. REHMAN	A. CHEHRAZI	Ş. KUBİLAY	
19:00-20:30	CONFERENCE DINNER									

September 12, 2018**Amphi 1** Chair: Yusuf YAYLI

- 9:30-10:15 Yen-Jen CHENG, **Chih-Wen WENG**, Feng-Lei FAN
A conjecture on the spectral radius of a bipartite graph
- 10:15-11:00 **Mehrdad LAKESTANI**
Proximal algorithm for compressive sensing
- 11:00-11:15 Coffee Break

Amphi 1 Chair: Mehrdad LAKESTANI

- 11:15-12:00 **Sedaghat SHAHMORAD**, Younes TALAEI
Review of the Tau method
- 12:00-12:45 **Yusuf YAYLI**
Quaternions and their applications
- 12:45-14:15 Lunch Break

Amphi 2 Chair: Necat GÖRENTAŞ

- 14:15-14:35 **Fulya YORUK DEREN**
Existence of solutions for a system of coupled fractional boundary value problems
- 14:35-14:55 **Tugba SENLIK CERDIK**, Fulya YORUK DEREN, Nuket Aykut HAMAL
Existence of positive solutions for boundary value problems of nonlinear fractional differential equations
- 14:55-15:15 **F. Ayca CETINKAYA**, Ilknur AYDIN
Spectral properties of a q -fractional boundary value problem
- 15:15-15:35 **Asghar AHMADKHANLU**, Vedat Suat ERTURK
Existence of positive solutions for a singular fractional boundary value problem
- 15:35-15:50 Coffee Break
- 15:50-16:10 POSTER PRESENTATIONS

Amphi 2 Chair: Fulya YORUK DEREN

- 16:10-16:30 **Hüseyin IŞIK**
A new class of set-valued contractions and related results
- 16:30-16:50 **Erdoğan ŞEN**
Generalized class of boundary value problems with a constant retarded argument
- 16:50-17:10 **Cesim TEMEL**
Krasnoselskii fixed point theorem for singlevalued operators and multivalued operators

Room 101 Chair: Bahar KALKAN

- 14:15-14:35 **Rabia Nagehan UREGEN**
On uniformly pr -ideals in commutative rings
- 14:35-14:55 **Jafar A'ZAMI**, Maryam KHAJEPOUR
Reduction and coreduction of modules
- 14:55-15:15 **Rezvan VARMAZYAR**
On 2-absorbing ideals
- 15:15-15:35 **Seda İĞRET ARAZ**
On numerical solution of an optimal control problem involving hyperbolic equation
- 15:35-15:50 Coffee Break
- 15:50-16:10 POSTER PRESENTATIONS

September 12, 2018**Room 101** Chair: Ramazan YAZGAN16:10-16:30 **Tolga AKTURK**, Hasan BULUT

Modified expansion function method to the nonlinear problem

16:30-16:50 Eda GUNAYDIN, Yusuf GUREFE, **Tolga AKTURK**

On the kink type and singular solitons solutions to the nonlinear partial differential equation

16:50-17:10 **Zeliha KÖRPINAR**, Mustafa İNÇ

On numerical solutions for fractional (1+1)-dimensional Biswas-Milovic equation

Room 102 Chair: Sultan ERDUR14:15-14:35 **Ahmad KHOJALI**

A note on the annihilator of certain local cohomology modules

14:35-14:55 **Hande GÜNAY AKDEMİR**

Simulation studies for credibility-based multi-objective programming problems with fuzzy parameters

14:55-15:15 **Fatih KUTLU**, Özkan ATAN

Review on fuzzy thermal image processing applications

15:15-15:35 Fatih KUTLU, **Feride TUĞRUL**, Mehmet ÇİTİL

On temporal intuitionistic fuzzy De Morgan triplets

15:35-15:55 Coffee Break

15:55-16:10 POSTER PRESENTATIONS

Room 102 Chair: Hatice TAŞKESEN16:10-16:30 **Asuman YILMAZ**, Mahmut KARA

Statistical inference for the inverse Weibull distribution

16:50-17:10 **Derya ARSLAN**

Second-order difference approximation for nonlocal boundary value problem with boundary layers

Room 103 Chair: Murat LUZUM14:15-14:35 **Hatice KUŞAK SAMANCI**, Muhsin İNCESU

Investigating a quadratic Bezier curve according to N-Bishop frame

14:35-14:55 **Esra ERKAN**, Salim YÜCEThe theory of Bézier curves in \mathbb{E}^4 14:55-15:15 **Selçuk BAŞ**, Talat KÖRPINAR, Rıdvan Cem DEMİRKOL, Mustafa YENEROĞLU

Roller coaster surface according to modified orthogonal frame in Euclidean space

15:15-15:35 **Mohammad Bagher KAZEMI BALGESHIR**

Invariant submanifolds of statistical Kenmotsu manifolds and their curvatures

15:35-15:50 Coffee Break

15:50-16:10 POSTER PRESENTATIONS

September 12, 2018**Room 103** Chair: Fatih KUTLU

- 16:10-16:30 **Gökhan SOYDAN**, Gamze SAVAŞ ÇELİK
Elliptic curves containing sequences of consecutive cubes
- 16:30-16:50 **Süleyman ŞENYURT**, Abdussamet ÇALIŞKAN
Curves and ruled surfaces according to alternative frame in dual space
- 16:50-17:10 **Tülay YILDIRIM**
Characterization of regular morphisms in terms of abelian categories

Room 104 Chair: Ömer KÜŞMÜŞ

- 14:15-14:35 **Elif ERTEM AKBAŞ**, Mustafa GÖK
Misconceptions regarding representativeness in probability subject of high school students: Van case
- 14:35-14:55 **Adel P. KAZEMI**
Total dominator coloring of a graph
- 14:55-15:15 **Nisar A. LONE**
An interplay between Riemann integrability and weaker forms of continuity
- 15:15-15:35 **Nagehan ALSOY-AKGÜN**
Numerical study of nanofluids under DDMC in a lid driven cavity
- 15:35-15:50 Coffee Break
- 15:50-16:10 POSTER PRESENTATIONS

Room 104 Chair: Chih-Wen WENG

- 16:10-16:30 **Mehmet Şerif ALDEMİR**, Abdalla Khdir Abdalla MANGURI
On stratified domination and Zagreb indices
- 16:30-16:50 **Mehmet Şerif ALDEMİR**, Abdalla Khdir Abdalla MANGURI
On eccentricity based indices of generalized Petersen graphs
- 16:50-17:10 **Süleyman EDİZ**, Murat CANCAN, Mehmet Şerif ALDEMİR
On ve -degrees in direct and strong products of two graphs

Room 105 Chair: I. Hakkı DENİZLER

- 14:15-14:35 **Ugur DURAN**, Mehmet ACIKGOZ, Serkan ARACI
A note on q -Fubini polynomials
- 14:35-14:55 **Emek DEMIRCI AKARSU**
Random process generated by the short incomplete Gauss sums
- 14:55-15:15 **Umit SARP**, Daeyoul KIM, Sebahattin IKIKARDES
Some applications about Mobius function
- 15:15-15:35 **Murat BEKAR**, YUSUF YAYLI
Some algebraic properties of elliptic biquaternions
- 15:35-15:50 Coffee Break
- 15:50-16:10 POSTER PRESENTATIONS

September 12, 2018**Room 105** Chair: Şenay BAYDAŞ

- 16:10-16:30 **FATMA TUTAR**, Şenay BAYDAŞ, Bülent KARAKAŞ
Galois theory and palindromic polynomials
- 16:30-16:50 **Bahar KALKAN**, Şenay BAYDAŞ, Bülent KARAKAŞ
Inverse kinematics computation for a 6-DOF articulated robot arm using conformal geometric algebra
- 16:50-17:10 **Vedat DÖRMA**, Bülent KARAKAŞ, Şenay BAYDAŞ
Kinematics of 4R and 2RPR mechanisms in Clifford algebra

Room 106 Chair: Cesim TEMEL

- 14:15-14:35 Elena BESPALOVA, **Nataliia YAREMCHENKO**
Quasilinearization method in problems of the subcritical deformation of flexible shell systems
- 14:35-14:55 **Ramazan OZARSLAN**, Erdal BAS, Ahu ERCAN
Singular eigenvalue problems via Hilfer derivative
- 14:55-15:15 **Onur SALDIR**, Mehmet Giyas SAKAR, Fevzi ERDOGAN
A numerical approach for time-fractional Kawahara equation with reproducing kernel method
- 15:15-15:35 Zeliha KÖRPINAR, **Fatih COŞKUN**
On the cubic nonlinear Shrodinger's equation with repulsive delta potential
- 15:35-15:50 Coffee Break
- 15:50-16:10 POSTER PRESENTATIONS

Room 106 Chair: Hayri TOPAL

- 16:10-16:30 **Sumeyra UCAR**, Nihal YILMAZ ÖZGÜR
Canonical finite Blaschke products and decomposibility
- 16:30-16:50 Roselaine Neves MACHADO, **Luiz Guerreiro LOPES**
Ehrlich-Aberth's type method with King's correction for the simultaneous approximation of polynomial zeros
- 16:50-17:10 **Omer KUSMUS**, I. Hakkı DENİZLER, Necat GORENTAS
Idempotent unit group in commutative group rings of direct products

Room 107 Chair: Mehmet Giyas SAKAR

- 14:15-14:35 Hıfı ALTINOK, Mithat KASAP, **Derya DENİZ**
Strongly Cesaro summability of order β with respect to a modulus function
- 14:35-14:55 Hıfı ALTINOK, **Derya DENİZ**, Mithat KASAP
Some properties of sequence class $S^\beta(\Delta, F, f)$ defined by a modulus function
- 14:55-15:15 **Mahmut KARAKUŞ**, Tunay BİLGİN
 Λ - matrix as a summability operator and completeness of certain normed spaces via weakly unconditionally Cauchy series
- 15:15-15:35 **Yasin KAYA**
The maximal function in Sobolev spaces
- 15:35-15:50 Coffee Break
- 15:50-16:10 POSTER PRESENTATIONS

September 12, 2018

Room 107 Chair: Mahmut KARAKUŞ

16:10-16:30 **Kazem HAGHNEJAD AZAR**

Some notes on the order-to-topology continuous operators

16:30-16:50 **Mehmet Gıyas SAKAR**, Onur SALDIR, Fevzi ERDOGAN

Reproducing kernel method with Bernstein polynomials for fractional boundary value problems

16:50-17:10 **İsmail Hakkı DENİZLER**

An analogue of the Artin-Rees Lemma for Artinian modules

15:50-16:10 POSTER PRESENTATIONS

Mohammed Salah ABDELOUAHAB, Safa BOURAFA

On Caputo and Riemann-Liouville fractional-order derivatives with fixed memory length

Ecem ACAR, Aydın İZGİ

On approximation by generalized Bernstein-Durrmeyer operators

Filiz KANBAY, Nurten VARDAR

A fuzzy methodology on surface representation of greenhouse gas estimation

Mohammad Bagher MOGHIMI

Asymptotic aspect of some functional equations

İlkay YASLAN KARACA, **Aycan SİNANOĞLU ARISOY**

Existence of positive solutions for second order impulsive boundary value problems on the half-line

September 12, 2018 Wednesday										
Time	Amphi 1	Amphi 2	Room 101	Room 102	Room 103	Room 104	Room 105	Room 106	Room 107	
Chair	Y. YAYLI									
9:30-10:15	C. W. WENG									Invited Speaker
10:15-11:00	M. LAKESTANI									Invited Speaker
11:00-11:15										Coffee Break
Chair	M. LAKESTANI									
11:15-12:00	S. SHAH-MORAD									Invited Speaker
12:00-12:45	Y. YAYLI									Invited Speaker
12:45-14:15										Lunch Break
Chair	N. GÖRENTAŞ	S. ER-DUR	B. KALKAN	M. LUZUM	Ö. KÜSMÜŞ	İ. DENİZLER	C. TEMEL	M. SAKAR	G. SAKAR	
14:15-14:35	F. YÖRÜK DEREN	A. KHO-JALI	R. N. ÜRE-GEN	H. KUŞAK SAMANCI	E. ERTEM AKBAŞ	U. DURAN	N. YAREM-CHENKO	D. DENİZ		
14:35-14:55	T. ŞENLİK ÇERDİK	H. GÜNAY AKDEMİR	J. AZAMI	E. ERKAN	A. KAZEMI PILEDARAQ	E. DEMİRCİ AKARSU	R. OZARSLAN	D. DENİZ		
14:55-15:15	F. ÇETİNKAYA	F. KUTLU	R. VAR-MAZYAR	S. BAŞ	N. A. LONE	Ü. SARP	O. SALDIR	M. KARAKUŞ		
15:15-15:35	A. AHMAD-KHANLU	F. TUĞRUL	S. İGRET ARAZ	M. B. KAZEMI	N. ALSOY- AKGÜN	M. BEKAR	F. COŞKUN	Y. KAYA		
15:35-15:50										Coffee Break
15:50-16:10										Poster Presentations
Chair	F. YÖRÜK DEREN	H. TAŞKESEN	R. YAZ-GAN	F. KUTLU	C. WENG	Ş. BAYDAŞ	H. TOPAL	M. KARAKUŞ	K. HAGH-NEJAD AZAR	G. SAKAR
16:10-16:30	H. IŞIK	A. YILMAZ	T. AKTÜRK	G. SOY-DAN	M. Ş. AL-DEMİR	F. TUTAR	S. UCAR	K. HAGH-NEJAD AZAR		
16:30-16:50	E. ŞEN		T. AKTÜRK	S. ŞENYURT	M. Ş. AL-DEMİR	B. KALKAN	L. LOPES	M. SAKAR		
16:50-17:10	C. TEMEL	D. AR-SLAN	Z. KÖRPİNAR	T. YILDIRIM	S. EDİZ	V. DÖRMA	Ö. KÜSMÜŞ	İ. DENİZLER		

Social Programme*September 13, 2018*

8:00-16:00 SOCIAL PROGRAM

8:00 Departure from university campus

9:00-10:30 Van Breakfast

10:30 Departure for Akhdamar Island

11:30-13:00 Visit to Akhdamar Island

13:00 Departure for lunch

14:30-16:00 Lunch (in city centre)

16:00 Return to university campus

September 13, 2018 Thursday	
Time	Program
8:00	Departure from university campus
09:00-10:30	Van Breakfast
10:30	Departure for Akhdamar Island
11:30-13:00	Visit to Akhdamar Island
13:00	Departure for lunch
14:30-16:00	Lunch (in city centre)
16:00	Return to university campus

Abstracts of invited speakers

Nonlocality and fractional calculus: Finding the best kernel

DUMITRU BALEANU^{1,2}

¹Cankaya University, Ankara, Turkey

²Institute of Space Sciences, Bucharest, Romania

emails: ¹dumitru@cankaya.edu.tr

Fractional calculus deals with the study of so-called fractional order integral and derivative operators over real or complex domains, and their applications. During the last decades, the fractional differentiation has drawn increasing attention in the study of so-called anomalous behaviors, where scaling power law of fractional order appears universal as an empirical description of such complex phenomena [1, 3]. In my talk, for a given real world models, I will present and compare the importance of singular and nonsingular kernels in capturing the non-local effects. A special attention will be devoted to the Mittag-Leffler kernels [2].

MSC 2010: 26A33, 33E12, 32A17

Keywords: Nonlocality, fractional calculus, Mittag-Leffler kernels

References

- [1] D. Baleanu, K. Diethelm, E. Scalas and J. J. Trujillo, Fractional Calculus: Models and Numerical Methods. *World Scientific Publishing, Berlin*, 2012.
- [2] D. Baleanu and O. Mustafa, Asymptotic Integration and Stability for Differential Equations of Fractional Order. *World Scientific Publishing, Berlin*, 2015.
- [3] A. Atangana and D. Baleanu, New fractional derivatives with non-local and non-singular kernel: theory and application to heat transfer model. *Thermal Sci.* **20** (2016), no. 2, 763-769.

A conjecture on the spectral radius of a bipartite graph

YEN-JEN CHENG¹, CHIH-WEN WENG², FENG-LEI FAN³

^{1,2}Department of Applied Mathematics, National Chiao Tung University, Taiwan

³Department of Photonics, Harbin Institute of Technology, China

emails: ¹yjc7755.am01g@nctu.edu.tw; ²weng@math.nctu.edu.tw

The spectral radius of a graph is the maximum eigenvalue of its adjacency matrix. Let G be a bipartite graph with e edges and without isolated vertices. It was known that the spectral radius of G is at most the square root of e , and the upper bound is attained if and only if G is a complete bipartite graph. Bhattacharya, Friedland and Peled [1] conjectured that a non-complete bipartite graph which has the maximum spectral radius with given e and bi-order (p, q) is obtained from a complete bipartite graph by adding one vertex and a corresponding number of edges. We find a counter example of this conjecture. Under the additional assumption that e at least $pq - q$ or p at most 5, where p at most q , we prove a weaker version of the above conjecture that drops the non-complete assumption of the bipartite graph G . Our method is based on a new tight upper bound of the spectral radii of bipartite graphs with bi-order (p, q) and a prescribed degree sequence of the part of order p .

MSC 2010: 05C50, 15A18, 05C35, 15A42.

Keywords: Adjacency matrix, bipartite graph, degree sequence, spectral radius

References

- [1] A. Bhattacharya, S. Friedland and U. N. Peled, On the first eigenvalue of bipartite graphs. *Electron. J. Combin.* **15** (2008), R144.

Stability of Caputo non-instantaneous impulsive fractional differential equations with delays

SNEZHANA HRISTOVA¹, RAVI AGARWAL^{2,3}, RONALD O'REGAN⁴

¹Plovdiv University, Plovdiv, Bulgaria

²Texas A&M University-Kingsville, Kingsville, TX 78363, USA

³Florida Institute of Technology, Melbourne, FL 32901, USA

⁴National University of Ireland, Galway, Ireland

emails: ¹snehri@gmail.com; ²agarwal@tamuk.edu; ³donal.oregan@nuigalway.ie

Impulsive differential equations arise from real world problems to describe the dynamics of processes in which sudden, discontinuous jumps occur. Such processes are natural in biology, physics, engineering, etc. In the literature there are two popular types of impulses:

- *instantaneous impulses*- the duration of these changes is relatively short compared to the overall duration of the whole process;
- *non-instantaneous impulses* - an impulsive action, which starts abruptly at a fixed point and its action continues on a finite time interval.

In this talk Caputo fractional differential equations with non-instantaneous impulses and bounded delays are studied. Both basic approaches in the interpretation of the solutions of the fractional equation deeply connected with the presence of non-instantaneous impulses are discussed and illustrated on several examples.

There are several approaches in the literature to study stability, one of which is the Lyapunov approach. Some difficulties have been encountered when one applies the Lyapunov technique to Caputo fractional differential equations. The basic question which arises is the definition of the derivative of the Lyapunov like function along the given fractional equation. Initially a brief overview of the basic fractional derivatives of Lyapunov functions used in the literature is given and their advantages/disadvantages are discussed and illustrated on examples. Lyapunov functions and Razumikhin technique are applied to study stability properties of Caputo fractional differential equations with non-instantaneous impulses and bounded delays. Comparison results using this definition and scalar fractional differential equations are presented and several sufficient conditions for stability, uniform stability, asymptotic stability, Mittag-Leffler stability are established. Several examples are given to illustrate the theory. Also some applications to neural networks with bounded delays and impulsive perturbations acting as non-instantaneous impulses are presented.

MSC 2010: 34A08, 34K37, 34A37, 34K20

Keywords: Caputo fractional derivative, non-instantaneous impulses, Lyapunov functions, Caputo fractional Dini derivative, Razumikhin method

Acknowledgement: Research was partially supported by the Fund NPD, Plovdiv University, No. FP17-FMI-008.

Generalized Reimann calculus on fractal sets and curves and application

ALIREZA KHALILI GOLMANKHANEH

Islamic Azad University, Urmia, Iran

emails: alirezakhalili2002@yahoo.co.in

In this paper, we review the fractal calculus and basic concepts. The Cantor-like sets are considered as the support of functions and their derivatives and integrals are given. The generalized Cantor cubes are presented and fractal calculus is extended on them. As the applications of formalism, the super-, sub-, and normal-diffusion are characterized. More, the mathematical models are suggested for the fractal grating in optic and fractal space-time in quantum and classical mechanics.

MSC 2010: 81Q35, 28A80, 76M60

Keywords: Cantor sets, fractal calculus

References

- [1] A. Parvate and A. D. Gangal, Calculus on fractal subsets of real-line I: Formulation. *Fractals* **17** (2009), no. 1, 53-148.
- [2] A. Parvate and A. D. Gangal, Calculus on fractal subsets of real line II: Conjugacy with ordinary calculus. *Fractals* **19** (2011), no. 3, 271-290.
- [3] A. K. Golmankhaneh, A. Fernandez and D. Baleanu, Diffusion on middle- ξ Cantor sets. <https://arxiv.org/abs/1805.01536>.
- [4] A. K. Golmankhaneh and C. Tunc, On the Lipschitz condition in the fractal calculus. *Chaos, Soliton Fract.* **95** (2017), 140-147.
- [5] A. K. Golmankhaneh and D. Baleanu, Diffraction from fractal grating Cantor sets. *J. Mod. Opt.* **63** (2016), no. 14, 1364-1369.
- [6] A. K. Golmankhaneh and D. Baleanu, Non-local integrals and derivatives on fractal sets with applications. *Open Phys.* **14** (2016), no. 1, 542548.
- [7] A. K. Golmankhaneh and A. S. Balankin, Sub-and super-diffusion on Cantor sets: Beyond the paradox. *Phys. Lett. A* **382** (2018), no. 14, 960-967.

Proximal algorithm for compressive sensing

MEHRDAD LAKESTANI

University of Tabriz, Tabriz, Iran

email: lakestani@tabrizu.ac.ir

Proximal algorithm can be used for solving non-smooth, constrained and large-scale optimization problems [3]. Thus it can be used successfully for image processing purposes [1]. Let $A \in \mathbb{R}^{m \times n}$ with $m < n$ or $m \ll n$ (in compressed sensing), $b \in \mathbb{R}^m$, and $x \in \mathbb{R}^n$. A basis pursuit problem is a constrained minimization problem as follows:

$$\min_{x \in \mathbb{R}^n} \{ \|x\|_1 : Ax = b \}, \quad (1)$$

which gives the solution of the under determined linear system $Ax = b$ with minimal L_1 norm. In this paper we improve and use proximal gradient algorithm to solve basis pursuit and related sparse optimization problems [4, 2].

MSC 2010: 65K10, 49M99, 90C51

Keywords: Proximal algorithms, Compressed sensing, L_1 minimization, Basis pursuit

References

- [1] A. Beck and M. Teboulle, Gradient-Based Algorithms with Applications to Signal Recovery Problems. In: *Convex Optimization in Signal Processing and Communications*, Cambridge University Press (2010), 42-88; doi: 10.1017/CBO9780511804458.003.
- [2] S. Osher, Y. Mao, B. Dong, and W. Yin, Fast linearized Bregman iteration for compressive sensing and sparse denoising. *Commun. Math. Sci.* **8** (2010), no. 1, 93-111.
- [3] N. Parikh and S. Boyd, Proximal algorithms. *Foundations and Trends in Optimization* **1** (2013), no. 3, 123-231.
- [4] W. Yin, Analysis and generalizations of the Linearized Bregman method. *Siam J. Imaging Sciences* **3** (2010), no. 4, 856-877; doi: 10.1137/090760350.

Review of the Tau method

SEDAGHAT SHAHMORAD¹, YOUNES TALAEI²

^{1,2}University of Tabriz, Tabriz, Iran.

emails: ¹shahmorad@tabrizu.ac.ir; ²y_talaei@tabrizu.ac.ir;

The Tau method, introduced for the first time by C. Lanczos [1] in 1938, in order to find approximate solution for some physical problems. In 1956, C. Lanczos and E. L. Ortiz introduced the recursive approach of the Tau method based on Canonical polynomials for numerical study of ordinary differential equation of the form

$$\sum_{i=0}^{\nu} p_i(x)y^{(i)}(x) = f(x), \quad g_j(y) = d_j, \quad j = 1, \dots, \nu \quad a \leq x \leq b, \quad (1)$$

where $p_i(x)$, $f(x)$ are polynomials of finite degree and g_j are some linear functionals acting on $y(x)$ [2, 3]. The Operational approach of the Tau method is based on three simple operational matrices, that introduced for the first time in 1981 by E. L. Ortiz and H. Samara for numerical solution of nonlinear ordinary differential equations [4] and in 2002 it was extend by M. Hosseini and S. Shahmorad for numerical solution of linear integro-differential equations ([5]).

MSC 2010:

Keywords: The Tau method, polynomial solutions, Matrix formulation.

References

- [1] C. Lanczos, Trigonometric interpolation of empirical and analytic functions. *J. Math. and Physics.* **17** (1938), 123–199.
- [2] C. Lanczos, Applied Analysis. *Prentice-Hall, Englewood Cliffs*, 1956.
- [3] E. L. Ortiz, The Tau method, SIAM. *J. Numer. Anal.* **6** (1969), 480–492.
- [4] E. L. Ortiz and H. Samara, An operational approach to the Tau method for the numerical solution of nonlinear differential equations. *Computing.* **27** (1981), no. 4, 15–25.
- [5] M. Hossein Aliabadi, S. Shamoard, A matrix formulation of the Tau method for Fedholm and Volterra linear integro-differential equations estimation. *Korean J. Comput. Appl. Math* **9** (2002), 497–507.

Stability of linear switched impulsive systems with unstable subsystems

VITALII I. SLYN'KO¹, CEMİL TUNÇ²

¹ S. P. Timoshenko Institute of Mechanics of NAS of Ukraine, Kiev, Ukraine

² Van Yuzuncu Yil University, Van, Turkey

emails: ¹vitstab@ukr.net; ²cemtunc@yahoo.com

We consider a Cauchy problem for a linear switched impulsive system [1]

$$\begin{aligned} \frac{dx}{dt} &= A_{\sigma(t)}x(t), \quad t \in (\tau_k, \tau_{k+1}), \\ \Delta x(t) &= B_{\sigma(t)}x(t), \quad t = \tau_k, \quad x(t_0) = x_0, \end{aligned} \quad (1)$$

where $x \in \mathbb{R}^n$, $\sigma(t)$ is a left continuous piecewise constant function which values belonging to the finite set $\{1, \dots, N\}$, $A_m \in \mathbb{R}^{n \times n}$, $m = 1, \dots, N$, $\{\tau_k\}_{k=1}^{\infty} \subset \mathbb{R}$ is an increasing sequence of moments of impulsive action having a single concentration point at infinity, $t_0 < \tau_1$, $x_0 \in \mathbb{R}^n$, $\Delta x(t) = x(t+0) - x(t)$, $B_m \in \mathbb{R}^{n \times n}$, $m = 1, \dots, N$.

Let us define the structural sets of the linear impulsive system $\mathcal{A} = \{A_1, \dots, A_N\}$, $\mathcal{B} = \{B_1, \dots, B_N\}$. With each pair of matrices $(A_m, B_m) \in \mathcal{A} \times \mathcal{B}$ we will associate a positive number θ_m (residence time), so if $\sigma(\tau_k) = m$, then $\tau_{k+1} - \tau_k = \theta_m$. The triple (A_m, B_m, θ_m) defines the subsystem of the hybrid system (1)

$$\begin{aligned} \frac{dz}{dt} &= A_m z(t), \quad t \neq k\theta_m, \\ \Delta z(t) &= B_m z(t), \quad t = k\theta_m, \quad z(t_0) = z_0. \end{aligned} \quad (2)$$

Note that the system (1) is not assumed to be periodic, so Floquet's theory is not applicable in this case.

We propose a new method for studying the stability of the hybrid system (1) for the case when all the matrices of the set \mathcal{A} do not satisfy the Routh-Hurwitz condition, matrices from \mathcal{B} do not satisfy the Schur's condition, and the subsystems (2) are all unstable. The proposed method of investigation is based on the ideas of commutator calculus [2].

MSC 2010: 93D21, 34D20, 34A38, 34A3.

Keywords: Lyapunov's direct method, switched systems, impulsive systems.

Acknowledgement: This work was partially supported by the Ministry of Education and Science of Ukraine project 0116U004691.

References

- [1] A.M. Samoilenko, N. A. Perestyuk, Impulsive differential equations. With a preface by Yu. A. Mitropolskii and a supplement by S. I. Trofimchuk. Translated from the Russian by Y. Chapovsky. World Scientific Series on Nonlinear Science. Series A: Monographs and Treatises, 14. *World Scientific Publishing Co., Inc., River Edge, NJ*, 1995.
- [2] W. Magnus, On the exponential solution of differential equations for a linear operator. *Comm. Pure Appl. Math.* **7** (1954), 649-673.

Quaternions and their applications

YUSUF YAYLI

Ankara University, Ankara, Turkey

email: yayli@science.ankara.edu.tr

Quaternions are brother of complex numbers. Quaternions defined in four dimensional space have quite good applications in three-dimensional kinematics field. In the modeling of robot movements, quaternions comes out as a screw operator and a rotation operator. Another application field is the linear spherical interpolation. Quaternionic fractals are also topics that have been studied in recent years. In quantum physics, they facilitate for the construction of unitary matrices $SU(2)$.

This talk will be about the historical development of quaternions and their application to various fields.

MSC 2010: 11R52, 15B33, 70B10, 70E15.

Keywords: Quaternion, split quaternion, interpolation

References

- [1] K. Shoemake, Animating rotation with quaternion curves. *ACM Siggraph* **19** (1985), no. 3, 245-254.
- [2] E. Pervin and J. A. Webb, Quaternions in Computer Vision and Robotics. In: *Proc. IEEE Conf. on Computer Vision and Pattern Recognition*, Los Alamitos, CA (1983), 382383.
- [3] B. O'Neill, Semi Riemannian Geometry with Applications Storelativity. *Academic Press Inc., London*, 1983.
- [4] D. Mandic and V. Su Lee Goh, The Magic of Complex Numbers. Complex Valued Nonlinear Adaptive Filters. *John Wiley & Sons*, 2009.
- [5] R. Ghadami, J. Rahebi, and Y. Yaylı, Linear interpolation in Minkowski space. *International Journal of Pure and Applied Mathematics* **77** (2012), no. 4, 469-484.
- [6] R. Ghadami, J. Rahebi, and Y. Yaylı, Spline split quaternion interpolation in Minkowski space. *Advances in Applied Clifford Algebras* **23** (2013), no. 4, 849862.
- [7] S. Aslan and Y. Yaylı, Generalized constant ratio surfaces and quaternions. *Kuwait journal of Science* **44** (2017), no. 1, 42-47.
- [8] S. Kızıltuğ and Y. Yaylı, On the quaternionic Mannheim curves of AW(k)-type in Euclidean E3 space. *Kuwait Journal of Science* **42** (2015), no. 2, 128-140.
- [9] Y. Yaylı, Homothetic motions at E^8 with Cayley numbers. *Mech. Mach. Theory* **30** (1995), no. 3, 417-420.
- [10] Y. Yaylı, Homothetic motions at E^4 . *Mechanism and Machine Theory* **27** (1992), no. 3, 303-305.

Abstracts of participants' talks

Existence of positive solutions for a singular fractional boundary value problem

ASGHAR AHMADKHANLU¹, VEDAT SUAT ERTURK²

¹Azərbaycan Şahid Mədani University, Tabriz, Iran

²Ondokuz Mayıs University, Samsun, Turkey

emails: ¹ahmadkhanlu@azaruniv.ac.ir; ²vsetrurk@omu.edu.tr;

Fractional derivative, as an extension of ordinary derivative, is a suitable tool for modeling of various physical phenomena, chemical processes and engineering.

Furthermore, fractional calculus has been found many applications in classical mechanics and the calculus of variations, and is a very useful means for obtaining solutions of nonhomogenous linear ordinary and partial differential equations.

The study of the existence and uniqueness of solution or multiplicity of solutions of initial and boundary value problem, including fractional differential equations, has a lot of importance in theoretical arguments. Remarkable researches in concern with the existence and multiplicity of positive solutions for nonlinear fractional boundary value problems have been done using fixed point theorems up to now (see [1, 3, 4]).

The aim of this work is to study the fractional differential equation

$${}^c D_0^q u(t) + \lambda f(t, u(t), u'(t)) = 0 \quad 1 < q \leq 2, \quad t \in J = [0, 1]$$

where ${}^c D_0^q$ is the Caputo fractional derivative, subject to the boundary conditions

$$u'(0) = \lambda u(0) \quad u'(1) = 0.$$

Sufficient and necessary conditions will be presented for the existence and uniqueness of solution of this fractional boundary value problem.

MSC 2010: 34A08, 34B18, 26A33

Keywords: Boundary value problem, fractional derivative, fixed point theorem,

References

- [1] A. Ahmadkhanlu and M. Jahanshahi, On the existence and uniqueness of solution of initial value problem for fractional order differential equations on time scales. *Bull. Ira. Math. Soc.* **38** (2012), no. 1, 241-252.
- [2] J. F. Douglas, Some applications of fractional calculus to polymer science. In: *Advances in Chemical Physics*, John Wiley & Sons, Inc., **102** (1997), 121-191.
- [3] K. M. Furati and N. Tatar, An existence result for a nonlocal fractional differential problem. *J. Frac. Calculus* **26** (2004), 43-51.
- [4] S. B. Hadid, Local and global existence theorems on differential equations of non-integer order. *J. Fract. Calculus* **7** (1995), 101-105.

Strongly graded rings over hereditary Noetherian prime rings

EVRİM AKALAN¹, HIDETOSHI MARUBAYASHI², AKIRA UEDA³

¹ Hacettepe University, Ankara, Turkey

²Naruto University of Education, Tokushima, Japan

³Shimane University, Shimane, Japan

email: ¹eakalan@hacettepe.edu.tr

Let $R = \bigoplus_{n \in \mathbb{Z}} R_n$ be a strongly graded ring of type \mathbb{Z} where R_0 is a hereditary Noetherian prime ring. In this paper, we investigate and completely describe the structure of projective ideals of R and prove that R is a strongly G-HNP ring.

MSC 2010: 16W50, 16D40, 16A18

Keywords: Strongly graded ring, hereditary Noetherian prime ring, projective ideal

Analysis of behaviors of solutions of a coupled Volterra integro-differential equations

IREM AKBULUT¹, CEMİL TUNÇ²

¹ Siirt University, Siirt, Turkey

² Van Yuzuncu Yil University, Van, Turkey

emails: ¹iremmmatematik@gmail.com; ²cemtunc@yahoo.com

In this paper, we discuss the existence of periodic solutions for a coupled nonlinear Volterra integro-differential equations with delay. We use some theorems of fixed point theory to prove the result to be given here. We compare our results with that can be found in the literature.

MSC 2010: 45D05, 45M10, 45Jxx

Keywords: Volterra integro-differential equations, coupled equations, periodic solution, fixed point theorem

References

- [1] Y. Raffoul, Analysis of periodic and asymptotically periodic solutions in nonlinear coupled Volterra integro-differential systems. *Turkish J. Math.* **42** (2018), no. 1, 108-120.
- [2] T. A. Burton, P. W. Eloe and M. N. Islam, Periodic solutions of linear integro-differential equations. *Math. Nachr.* **147** (1990), 175-184.
- [3] J. Diblk, E. Schmeidel and M. Rikov, Existence of asymptotically periodic solutions of system of Volterra difference equations. *J. Difference Equ. Appl.* **15** (2009), no. 11-12, 1165-1177.
- [4] T. Furumochi, Asymptotically periodic solutions of Volterra difference equations. *Vietnam J. Math.* **30** (2002), 537-550.

Modified expansion function method to the nonlinear problem

TOLGA AKTURK¹, HASAN BULUT²

¹Ordu University, Ordu, Turkey

²Firat University, Elazig, Turkey

emails: ¹tolgaakturk@odu.edu.tr; ²hbulut@firat.edu.tr

In this article, the solutions of the DSW equation are obtained by using the modified expansion function method. Real and imaginary solutions are obtained according to the coefficients obtained from algebraic equation systems. Two and three dimensional graphics of the found solutions are drawn with the Mathematica program by selecting the appropriate parameters.

Drinfel'd-Sokolov-Wilson equation system,

$$u_t + pvv_x = 0, \quad (1)$$

$$v_t + qv_{xxx} + ruv_x + su_xv = 0. \quad (2)$$

MSC 2010: 35C07, 35C08, 35J60

Keywords: The modified expansion function method (MEFM), Drinfel'd-Sokolov-Wilson Equation (DSW), the soliton solutions

References

- [1] V. G. Drinfel'd and V. V. Sokolov, Equations of Korteweg-de Vries type and simple Lie algebras. By (G'/G) -Expansion Method. *Doklady Akademii Nauk SSSR*. **258** (1981), no. 1, 1116.
- [2] G. Wilson, The affine Lie algebra $C_2(1)$ and an equation of Hirota and Satsuma. *Physics Letters A* **89** (1982) no. 7, 332334.
- [3] S. J. H. He and X. H. Wu, Exp-function method for nonlinear wave equations. *Chaos, Solitons and Fractals* **30** (2006), 700–708.
- [4] R. Hirota, B. Grammaticos, and A. Ramani, Soliton structure of the Drinfel'd-Sokolov-Wilson equation. *Journal of Mathematical Physics* **27** (1986), no. 6, 1499–1505.
- [5] H. T. Chen, New double periodic solutions of the classical Drinfeld-Sokolov-Wilson equation. *Numerical Analysis and Applied Mathematics* **1048** (2008), 138–142.
- [6] J. Satsuma, M. J. Ablowitz, Two-dimensional lumps in nonlinear dispersive systems. *J. Math. Phys* **20** (1979), 1496–1503.
- [7] A. Yokus, H. M. Baskonus, T. A. Sulaiman and H. Bulut, Numerical simulation and solutions of the two-component second order KdV evolutionary system. *Numerical Methods for Partial Dif. Eq.* **34** (2018), no.1, 211-227.
- [8] E. Misirli and Y. Gurefe, Exact solutions of the Drinfel'd-Sokolov-Wilson equation using the Exp-function method. *Applied Mathematics and Comp.* **216** (2010), 2623-2627.

Some properties of conformal generic submersions

MEHMET AKİF AKYOL

Bingöl University, Bingöl, Turkey

email: mehmetakifakyol@bingol.edu.tr

Akyol and Şahin [2] introduced the notion of conformal semi-invariant submersions from almost Hermitian manifolds. The present talk deal with the study of conformal generic submersions from almost Hermitian manifolds which extends semi-invariant submersions, generic Riemannian submersions and conformal semi-invariant submersions a natural way. We mention some examples of such maps and obtain characterizations and investigate some properties, including the integrability of distributions, the geometry of foliations and totally geodesic foliations. Moreover, we obtain some conditions for such submersions to be totally geodesic and harmonic, respectively.

MSC 2010: 53C43, 53C20

Keywords: Kähler manifold, Riemannian submersion, generic Riemannian submersion, conformal submersion, conformal generic submersion, vertical distribution.

References

- [1] S. Ali and S. T. Fatima, Generic Riemannian submersions. *Tamkang Journal of Math.* **44** (2013), no. 4, 395-409.
- [2] M. A. Akyol and B. Şahin, Conformal semi-invariant submersions. *Communications in Contemporary Mathematics* **19** (2017), no. 2, 1650011; doi: <http://dx.doi.org/10.1142/S0219199716500115>.
- [3] P. Baird and J. C. Wood, Harmonic morphisms between Riemannian manifolds, London Mathematical Society Monographs, 29, *Oxford University Press The Clarendon Press. Oxford*, 2003.
- [4] A. Bejancu, Geometry of CR-submanifolds. *Kluwer Academic*, 1986.
- [5] B. Y. Chen, Differential geometry of real submanifolds in a Kaehler manifold, *Monatsh. Math.* **91** (1981), 257-274.
- [6] S. Gundmundsson and J. C. Wood, Harmonic morphisms between almost Hermitian manifolds. *Boll. Un. Mat. Ital. B.* **11** (1997), no. 2, 185-197.
- [7] B. O'Neill, The fundamental equations of a submersion. *Mich. Math. J.* **13** (1966), 458-469.
- [8] B. Şahin, Semi-invariant Riemannian submersions from almost Hermitian manifolds. *Canad. Math. Bull.* **56** (2011), 173-182.
- [9] B. Şahin, Riemannian Submersions, Riemannian Maps in Hermitian Geometry, and Their Applications. *Elsevier, Academic Press, Massachusetts, Cambridge*, 2017.
- [10] K. Yano, M. Kon, Structures on Manifolds. *World Scientific, Singapore*, 1984.

Approximate confidence interval based on winsorized mean for the coefficient of variation of positively skewed populations

HAYRIYE ESRA AKYÜZ

Bitlis Eren University, Bitlis, Turkey

email: heakyuz@beu.edu.tr

The coefficient of variation (CV), as an important measure of variation, has been used in many fields such as medicine, biology, physics, finance, toxicology, business, engineering, life insurance and survival analysis. It is free from the unit of measurement and it can be used for comparing the variability of two different populations. In this study are proposed a confidence interval based on winsorized mean for the population coefficient of variation in the skewed distributions. This confidence interval is based on the Bonett (2006) formula which calculates an confidence interval for the standard deviation of non-normal distributions. A simulation study was made to compare this confidence interval and existing confidence intervals in terms of the coverage probability and average width for normal and some skewed distributions. The number of simulation replications is $M = 50.000$ for each case and sample size is used as $n = 15, 25, 50, 100$ using the program written in MATLAB R2016a. Simulation study showed that the coverage probabilities of the proposed confidence interval were very close to nominal confidence level. In addition to, proposed approximate confidence interval also performed well in terms of average width. Wider confidence intervals were obtained for large values of the population coefficient of variation. As a result; it is recommended to use the approximate confidence interval for the coefficient of variation of positively skewed populations.

MSC 2010: 62F10, 62F12, 62F35

Keywords: Average width, coefficient of variation, coverage probability, winsorized mean

References

- [1] D. G. Bonett, Approximate confidence interval for standard deviation of nonnormal distributions. *Comput. Stat. Data. Anal.* **50** (2006), no. 3, 775-782; doi:10.1016/j.csda.2004.10.003.

Numerical study of nanofluids under DDMC in a lid driven cavity

NAGEHAN ALSOY-AKGÜN

Van Yüziüncü Yıl University, Van, Turkey

email: nagehanalsoyakgun@yyu.edu.tr

In this study the advantages of nanofluids on double diffusive mixed convection (DDMC) in a lid-driven cavity is analyzed by solving the velocity-vorticity form of the governing equations along with the energy and concentration equations. Numerical computations are conducted using the dual reciprocity boundary element method (DRBEM). Vorticity transport, energy and concentration equations are transformed to the form of modified Helmholtz equations by discretizing the time derivative terms first. The effects of Reynolds number (Re), Richardson numbers (Ri) and buoyancy ratio (N) for variation in volume fraction from 0 to 0.2 is presented for copper based nanofluid graphically and obtained results are good agreement with the results in [1].

MSC 2010:

Keywords: DRBEM, Mixed Convection, Nanofluid, Thermo-solutal buoyancy forces, Lid-driven

References

- [1] N. Reddy, K. Murugesan, Numerical investigation on the advantages of nanofluids under DDMC in a lid-driven cavity. *Heat Transfer-Asian Research* **46** (2017), 1065-1086.
- [2] N. Alsoy-Akgün and M. Tezer-Sezgin, DRBEM Solution of the thermo-solutal buoyancy induced mixed convection flow problems. *Engineering Analysis with Boundary Elements* **37** (2013), 513-526.

Some properties of sequence class $S^\beta(\Delta, F, f)$ defined by a modulus function

HIFSI ALTINOK¹, DERYA DENİZ², MİTHAT KASAP³

^{1,2}Firat University, Elazığ, TURKEY

³Şırnak University, Şırnak, TURKEY

emails: ¹hifsialtinok@gmail.com; ²deryadeniz485@yandex.com; ³fdd_mithat@hotmail.com

Aizpuru *et al.* [1] defined the f -density of the subset A of \mathbb{N} by using an unbounded modulus function. After then, Bhardwaj [4] introduced f -statistical convergence of order α with respect to a modulus function f for real sequences. In the present paper, we define the sequence class $S^\beta(\Delta, F, f)$ for $0 < \beta \leq 1$, where f is an unbounded modulus function, Δ is a difference operator in sequences of fuzzy numbers and give some inclusion theorems between $S^\beta(\Delta, F, f)$ and the classical sequence classes.

MSC 2010: 40A05, 40A25, 40A30, 40C05, 03E72

Keywords: Statistical convergence, sequence of fuzzy numbers, modulus function, difference operator

References

- [1] A. Aizpuru, M. C. Listan-Garcia, and F. Rambla-Barreno, Density by moduli and statistical convergence. *Quaest. Math.* **37** (2014), 525-530.
- [2] V. K. Bhardwaj and S. Dhawan, f -statistical convergence of order α and strong Cesaro summability of order α with respect to a modulus. *J. Inequal. Appl.* **2015** (2015), no. 332; doi:10.1186/s13660-015-0850-x.
- [3] Ü. Çakan, and Y. Altın, Some classes of statistically convergent sequences of fuzzy numbers generated by a modulus function. *Iranian Journal of Fuzzy Systems* **12** (2015), no. 3, 47-55.
- [4] R. Çolak, Statistical convergence of order α . In: *Modern Methods in Analysis and Its Applications*, New Delhi, India: Anamaya Pub. (2010), 121-129.
- [5] H. Kızılmaz, On certain sequence spaces. *Canad. Math. Bull.* **24** (1981), no. 2, 169-176.
- [6] M. Matloka, Sequences of fuzzy numbers. *BUSEFAL* **28** (1986), 28-37.

Strongly Cesaro summability of order β with respect to a modulus function

HIFSI ALTINOK¹, MİTHAT KASAP², DERYA DENİZ³

^{1,3}Fırat University, Elazığ, TURKEY

²Şırnak University, Şırnak, TURKEY

emails: ¹hifsialtinok@gmail.com; ²fdd_mithat@hotmail.com; ³deryadeniz485@yandex.com

In this study, we generalize and examine the sequence classes $w^\beta(F, f)$, $w^{\beta,0}(F, f)$ and $w^{\beta,\infty}(F, f)$, where f is an unbounded modulus function and $\beta \in (0, 1]$ is a real number, for sequences of fuzzy numbers and examine some inclusion relations between them.

MSC 2010: 40A05, 40A25, 40A30, 40C05, 03E72

Keywords: Sequence of fuzzy numbers, statistical convergence, modulus function, Cesàro summability

References

- [1] A. Aizpuru, M. C. Listan-Garcia, and F. Rambla-Barreno, Density by moduli and statistical convergence. *Quaest. Math.* **37**, (2014), 525-530.
- [2] H. Altınok, Y. Altın and M. Işık, Statistical convergence and strong p -Cesàro summability of order β in sequences of fuzzy numbers. *Iranian J. of Fuzzy Systems* **9** (2012), no. 2, 65-75.
- [3] H. Altınok and M. Kasap, f -statistical convergence of order β for sequences of fuzzy numbers. *Journal of Intelligent & Fuzzy Systems* **33** (2017), 705-712.
- [4] V. K. Bhardwaj and S. Dhawan, f -statistical convergence of order α and strong Cesaro summability of order α with respect to a modulus. *J. Inequal. Appl.* **2015** (2015), no.332; doi:10.1186/s13660-015-0850-x.
- [5] Ü. Çakan, and Y. Altın, Some classes of statistically convergent sequences of fuzzy numbers generated by a modulus function. *Iranian Journal of Fuzzy Systems* **12** (2015), no. 3, 47-55.
- [6] R. Çolak, Statistical convergence of order α . In: *Modern Methods in Analysis and Its Applications*, New Delhi, India: Anamaya Pub. (2010), 121-129.
- [7] M. Matloka, Sequences of fuzzy numbers, *BUSEFAL* **28** (1986), 28-37.

On eccentricity based indices of generalized Petersen graphs

MEHMET ŞERİF ALDEMİR¹, ABDALLA KHDIR ABDALLA MANGURI²

¹Van Yüzüncü Yıl University, Van, Turkey

² University of Sulaimani, Sulaymaniyah, Iraq

emails: ¹msaldemir@yyu.edu.tr; ²abdullah6nasim@gmail.com

In this study, we firstly calculate the eccentric connectivity and connective eccentricity indices for the generalized Petersen graphs.

MSC 2010: 05C12

Keywords: Eccentric connectivity index, Connective eccentricity index, Generalized Petersen graphs

On stratified domination and Zagreb indices

MEHMET ŞERÍF ALDEMİR¹, ABDALLA KHDIR ABDALLA MANGURI²

¹Van Yuzuncu Yil University, Van, Turkey

² University of Sulaimani, Sulaymaniyah, Iraq

emails: ¹abdullah6nasim@gmail.com, ²msaldemir@yyu.edu.tr

In this study, we firstly investigate the relationship between stratified domination number and Zagreb indices.

MSC 2010: 05C12

Keywords: The first Zagreb index, The second Zagreb index, Stratified domination

Inference algorithms for jump-diffusion approximations of multi-scale processes

DERYA ALTINTAN¹, HEINZ KOEPL²

¹Department of Mathematics, Selçuk University, Konya, Turkey

²Department of Electrical Engineering and Information Technology, Technische Universität Darmstadt, Darmstadt, Germany

emails: ¹altintan@selcuk.edu.tr; ²heinz.koepl@bcs.tu-darmstadt.de

In a biochemical system, the abundance of molecular species and the magnitude of reaction rates can vary in a wide range. This diversity leads to hybrid models which combine deterministic and stochastic modeling approaches. We proposed a jump-diffusion approximation to model biochemical processes with multi-scale nature [3]. The idea of the model is to partition reactions into fast, slow groups and to combine Markov chain updating scheme for the slow set with a diffusion (Langevin) approach updating scheme for the fast set. Based on the state vector representation of the jump-diffusion approximation which is defined as a summation of the random time change model and the Langevin equation, we proved that the joint probability density function of jump-diffusion approximation satisfies the hybrid master equation which is the summation of the corresponding chemical master equation and the Fokker-Planck equation [4].

In this study, we develop an inference algorithm to estimate the hidden states/parameters of reaction systems whose posterior distribution satisfies the hybrid master equation. To construct the algorithm, we combine particle filtering/smoothing methods [2] with Gibbs Monte Carlo Markov Chain scheme [1]. To illustrate the method, we implement the algorithm to biochemical processes.

MSC 2010: 60H30, 60J28, 92B05

Keywords: Deterministic modeling, stochastic modeling, diffusion (Langevin) approach, jump-diffusion approximation, chemical master equation, Fokker-Planck equation, Gibbs Monte Carlo Markov Chain

Acknowledgement: This work is supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK) Program no:3501 Grant, no. 115E252.

References

- [1] G. Casella, C. Robert, Monte Carlo Statistical Methods. Springer Texts in Statistics. *Springer*, 2nd edition, 2004.
- [2] A. Doucet, A. M. Johansen, A tutorial on particle filtering and smoothing: Fifteen years later. In : *Nonlinear Filtering Handbook*, Oxford University Press (2011), 656–704.
- [3] A. Ganguly, D. Altntan, and H. Koepl, Jump-diffusion approximation of stochastic reaction dynamics: Error bounds and algorithms. *Multiscale Model. Simul.* **13** (2015), no. 4, 1390–1419.
- [4] D. T. Gillespie, The chemical Langevin and Fokker-Planck equations for the reversible isomerization reaction. *J. Phys. Chem. A* **106** (2002), 5063–5071.

On the global exponential stability of nonlinear neutral differential equations with time-varying delays

YENER ALTUN¹, CEMİL TUNÇ²

^{1,2}Van Yüzüncü Yıl University, Van, Turkey

emails: ¹yener-altun@yahoo.com; ²cemtunc@yahoo.com;

In this study, we investigated the global exponential stability of the zero solution of a neutral differential equation with time-lags. We find sufficient conditions which guarantee the global exponential stability of the zero solution of the equation. We benefit from the Lyapunov functional as a basic tool and the obtained result includes and improves some ones in the literature. An example is given to illustrate the applicability and correctness of the obtained result by MATLAB-Simulink.

MSC 2010: 34K20, 34K40 93D05.

Keywords: Neutral differential equation, global exponential stability, Lyapunov functional, matrix inequality, multiple delays.

References

- [1] H. Chen, Some improved criteria on exponential stability of neutral differential equation. *Adv. Difference Equ.* **2012** (2012), no. 170, 1-9.
- [2] P. Keadnarmol and T. Rojsiraphisal, Globally exponential stability of a certain neutral differential equation with time-varying delays. *Adv. Difference Equ.* **2014** (2014), no. 32, 1-10.
- [3] C. Tunç, Exponential stability to a neutral differential equation of first order with delay. *Ann. Differential Equations* **29** (2013), no. 3, 253256.

On the exponential stability in nonlinear neutral differential equations

YENER ALTUN¹, CEMİL TUNÇ², ABDULLAH YİĞİT³

^{1,2,3} Van Yüzüncü Yıl University, Van, Turkey

emails: ¹yeneraltun@yyu.edu.tr; ²cemtunc@yahoo.com; ³a-yigit63@hotmail.com

In this work, we consider a nonlinear time-varying delay system of neutral equations with periodic coefficients in the form

$$\frac{d}{dt}(y(t) + Dy(t - \tau(t))) = A(t)y(t) + B(t)y(t - \tau(t)) + F(t, y(t), y(t - \tau(t)))$$

where

$$\|F(t, u, v)\| \leq q_1\|u\|^{1+w_1} + q_2\|v\|^{1+w_2}, \quad q_1, q_2, w_1, w_2 > 0$$

We obtain some new estimates characterizing the exponential decay of solutions at infinity and the attraction sets of the zero solution.

MSC 2010: 34K20, 34K40

Keywords: Neutral equation, Lyapunov- Krasovskii functional.

References

- [1] G. V. Demidenko, and I. I. Matveeva, Estimates for solutions to a class of nonlinear time-delay systems of neutral type. *Electron. J. Differential Equations* **2015** (2015), No. 34, 14 pp.
- [2] G. V. Demidenko, and I. I. Matveeva, Exponential stability of solutions to nonlinear time-delay systems of neutral type. *Electronic Journal of Differential Equations* **2016** (2016), No. 19, 1-20.
- [3] M. A. Skvortsova, Asymptotic properties of solutions to systems of neutral type differential equations with variable delay. *Journal of Mathematical Sciences* **205** (2015), no. 3, 455-463.

On a new type of q -Baskakov-Kantorovich operators

NAZLIM DENİZ ARAL¹, ZEYNEP SEVİNÇ²

^{1,2} Bitlis Eren University, Bitlis, Turkey

emails: ¹ndaral@beu.edu.tr; ²z.sevinc13@gmail.com

In this work, we have introduced a new type of q -analogous of Baskakov-Kantorovich operators and investigated their statistical approximation properties. By using a weighted modulus of smoothness, we have given some direct estimations for error in the case $0 < q < 1$.

MSC 2010: 41A36, 41A30, 41A25

Keywords: q -analysis, q -Baskakov-Kantorovich operators

References

- [1] N. I. Mahmudov, Statistical approximation of Baskakov and Baskakov-Kantorovich operators based on the q -integer. *Cent. Eur. J. Math.* **8** (2010), no. 4, 816-826.
- [2] V. Gupta, C. Radu, Statistical approximation properties of q -Baskakov-Kantorovich operators. *Cent. Eur. J. Math.* **7** (2009), no. 4, 809-818.
- [3] E. Şimşek On a new type of q -Baskakov operators. *Süleyman Demirel University, Journal of Natural and Applied Sciences* (2009), no. 4, 809-818.

Second-order difference approximation for nonlocal boundary value problem with boundary layers

DERYA ARSLAN

Ministry of National Education, Bitlis, Turkey

email: ayredlanu@gmail.com

We consider uniform finite difference method on Bakhvalov mesh for a linear singularly perturbed multi-point boundary value problem

$$\begin{aligned}\varepsilon^2 u''(x) + \varepsilon a(x) u'(x) - b(x)u(x) &= f(x), & 0 < x < 1, \\ u(0) &= A, \\ u(1) - \gamma u(l_1) &= B, & 0 < l_1 < 1,\end{aligned}$$

where $0 < \varepsilon \ll 1$ is a small perturbation parameter; A , B and γ are given constants; $a(x) \geq 0$ and $b(x) \geq \beta > 0$; $f(x)$ and $a(x)$ are assumed to be sufficiently continuously differentiable functions in $[0, 1]$. Moreover the solution $u(x) = 0$ and $u(x) = 1$. In this study, we give asymptotic properties of the exact solution. We discretize the problem on a nonuniform mesh and obtain finite difference scheme. Finally, error estimation showed that the proposed method is the second-order uniform convergent independently of the perturbed term ε in the discrete maximum norm.

MSC 2010: 65L10, 65L11, 65L12, 65L15, 65L20, 65L70, 34B10

Keywords: Singular perturbation, finite difference scheme, Bakhvalov mesh, second-order convergence

References

- [1] G. M. Amiraliyev, Difference method for a singularly perturbed initial value problem. *Turkish J. Math.* **22** (1998), 283-294.
- [2] G. M. Amiraliyev and M. Cakir, A uniformly convergent difference scheme for singularly perturbed problem with convective term and zeroth order reduced equation. *International Journal of Applied Mathematics* **2** (2000), no. 12, 1407-1419.
- [3] G. M. Amiraliyev and M. Cakir, Numerical solution of the singularly perturbed problem with nonlocal condition. *Applied Mathematics and Mechanics* **23** (2002), no. 7, 755-764.
- [4] D. Arslan, Finite difference method for solving singularly perturbed multi-point boundary value problem. *Journal of the Institute of Natural and Applied Sciences* **22** (2017), no. 2, 64-75.

Approximation by bivariate Bernstein-Kantorovich operators on a triangular domain

REŞAT ASLAN¹, AYDIN İZGİ²

^{1,2} Harran University, Şanlıurfa, Turkey

emails: ¹resat.aslan@iskur.gov.tr; ²a.izgi@harran.edu.tr

The aim of this paper is to study the convergence and approximation properties of the bivariate Bernstein-Kantorovich operators on a triangular domain. Our operator is defined as below

$$R_n(f; x, y) = \sum_{k=0}^n \sum_{l=0}^{n-k} \varphi_{n,k,l}(x, y) \left(\frac{n+1}{2}\right)^2 \int_{2^{\frac{k+1}{n+1}-1}}^{2^{\frac{k+1}{n+1}-1}} \int_{2^{\frac{l+1}{n+1}-1}}^{2^{\frac{l+1}{n+1}-1}} f(s, t) ds dt$$

where

$$\varphi_{n,k,l}(x, y) = \binom{n}{k} \binom{n-k}{l} \left(\frac{1+x}{2}\right)^k \left(\frac{1+y}{2}\right)^l \left(1 - \frac{1+x}{2} - \frac{1+y}{2}\right)^{n-k-l}$$

The approximation properties were researched and approximation degree of this operator by means of the partial and complete modulus of continuity on the triangular domain were investigated and numerical examples were given and graphics were drawn by Mapple programme. We estimate the order of approximation by Voronovskaja type result and also demonstrate the convergence of the operators R_n to a certain function and the comparison of the convergence of the bivariate Bernstein-Kantorovich operators to the function through illustrations using .

MSC 2010: 41A36, 41A35, 41A10

Keywords: Approximation theory, linear positive operators, partial and complete modulus of continuity, bernstein-kantorovich operators, degree of approximation

References

- [1] S. Bernstein, Démonstration du théorème de Weierstrass, fondée sur le calcul des probabilités. *Commun. Soc. Math. Kharkow* **13** (1912), no. 2, 1-2.
- [2] L. V. Kantorovich, Sur certains développements suivant les polynômes de la forme de S. Bernstein, I, II. *C.R. Acad. URSS* **563-568** (1930), 595-600.
- [3] H. Hacısalihoğlu and A. Hacıyev, Lineer Pozitif Operatör Dizilerinin Yakınsaklığı. *Ankara University, Ankara*, 1995.
- [4] P. P. Korovkin, On convergence of linear positive operators in the space of continuous functions. *Dokl Akad Nauk SSSR* **90** (1953), 961-964.
- [5] V. I. Volkov, On the convergence of sequences of linear positive operators in the space of continuous function of two variable. *Math. Sb. N. S.* **43** (1957), no. 85, 504 (Russian).

Synchronization control of two chaotic systems via a novel fuzzy control method

ÖZKAN ATAN¹, FATİH KUTLU²

^{1,2}Van Yüzüncü Yıl University, Van, Turkey
emails: ¹oatan@yyu.edu.tr; ²fatihkutlu@yyu.edu.tr

In this study, synchronization control will be performed using intuitionistic fuzzy method for synchronization of two chaotic systems. The stability range of the system will be determined by the Lyapunov method and synchronization control will be provided according to this method. The designed synchronization method is compared with the studies in the literature.

MSC 2010: 93C42, 34C28, 94D05

Keywords: Chaotic systems; chaotic synchronization; fuzzy control, intuitionistic fuzzy sets.

References

- [1] T. H. Lee, C. P. Lim, S. Nahavandi, and J. H. Park, Network-based Synchronization of TS Fuzzy Chaotic Systems with Asynchronous Samplings. *Journal of the Franklin Institute* (2018).
- [2] A. Senouci and A. Boukabou, Fuzzy modeling, stabilization and synchronization of multi-scroll chaotic systems. *Optik-International Journal for Light and Electron Optics* **127** (2016), no. 13, 5351-5358.
- [3] M. R. Akbarzadeh-T, S. A. Hosseini, and M. B. Naghibi-Sistani, Stable indirect adaptive interval type-2 fuzzy sliding-based control and synchronization of two different chaotic systems. *Applied Soft Computing* **55** (2017), 576-587.
- [4] W. Wang and Y. Fan, Synchronization of Arneodo chaotic system via backstepping fuzzy adaptive control. *Optik-International Journal for Light and Electron Optics* **126** (2015), no. 20, 2679-2683.
- [5] A. Boulkroune, A. Bouzeriba and T. Bouden, Fuzzy generalized projective synchronization of incommensurate fractional-order chaotic systems. *Neurocomputing* **173** (2016), 606-614.
- [6] S. Mobayen, Chaos synchronization of uncertain chaotic systems using composite nonlinear feedback based integral sliding mode control. *ISA transactions* **77** (2018), 100-111
- [7] S. F. Sarcheshmeh, R. Esmaelzadeh and M. Afshari, Chaotic satellite synchronization using neural and nonlinear controllers. *Chaos, Solitons & Fractals* **97** (2017), 19-27.

On the structure of Ricci solitons on gradient Einstein-type manifolds

MERVE ATASEVER¹, SEZGİN ALTAY DEMİRBAĞ²

^{1,2}Istanbul Technical University, İstanbul, Turkey

emails: ¹atasever17@itu.edu.tr; ²saltay@itu.edu.tr

There has been increasing interest especially on the study of Einstein manifolds and their several generalizations in Riemannian geometry. We say that (M, g) is a gradient Einstein-type manifold if we have

$$\alpha Ric + \beta Hess(f) + \mu df \otimes df = (\rho + \lambda)g, \quad (1)$$

for some $\alpha, \beta, \mu, \rho \in \mathbb{R}$, and $f \in C^\infty(M)$. In this paper we study the notion of gradient Einstein-type structure on a Riemannian manifolds such as gradient Ricci solitons and quasi-Einstein manifolds. Then, some examples for this kind of manifolds will be given in the following part of the paper.

MSC 2010: 53B15, 53B20, 53C21, 53C25

Keywords: Ricci solitons, Gradient Einstein-type manifolds, Quasi-Einstein manifolds, Parallel vector fields, Warped products

References

- [1] G. Catino, P. Mastrolia, D. Monticelli, and M. Rigoli, On the geometry of gradient Einstein-type manifolds. *Pac. J. of Math.* **286** (2016), no. 1, 39-67.
- [2] S. Altay Demirbağ and S. Güler, Rigidity of (m, ρ) -quasi Einstein manifolds. *Math. Nach.* **290** (2017), no. 14-15, 2100-2110.

A non-local model for E and N cadherin-dependent cell-cell adhesion

EMINE ATICI ENDES¹, JONATHAN A. SHERRATT²

^{1,2} Heriot-Watt University, Edinburgh, UK

emails: ¹eae2@hw.ac.uk; ²j.a.sherratt@hw.ac.uk

Cell adhesion molecules, such as Epithelial (E)-cadherin and Neural (N)-cadherin, have an essential importance on the binding of one cell to another at the cell surface. Known also as cell-cell adhesion this process plays a critical role on tissue formation during early embryo development, immune responses and wound healing. All these biological functions require the coordinated movement of cells in particular ways to specific locations. However; it is not clear how cell junctions control this coordinated migration. Due to this fact, we examine throughly the role of E- and N-cadherins on cell migration phenomenon. In other words, we investigate how E- and N-cadherins affect cell direction during migration. In order to understand their impacts on the direction of cell movement, we develop a new continuous mathematical model consisting of two different direction functions that represent E cadherin and N cadherin, respectively.

The model with non-local adhesion term:

$$\underbrace{\frac{\partial u}{\partial t}}_{\text{change in cell density}} = D \underbrace{\frac{\partial^2 u}{\partial x^2}}_{\text{cell diffusion}} - \underbrace{\frac{\partial}{\partial x}(uK(u))}_{\text{adhesion force}} \quad (\text{Mass Conservation Equation})$$

where

$$K(u) = \frac{\phi}{R} \int_{-R}^R \alpha \left(g_1(u(x+x_0))w_1(x_0) + g_2(u(x+x_0))w_2(x_0) \right) dx_0$$

$K(u)$ is the non-local term, α reflects the strength of adhesion force, $g_1(u(x+x_0))$ and $g_2(u(x+x_0))$ are the nature of the forces, and $w_1(x_0)$ and $w_2(x_0)$ describe the direction and magnitude of the force between cells.

MSC 2010: 35R09, 35C07

Keywords: Cell-cell adhesion, Integro-PDE model, E and N-cadherins

Acknowledgement: This research is supported by The Ministry of National Education a Centre for Postgraduate Training (YLSY Program) funded by The Turkish Government, Ankara, Turkey

References

- [1] N. J. Armstrong, K. J. Painter and J. A. Sherratt, A continuum approach to modeling cell-cell adhesion. *J. Theor. Biol.* **243** (2006), 98-113.
- [2] J. T. Nardini, D. A. Chapnick, X. D. Liu and D. M. Bortz, Modeling keratinocyte wound healing dynamics: cell-cell adhesion promotes sustained collective migration. *J. Theor. Biol.* **400** (2016), 103-117.
- [3] R. B. Hazan, G. R. Phillips, R. F. Qiao, L. Norton and S. A. Aaronson, Exogenous expression of N-cadherin in breast cancer cells induces cell migration, invasion, and metastasis. *J. Cell Biol.* **148** (2000), 779-790.

Approximating the stochastic evolution via difference equations

OZGUR AYDOGMUS

Social Sciences University of Ankara, Ankara, Turkey

email: ozgur.aydogmus@asbu.edu.tr

We consider a chain-binomial process modeling the evolution of behavioral traits in a population. Mean field equations of the model are found and analyzed. The behavior of the chain-binomial process is probabilistically linked to the mean field equation. As a result of this link, we are able to show that the mean fixation time is an exponentially increasing function of time if there exist an interior evolutionary stable state. We also present simulation results for the process to validate our analytical findings.

MSC 2010: 91A22, 60J10, 39A30

Keywords: evolutionary game theory, difference equations, chain-binomial processes

Reduction and coreduction of modules

JAFAR A'ZAMI¹, MARYAM KHAJEPOUR²

^{1,2}University of Mohaghegh Ardabili, Ardabil, Iran

emails: ¹jafar.azami@gmail.com; ²maryamkhajepour@uma.ac.ir

Throughout this paper, all rings are commutative rings with identity and all modules are unital. Let R be a ring and M be an R -module and N, K be submodules of M . The product of N and K is defined as $NK = (N : M)(K : M)M$. (see [4]). Let M be an R -module and N, K be submodules of M . We say that K is a reduction of N , if $K \subseteq N$ and there exists a natural number s such that $KN^s = N^{s+1}$. Let M be an R -module and N, K be submodules of M such that $N \subseteq K$. We say that K is a coreduction of N , if there exists a natural number s such that $(0 :_M \text{Ann}(K)\text{Ann}^s(N)) = (0 :_M \text{Ann}^{s+1}(N))$. We denote it by $C(KN^s) = C(N^{s+1})$. In this paper, we prove some relations about this notions over multiplication and comultiplication modules.

MSC 2010: 13C13, 13E15

Keywords: Reduction, integral closure, multiplication modules, integrally dependent

References

- [1] Z. Abd El-Bast and P. F. Smith, Multiplication modules. *Comm. Algebra* **16** (1988), no. 4, 755-779.
- [2] M. M. Ali, Idempotent and nilpotent submodules of multiplication modules for associated primes of local cohomology modules. *Comm. Algebra* **36** (2008), no. 12, 4620-4642.
- [3] M. M. Ali, Residual submodules of multiplication modules. *Beitr. Algebra Geom.* **46** (2005), no. 2, 405-422.
- [4] H. Ansari-Toroghy and F. Farshadifar, Product and dual product of submodules. *Far East J. Math. Sci.* **25** (2007), no. 3, 447-455.
- [5] M. P. Brodmann and R. Y. Sharp, Local Cohomology: An Algebraic Introduction with Geometric Applications. *Cambridge University Press, Cambridge*, 1998.

Roller coaster surface according to modified orthogonal frame in Euclidean space

SELÇUK BAŞ¹, TALAT KÖRPINAR², RIDVAN CEM DEMİRKOL³, MUSTAFA YENEROĞLU⁴

^{1,2,3}Muş Alparslan University, Muş, Turkey

⁴Fırat University, Elazığ, Turkey

emails: ¹selcukbas79@hotmail.com; ²talatkorpinar@gmail.com; ³rcdemirkol@gmail.com;
⁴mustafayeneroglu@gmail.com

In this study, Roller Coaster surfaces according to modified orthogonal frame is introduced in Euclidean space 3-space. The Gaussian curvature, mean curvature, first and second fundamental form of coefficients of Roller Coaster surfaces of are examined. Then, we obtain some characterizations of Roller Coaster surfaces in the Euclidean space 3-space.

MSC 2000: 53A04, 53A05

Keywords: Modified orthogonal frame, roller coaster surfaces, Gaussian curvature, mean curvature

References

- [1] L. Cui, D. Wang and J. S. Dai, Kinematic geometry of circular surfaces with a fixed radius based on Euclidean invariants. *Journal of Mechanical Design* **131** (2009), 101009.
- [2] D. Y. Kwon , F. C. Park and D. P. Chi, Inextensible flows of curves and developable surfaces. *Appl. Math. Lett.* **18** (2005), 1156-1162.
- [3] S. Izumiya, S. Saji and N. Takeuchi, Circular surfaces. *Commun Advances in Geometry* **7** (2005), 295-313.

On mechanisms in three-dimensional Minkowski space

ŞENAY BAYDAŞ¹, BÜLENT KARAKAŞ²

^{1,2}Van Yüzüncü Yıl University, Van, Turkey

emails: ¹senaybaydas@gmail.com; ²bulentkarakas@gmail.com;

In Euclidean space a mechanism is designed by Denavit-Hartenberg representation. In the Minkowski 3-space, the rotations about the standard spacelike coordinate axes through the hyperbolic angle are represented with the orthonormal matrices. In this paper, a mechanism is designed in three-dimensional Minkowski space with D-H parameters.

MSC 2010: B3B30, 70B10

Keywords: Denavit-Hartenberg parameters, Lorentzian geometry, space-like vector

References

- [1] J. G. Ratclie, Foundations of Hyperbolic Manifolds. *Springer, New York*, 2006.
- [2] O. Nesovic, On rotation about lightlike axis in three-dimensional Minkowski space. *Adv. App. Clifford Algebras* **26** (2016), 237-251.
- [3] S. B. Niku, Introduction to Robotics. *Prentice Hall, New Jersey*, 2001.
- [4] B. Oneill, Semi- Riemannian Geometry with Application to Relativity. *Academic Press, New York*, 1983.

Groups whose codegree graphs have no triangle

HOSHANG BEHRAVESH¹, MEHDI GHAFARZADEH², MOHSEN GHASEMI³

^{1,3} Urmia University, Urmia, Iran

² Islamic Azad University, Khoy, Iran

emails: ¹h.behraves@urmia.ac.ir; ²ghaffarzadeh@iaukhoy.ac.ir; ³m.ghasemi@urmia.ac.ir

For a character χ of a finite group G , the number $\text{cod}(\chi) = |G : \text{Ker}(\chi)|/\chi(1)$ is called the codegree of χ . The codegree graph $\Gamma(G)$ is a graph whose vertex set is the all primes dividing some codegree of a character of G and there is an edge between two distinct primes p and q if pq divides some codegree of a character of G . In this paper, we show that if G is a finite group whose codegree graph has no triangle then $\Gamma(G)$ has at most 5 vertices.

MSC 2010: 20C15, 20D05

Keywords: finite group, irreducible character, codegree graph

References

- [1] D. Chillag, A. Mann and O. Manz, The co-degrees of irreducible characters. *Israel J. Math* **73** (1991), 207–223.
- [2] N. Du, M. Lewis, Codegrees and nilpotence class of p -groups. *J. Group Theory* **19** (2016), no. 4, 561–568

Some algebraic properties of elliptic biquaternions

MURAT BEKAR¹, YUSUF YAYLI²

¹Gazi University, Polatli/Ankara, Turkey

²Ankara University, Ankara, Turkey

emails: ¹muratbekar@gazi.edu.tr; ²yayli@science.ankara.edu.tr

In this study, firstly, we give a brief summary of the concepts elliptic numbers, real quaternions and complex quaternions. Afterwards, we consider the algebra of elliptic biquaternions and give some algebraic properties of this algebra.

MSC 2010: 11R52, 53A17, 53A35

Keywords: Elliptic number, real quaternion, complex quaternion, elliptic biquaternion

References

- [1] W. R. Hamilton, Lectures on Quaternions. *Hodges and Smith, Dublin*, 1853.
- [2] M. Bekar and Y. Yayli, Involutions of complexified quaternions and split quaternions. *Adv. Appl. Cliff. Algeb.* **23** (2013), no. 2, 283-299; doi:10.1007/s00006-012-0376-y

Quasilinearization method in problems of the subcritical deformation of flexible shell systems

ELENA BESPALOVA¹, NATALIYA YAREMCHENKO²

^{1,2}Institute of mechanics of NAS of Ukraine, Kiev, Ukraine

emails: ¹elena.bespalova@ukr.net; ²nataliya.petrivna@ukr.net

Compound systems of flexible shells with various geometry and structure represent design schemes of many constructions of modern engineering. Investigation of elastic deformation of such systems necessitates to solve nonlinear boundary-value problems for systems of high-order differential equations, which meets with certain computational difficulties even for advanced computer engineering.

In the report, the technique for determining the stress-strain state of compound shells within the wide range of acting loads up to their limiting critical values is proposed. It combines the Newton-Kantorovich-Raphson linearization method (quasilinearization method) [1] and the orthogonal-sweep method [2]. For the technique, the quadratic convergence of the process of successive approximations and high accuracy in solving linearized problems are typical.

Values of the critical loads are determined using a computational criterion by which $\lambda_{cr}^{lim} \in [\lambda_{n-1}, \lambda_n]$, where λ_{n-1} is the maximum value of the load at which the process monotonically converges, λ_n is its minimum value at which the conditions of monotonic convergence are not fulfilled ($n = 1, 2, \dots$ is the step of the iteration process).

Using, as an example, a space-rocket apparatus, it is shown that the process of successive approximations $\varepsilon = \varepsilon(n)$ (ε is a certain chosen characteristic) may be of various nature, have different convergence rapidity and contains in a number of cases very shallow segments (plateau) depending on the geometrical and physical features of the object being studied. Such unsuspected character of this process demands enhanced attention in introducing automation in calculations of the subcritical state of shell systems.

Keywords: flexible shell systems, nonlinear boundary-value problems, quasilinearization method, orthogonal-sweep method, analysis

References

- [1] R. E. Bellman, R. E. Kalaba, Quasilinearization and nonlinear boundary-value problems. *Am Elsevier Publ. Comp. INC, New York*, 1965.
- [2] S. K. Godunov, Numerical solution of boundary-value problems for systems of linear ordinary differential equations. *Uspekhi Mat. Nauk.* **16** (1961), no. 3, 171-174.

MHD natural convection flow in a porous cavity

CANAN BOZKAYA

Middle East Technical University, Ankara, Turkey

email: bcanan@metu.edu.tr

A numerical investigation of natural convection flow in a cavity filled with a fluid-saturated porous medium in the presence of uniform magnetic field is performed. The steady, viscous, incompressible flow inside the porous medium is assumed to obey the Darcy law. The fluid physical properties are constant except the density in the body force term which is treated according to Boussinesq approximation. The fluid and porous medium are in thermal equilibrium. The governing equations subject to appropriate boundary conditions are solved by using the dual reciprocity boundary element method (DRBEM) which transforms the differential equations into equivalent boundary integral equations by treating the non-homogeneity through a radial basis function approximation. A parametric study illustrating the influence of the physical parameters on the flow and heat transfer characteristics is carried out and the results are visualized in terms of the streamlines, isotherms and the average Nusselt number.

MSC 2010: 65N38, 76M15, 76R10, 76W05

Keywords: MHD, cavity, porous medium, DRBEM

References

- [1] R. U. Haq, F. A. Soomro, T. Mekkaoui, Q. M. Al-Mdallal, MHD natural convection flow enclosure in a corrugated cavity filled with a porous medium. *Int. J. Heat. Mass. Tran.* **121** (2018), 1168-1178; doi:10.1016/j.ijheatmasstransfer.2018.01.063.
- [2] I. A. Badruddin, A. A. A. Al-Rashed, N. J. S. Ahmed, S. Kamangar, K. Jeevan, Natural convection in a square porous annulus. *Int. J. Heat. Mass. Tran.* **55** (2012), 7175-7187; doi:10.1016/j.ijheatmasstransfer.2012.07.034.

A general approach to find generating sets of certain finite subsemigroups of symmetric inverse semigroup

LEYLA BUGAY

Çukurova University, Adana, Turkey

email: ltanguler@cu.edu.tr

Let I_n be the inverse semigroup consists of all partial bijections on $X_n = \{1, \dots, n\}$ which is called symmetric inverse semigroup. It is known from Wagner-Preston Theorem that every finite inverse semigroup is isomorphic to a subsemigroup of a suitable symmetric inverse semigroup. Hence the symmetric inverse semigroups and their subsemigroups have an important role in Inverse Semigroup Theory like as the symmetric groups in Group Theory. Let $\alpha \in I_n$. Then α is called *isotone* (*antitone*) if $x < y \Rightarrow x\alpha < y\alpha$ ($x < y \Rightarrow x\alpha > y\alpha$) for $\forall x, y \in \text{dom}(\alpha)$, and α is called *monotone* if α is isotone or antitone. Clearly the set of all monotone partial bijections is a subsemigroup of I_n , denoted by $PODI_n$, and also, for $0 \leq r \leq n - 1$, $PODI_{n,r} = \{\alpha \in PODI_n : |\text{im}(\alpha)| \leq r\}$ is a subsemigroup of $PODI_n$. In this talk we give a new and general approach to find any generating set of $PODI_{n,r}$ by using digraphs.

MSC 2010: 20M20

Keywords: Partial bijection, monotone map, generating set

Acknowledgement: My sincere thanks are due to Prof. Dr. Hayrullah Ayık for his helpful suggestions and encouragement.

References

- [1] V. H. Fernandes, Semigroups of order preserving mappings on a finite chain. A New class of Divisors. *Semigroup Forum* **54** (1997), 230–236.
- [2] V. H. Fernandes, The monoid of all injective order-preserving partial transformations on a finite chain. *Semigroup Forum* **62** (2001), 178–204.
- [3] V. H. Fernandes, G. M. S. Gomes and M. M. Jesus, Presentations for some monoids of injective partial transformations on a finite chain. *Southeast Asian Bulletin of Mathematics* **28** (2004), 903–918.
- [4] G. M. S. Gomes and J. M. Howie, On the ranks of certain semigroups of order-preserving transformations. *Semigroup Forum* **45** (1992), no:3, 272–282.
- [5] P. Zhao and V. H. Fernandes, The ranks of ideals in various transformation monoids. *Comm. Algebra* **43** (2015), 674–692.

On ve -degrees in direct and strong products of two graphs

MURAT CANCAN¹, SÜLEYMAN EDİZ², MEHMET ŞERİF ALDEMİR³

^{1,2,3} Van Yuzuncu Yil University, Van, Turkey

emails: ¹mcancan@yyu.edu.tr; ²suleymanediz@yyu.edu.tr; ³msaldemir@yyu.edu.tr

Let G be a graph and v be a vertex of G . The ve -degree of the vertex v defined as the number of different edges incident to the vertices of the open neighborhood of v . In this study we investigate the ve -degrees in Cartesian product of two graphs.

MSC 2010: 05C07

Keywords: ve -degree, Cartesian product, Graph operation

Extortion strategies in non-symmetric iterated Prisoner's Dilemma

CANSU CENGİZ¹, SERKAN ALI DUZCE²

^{1,2} Anadolu University, Eskisehir, Turkey

emails: ¹cansucengiz@anadolu.edu.tr; ²saduzce@anadolu.edu.tr

Good strategies are described for the memory-one symmetric games [1, 2]. Iterated Prisoners Dilemma (IPD) is a symmetric game that players get same payoffs in similar situations with cooperation and defection. The symmetric version of IPD is suitable for evolutionary games, but classic game theory payoffs have to be given in terms of utility functions, which measure the preferences of the players. When interpersonal comparison of utilities is excluded, symmetric game is not suitable. In our study, it is investigated that extortioner player can enforce a different extortionate share in non-symmetric IPD games where players gain different payoffs. Extortioner player gains more than the other player out of their payoffs.

MSC 2010: 91A05, 91A12, 91A20

Keywords: Iterated Prisoner's Dilemma, good Strategy, extortion strategy

References

- [1] E. Akin, Good strategies for the iterated Prisoner's Dilemma. *ArXiv 1211.0969,v.2* (2013).
- [2] E. Akin, What you gotta know to play good in the iterated Prisoner's Dilemma. *Games* **6** (2015), 175–190.
- [3] W. Press and F. Dyson Iterated Prisoner's Dilemma contains strategies that dominate any evolutionary opponent. *PNAS* **109** (2012), no. 26, 10409–10413.
- [4] C. Hilbe, M. Nowak and K. Sigmund, The evolution of extortion in iterated Prisoner's Dilemma games. *PNAS* **110** (2013), no. 17, 6913–6918.
- [5] A. J. Stewart and Joshua B. Plotkin, From extortion to generosity, evolution in the Iterated Prisoner's Dilemma. *PNAS* **110** (2013), no. 38, 15348–15353.

Spectral properties of a q -fractional boundary value problem

F. AYCA CETINKAYA¹, ILKNUR AYDIN²

^{1,2}Mersin University, Mersin, Turkey

emails: ¹faycacetinkaya@mersin.edu.tr; ²aydnilknur95@gmail.com

This paper deals with a boundary value problem which is generated by a differential equation with q -Jackson derivative and a discontinuous weight function. The interval, in which the boundary value problem is defined, is finite. By modifying some techniques of [1,2] and [3] we investigate the spectral properties of the above-mentioned boundary value problem.

MSC 2010: 34B08, 34L05

Keywords: q -Jackson derivative, Sturm-Liouville operator, eigenvalues and eigenfunctions

References

- [1] V. Kac and P. Cheung, Quantum Calculus. *Springer-Verlag, New York*, 2002.
- [2] M. H. Annaby and Z. S. Mansour, Basic Sturm-Liouville Problems. *J. Phys. A: Math. Gen.* **38** (2005), 3775-3797.
- [3] M. H. Annaby and Z. S. Mansour q -Fractional Calculus. *Springer-Verlag, New York*, 2012.

Mathematical aspects of quantum cryptography

AKRAM CHEHRAZI¹, TURGUT HANOYMAK²

¹Azərbaycan Şahid Mədani University, Tabriz, Iran

²Van Yüzüncü Yıl University, Van, Turkey

emails: ¹achehrazi95@gmail.com; ²turguthanoymak@gmail.com;

The integer factorization problem is known to be one of the hardest in mathematics. In modern cryptography, RSA encryption algorithm, which is highly practical and widespread, is based on the integer factorization problem. No efficient classical algorithm for the factorization of large number is known. In 1994, Peter Shor proposed an algorithm for this problem which runs in polynomial time on quantum computer. The scope of this study covers the fundamental concepts of quantum cryptography, quantum computation and the basics of Shor's quantum algorithm.

MSC 2010: 94A60, 94A62, 68P25

Keywords: Superposition principle, qubit, density matrix, Shor's factoring algorithm

References

- [1] C. Bennett and G. Brassard, Quantum Cryptography: Public Key Distribution and Coin Tossing. *International Conference on Computers, Systems and Signal Processing, Bangalore, India, 1984.*
- [2] T. Curcic, M. E. Filipkowski, A. Chtchelkanova, P. A. D'Ambrosio, S. A. Wolf, M. Foster and D. Cochran, Quantum networks: from quantum cryptography to quantum architecture. *ACM SIGCOMM Computer Communication Review* **34** (2004), 3-8.
- [3] C. Elliott, D. Pearson and G. Troxel, Quantum cryptography in practice. *Proceedings of the 2003 Conference on Applications, Technologies, Architectures and Protocols for Computer Communications. ACM, 2003.*
- [4] C. E. Shannon, Communication Theory of Secrecy Systems. *Bell System Technical J.* **28** (1949), no. 4, 656-715.
- [5] G. S. Vernam, Cipher Printing Telegraph Systems for Secret Wire and Radio Telegraphic Communications. *J. Am. Inst. Electrical Eng.* **45** (1926), 109-115.
- [6] P. W. Shor, Polynomial-Time Algorithms for Prime Factorization and Discrete Logarithms on a Quantum Computer. *SIAM J. Comp.* **26** (1997), no. 5, 1484-1509.
- [7] K. P. Arya, M. S. Aswal and V. Kumar, Comparative study of asymmetric key cryptography algorithms. *International Journal of Computer Science and Communication Networks* **5** (2011), 17-21.

Random process generated by the short incomplete Gauss sums

EMEK DEMIRCI AKARSU

Recep Tayyip Erdogan University, Rize, Turkey

email: emek.akarsu@erdogan.edu.tr

In this talk we investigate a random process generated by the short incomplete Gauss sums and establish an analog of weak invariance principle for these sums. A generalization of the limit distribution of the short incomplete Gauss sums given by theta sums on the metaplectic horocycles is analyzed. This talk is an extension of the author's earlier work on the subject. [1]

MSC 2010: 11L05

Keywords: Short Gauss sums; weak invariance principle, random process, metaplectic group.

References

- [1] E. Demirci Akarsu, Short incomplete Gauss sums and rational points on metaplectic horocycles. *Int. J. Number Thr.* **10** (2014), no. 6, 1553–1576; doi: 10.1142/S1793042114500444.

A new approach to a bending energy of elastica for space curves in De-Sitter space

RIDVAN CEM DEMİRKOL¹, TALAT KÖRPİNAR², VEDAT ASİL³, SELÇUK BAŞ⁴

^{1,2,4}Mus Alparslan University, Mus, Turkey

³Fırat University, Elazığ, Turkey

emails: ¹rcdemirkol@gmail.com; ²talatkorpınar@gmail.com; ³vedatasil@gmail.com;
⁴s.bas@alparslan.edu.tr

In this paper, we firstly introduce kinematics properties of a moving particle lying in De-Sitter space S_1^3 . We assume that the particle corresponds to a different type of space curves such that they are characterized by using Frenet vector fields in De-Sitter spacetime. Based on this assumption, we present geometrical understanding of the energy on the particle in each Frenet vector fields depending on being a spacelike or timelike curve in S_1^3 . Then, we also determine the bending elastic energy functional for the same particle in S_1^3 by assuming the particle has a bending feature of elastica. Finally, we prove that bending energy formula can be represented by the energy on the particle in each Frenet vector field. We conclude our results by providing energy variation sketches with respect to time for different cases.

MSC 2010: 53C41, 53A10

Keywords: Energy, De-Sitter space, Frenet vector fields, elastica

References

- [1] J. Weber, Relativity and Gravitation. *Interscience, New York*, 1961.
- [2] J. Guven, D. M. Valencia, J. Vazquez-Montejo, Environmental bias and elastic curves on surfaces. *Phys. A: Math Theory* **47** (2014), 355201.
- [3] T. Mert, B. Karlığa, Timelike surfaces with constant angle in de -Sitter space. *Bol. Soc. Paran. Mat.* **35** (2017), 79-93.
- [4] C. M. Wood, On the energy of a unit vector field. *Geom. Dedic.* **64** (1997), 319-330.
- [5] A. Altın, On the energy and Pseduoangle of Frenet vector fields in R_n^v . *Ukrainian Mathematical J.* **63** (2011), 969-975.

An analogue of the Artin-Rees Lemma for Artinian modules

İSMAİL HAKKI DENİZLER

Van Yüzyüncü Yıl University, Van, TURKEY

email: ismailhd@yyu.edu.tr

In this study we state a useful theorem for Noetherian modules, known as the “Artin-Rees Lemma” after E. Artin and D. Rees who discovered it independently. We will show later how such an Artinian analogue can be deduced from the original result, in a direct and simple way by using Matlis duality[1].

Artin-Rees Lemma: Let R be a Noetherian ring and N a finitely generated R -module (so Noetherian). Let M be a submodule of N , and let I be a proper ideal of R . Then there exists a positive integer c such that for every $n > c$ we have

$$I^n N \cap M = I^{n-c}(I^c N \cap M).$$

(See, for example [2, (8.5)].)

Later we give the analogue result of the Artin-Rees lemma for Artinian modules;

Let A be an Artinian R -module, and let B be a submodule of A . Then for an ideal I of R there exists a non-negative integer c such that

$$B + (0 :_A I^n) = (B + (0 :_A I^c) :_A I^{n-c})$$

for all $n \geq c$, $n \in \mathbb{N}$.

We prove this lemma in two steps. First, we consider the special case in which we assume R is a complete semi-local (Noetherian) ring.

Later we complete the proof by supposing that R is an arbitrary (non-trivial commutative) ring. Since A is Artinian, then by [3], there exists a complete semi-local (Noetherian) ring R' such that the module A is, in the natural way, a faithful Artinian R' -module; moreover, a subset of A is an R -module if and only if it is an R' -submodule.

MSC 2010: 13E10, 13E05, 13A15

Keywords: Artinian modules, Noetherian modules, Ideal theory

References

- [1] E. Matlis, Injective modules over Noetherian rings. *Pacific J. Math.* **8** (1958), 511-528.
- [2] H. Matsumura, Commutative Ring Theory. *Cambridge University Press, Cambridge*, 1986.
- [3] R. Y. Sharp, Artinian modules over commutative rings. *Math. Proc. Camb. Phil. Soc.* **111** (1992), 25-33.

Kinematics of 4R and 2RPR mechanisms in Clifford algebra

VEDAT DÖRMA¹, BÜLENT KARAKAŞ², ŞENAY BAYDAŞ³

^{1,2,3}Van Yuzuncu Yil University, Van, Turkey

emails: ¹vedatdorma65@gmail.com; ²bulentkarakas@gmail.com; ³senay.baydas@gmail.com

In this paper, Clifford product, Denavit-Hartenberg representation, the forward kinematic equations and forward kinematics of motor algebra are given. Forward kinematics, motor equation and kinematic equations of the 4R and 2RPR mechanisms in Clifford algebra are obtained. Additionally, Matlab program is used for examples.

MSC 2010: 15A66, 68T40, 70B15, 53A17

Keywords: Clifford algebra, Denavit-Hartenberg representation, SCARA robot

References

- [1] E. Bayro-Corrochano and D. Kahler, Motor algebra approach for computing the kinematics of robot manipulators. *Journal of Robotic Systems* **17** (2000), no. 9, 495-516.
- [2] H. H. Hacısaliholu, Hareket Geometrisi ve Kuaterniyonlar Teorisi. *Gazi Üniversitesi, Fen-Edebiyat Fakültesi, Ankara*, 1983.
- [3] S. B. Niku, Introduction to Robotics. *Prentice Hall, New Jersey*, 2001.
- [4] J. M. McCarthy, Introduction to Theoretical Kinematics. *The MIT Press, U.S.A.*, 1990.
- [5] J. M. McCarthy, Geometric Design of Linkages. *Springer-Verlag, New York*, 2000.

A note on q -Fubini polynomials

UGUR DURAN¹, MEHMET ACIKGOZ², SERKAN ARACI³

¹Iskenderun Technical University, Hatay, Turkey

²University of Gaziantep, Gaziantep, Turkey

³Hasan Kalyoncu University, Gaziantep, Turkey

emails: ¹mtdrnugur@gmail.com; ²acikgoz@gantep.edu.tr; ³mtsrkn@hotmail.com

The main aim of this study is to introduce a new extension of Fubini polynomials based on q -numbers. Then, we investigate some of their properties including recurrence relations, differentiate properties and explicit formulas.

MSC 2010: 11B835, 05A19

Keywords: q -numbers, Fubini polynomials, Stirling numbers of the second kind

References

- [1] T. Kim, D.S. Kim and G.-W. Jang, A note on Degenerate Fubini Polynomials. *Proc. Jangjeon Math. Soc.* **20** (2017), no. 4, 521-531.

On *ve*-degrees in direct and strong products of two graphs

SÜLEYMAN EDİZ¹, MURAT CANCAN², MEHMET ŞERİF ALDEMİR³

^{1,2,3}Van Yuzuncu Yil University, Van, Turkey

emails: ¹mcancan@yyu.edu.tr; ²suleymanediz@yyu.edu.tr; ³msaldemir@yyu.edu.tr

Let G be a graph and v be a vertex of G . The *ve*-degree of the vertex v defined as the number of different edges incident to the vertices of the open neighborhood of v . In this study we investigate the *ve*-degrees in direct and strong products of two graphs.

MSC 2010: 05C07

Keywords: *ve*-degree, direct product, strong product

On the existence of periodic solutions of third order nonlinear differential equations with multiple delays

SULTAN ERDUR¹, CEMIL TUNÇ²

^{1,2}Van Yuzuncu Yil University, Van, Turkey

emails: ¹serdur82@gmail.com; ²cemtunc@yahoo.com

In this paper, we investigate the existence of periodic solutions of third order nonlinear differential equations with multiple delays by using Lyapunov's second method. We establish sufficient conditions which guarantee the existence of periodic solutions of the considered equations. We give an example to visualize. The obtained results include and improve some results in the literature.

MSC 2010: 37B25, 34C25, 34A34

Keywords: Delay differential equations, Lyapunov method, periodic solution, third order

References

- [1] E. N. Chukwu, On the boundedness and the existence of a periodic solution of some nonlinear third order delay differential equation. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* **64** (1978), no. 5, 440-447.
- [2] H. O. Tjumla and B. Tchegnani, Stability, boundedness and existence of periodic solutions of some third and fourth order nonlinear delay differential equations. *J. Nigerian Math. Soc.* **19** (2000), no. 3, 9-19.
- [3] T. Yoshizawa, Stability theory and the Existence of Periodic Solutions and Almost Periodic Solutions. *Springer-Verlag*, 1975.
- [4] Y. Zhu, On stability, boundedness and existence of periodic solution of a kind of third order nonlinear delay differential system. *Ann. of Diff. Eqs.* **8** (1992), no. 2, 249-259.

The theory of Bézier curves in \mathbb{E}^4

ESRA ERKAN¹, SALİM YÜCE²

^{1,2}Yildiz Technical University, Istanbul, Turkey

emails: ¹eserkan@yildiz.edu.tr; ²sayuce@yildiz.edu.tr

In this work, we aim to study Bézier curves that is important in Computer Aided Geometric Design and given by the equation, [2]

$$P(t) = \sum_{i=0}^n B_{i,n}(t)b_i, \quad 0 \leq t \leq 1$$

where the b_i 's represent the $n + 1$ control points in Euclidean space E^4 with regard to differential geometry. We derive Serret-Frenet elements of a Bézier curve at its all points, starting and ending points in Euclidean space E^4 .

MSC 2010: 53A04, 65D07, 68U05

Keywords: Bézier curve, Serret-Frenet frame, triple (ternary) product

Acknowledgement: This work has been supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK).

References

- [1] V. B. Anand, Computer Graphics and Geometric Modeling for Engineers. *John Wiley and Sons Inc., New York*, 1992.
- [2] A. R. Forrest, Curves and Surfaces for Computer-Aided Design. *PhD, University of Cambridge, United Kingdom, Cambridge*, 1968.
- [3] G. Farin, Curves and Surfaces For Computer Aided Geometric Design A Practical Guide. *Academic Press, San Diego*, 1996.
- [4] D. Marsh, Applied Geometry for Computer Graphics and CAD. *Springer-Verlag, London Berlin Heidelberg*, 2004.
- [5] O. Aléssio, M. Döldül, B. U. Döldül, A. S. Badr and N. H. Abdel-All, Differential geometry of non-transversal intersection curves of three parametric hypersurfaces in Euclidean 4-space. *Computer Aided Geometric Design* **31** (2014), 712-727; doi:10.1016/j.cagd.2014.09.003.
- [6] M. Z. Williams and F. M. Stein, A triple product of vectors in four-space. *Mathematics Magazine* **37** (1964), no. 4, 230-235; doi:10.2307/2688595.
- [7] R. Shaw, Vector cross products in n dimensions. *International Journal of Mathematical Education in Science and Technology* **18** (1987), no. 6, 803-816; doi:10.1080/0020739870180606.
- [8] S. A. Badr, N. H. Abdell-All, O. Alssio, M. Dldl and B. U. Dldl, Non-transversal intersection curves of hypersurfaces in Euclidean 4-space. *Journal of Computational and Applied Mathematics* **288** (2015), 81-98; doi:10.1016/j.cam.2015.03.054.
- [9] O. Aléssio, Differential geometry of intersection curves in \mathbb{R}^4 of three implicit surfaces. *Computer Aided Geometric Design* **26** (2009), no. 4, 455-471.

Misconceptions regarding representativeness in probability subject of high school students: Van case

ELİF ERTEM AKBAŞ¹, MUSTAFA GÖK²

^{1,2}Van Yuzuncu Yil University, Van, Turkey

emails: ¹elifertem@yyu.edu.tr; ²mustafagok@yyu.edu.tr

Probability is a measure of the certainty of an event. Probability, which has an important role in students' preferences in daily life, includes many objects that are difficult to understand. One of these is representativeness, which represents an exemplary situation that could possibly arise as a result of an event. For example, most people think that when tossing a coin, a sequence of six tails the TTTTTT sequence is less probable than the THHTHT sequence. Therefore, in this study, it was aimed to determine the misconceptions in the probability regarding representativeness of high school students (9th, 10th and 11th grade). In the study, the case study, one of the qualitative research methods, was used. The participant of the study constitutes 177 high school students selected by purpose of sampling method. The "Representativeness in Statistical Reasoning: Identifying and Assessing Misconceptions" test, which was developed as a data collection tool [1] and calculated with a reliability of 0.84, was adapted to Turkish. Findings of the study reveal that about half of the students are conceptual misconceptions about the representativeness of probability. It has also been found that as class levels increase, misconceptions about the representativeness of probability decrease.

MSC 2010: 97D70, 97A99

Keywords: Probability, representativeness, misconceptions, high school students

References

- [1] A. Baki, Kuramdan Uygulamaya Matematik Eğitimi. *Harf Eğitim, Ankara*, 2008.
- [2] J. Franklin, Probability theory: The logic of science. *Mathematical Intelligencer* **27** (2005), no. 2, 83-85.
- [3] R. Gürbüz, Olasılık konusunda geliştirilen materyallere dayalı öğretime ilişkin öğretmen ve öğrenci görüşleri. *Kastamonu Eğitim Dergisi* **15** (2007), no. 1, 259-270.
- [4] L. S. Hirsch and A. M. O'Donnell, Representativeness in statistical reasoning: Identifying and assessing misconceptions. *Journal of Statistics Education* **9** (2001), no. 2, 1-22; doi:10.1080/10691898.2001.11910655.
- [5] S. Bulut, Investigation of performances of prospective mathematics teachers on probability. *Hacettepe Üniversitesi Eğitim Fakültesi Dergisi* **20** (2001), 33-39.
- [6] S. Munisamy and L. Doraisamy, Levels of understanding of probability concepts among secondary school pupils. *International Journal of Mathematical Education in Science and Technology* **29** (1998), no. 1, 39-45.

New exponential stability criteria for certain neutral differential equations with interval discrete and distributed time-varying delays

MELEK GÖZEN¹, CEMİL TUNÇ²

^{1,2} Van Yuzuncu Yil University, Van, Turkey

emails: ¹melekgozen2013@gmail.com; ²cemtunc@yahoo.com

In this paper, we give new sufficient conditions for the exponential stability of solutions to certain neutral differential equations with discrete and distributed time-varying delays. Based on the some new definitions of a class of Lyapunov-Krasovskii functionals, a model transformation, the decomposition technique of constant coefficients, the Leibniz-Newton formula and usage of a zero equation, some new delay-range-dependent exponential stability criteria are derived in terms of the linear matrix inequality (LMI) for the equations considered. We give an example to illustrate the effectiveness and improvement of the results given.

MSC 2010: 34K20, 93D09, 93D20.

Keywords: Neutral differential equation, time-varying delay, exponentially stable.

References

- [1] H. Chen and X. Meng, An improved exponential stability criterion for a class of neutral delayed differential equations. *Appl. Math. Lett.* **24** (2011), no. 11, 1763-1767.
- [2] W. Chatbupapan and K. Mukdasai, New delay-range-dependent exponential stability criteria for certain neutral differential equations with interval discrete and distributed time-varying delays. *Adv. Difference Equ.* **2016** (2016), no. 324, 18 pp.
- [3] M. Gözen and C. Tunç, On exponential stability of solutions of neutral differential systems with multiple variable delays. *Electron. J. Math. Anal. Appl.* **5** (2017), no. 1, 17-31.
- [4] C. Tunç, Exponential stability to a neutral differential equation of first order with delay. *Ann. Differential Equations* **29** (2013), no. 3, 253-256.

On the kink type and singular solitons solutions to the nonlinear partial differential equation

EDA GUNAYDIN¹, YUSUF GUREFE², TOLGA AKTURK³

^{1,3} Ordu University, Ordu, Turkey

² Usak University, Usak, Turkey

emails: ¹eda-gunaydin@windowslive.com; ²yusuf.gurefe@usak.edu.tr; ³tolgaakturk@odu.edu.tr

In this study, the solutions of the Kadomtsev-Petviashvili equation were obtained by using the modified expansion function method. With this method, two and three dimensional graphics are drawn by selecting appropriate parameter values. It can be seen that the shapes of the obtained graphs correspond to the kink type and other soliton solutions graph. All the obtained solutions were checked with the help of the Wolfram Mathematica software, which provided the KP equation.

Following form the KP equation,

$$(u_t + 6uu_x + u_{xxx})_x - 3u_{yy} = 0, \quad (1)$$

MSC 2010: 35C07, 35C08, 35J60

Keywords: The modified expansion function method (MEFM), Kadomtsev-Petviashvili (KP) equation, the kink type soliton solution

References

- [1] M. Wen-Xiu and Y. Zhou, Lump solutions to nonlinear partial differential equations via Hirota bilinear forms. *Journal of Differential Equations* **264** (2018), no. 4, 2633-2659.
- [2] W. X. Ma, Lump solutions to the Kadomtsev–Petviashvili equation. *Phys. Lett. A* **379** (2015), no. 36, 1975–1978.
- [3] S. V. Manakov, V. E. Zakharov, L. A. Bordag and V. B. Matveev, Two-dimensional solitons of the Kadomtsev–Petviashvili equation and their interaction. *Phys. Lett. A* **63** (1977), 205–206.
- [4] A. A. Minzoni and N. F. Smyth, Evolution of lump solutions for the KP equation. *Wave Motion* **24** (1996), 291–305.
- [5] J. Satsuma and M. J. Ablowitz, Two-dimensional lumps in nonlinear dispersive systems. *J. Math. Phys.* **20** (1979), 1496–1503.
- [6] H. M. Baskonus, H. Bulut and A. Atangana. On the complex and hyperbolic structures of the longitudinal wave equation in a magneto-electro-elastic circular rod. *Iop Publishing* **25** (2016), no. 3, 035022.
- [7] A. Yokus, H. M. Baskonus, T. A. Sulaiman and H. Bulut, Numerical simulation and solutions of the two-component second order KdV evolutionary system. *Numerical Methods for Partial Dif. Eq.* **34** (2018), no. 1, 211-227.

Lorentzian homogeneous generalized Ricci solitons of dimension $n \geq 3$

SİNEM GÜLER^{1,2}

¹Istanbul Technical University, Istanbul, Turkey

²Istanbul Sabahattin Zaim University, Istanbul, Turkey

emails: ¹singuler@itu.edu.tr; ²sinem.guler@izu.edu.tr

In this study, we investigate the existence of Lorentzian homogeneous (generalized) Ricci solitons of dimension $n = 3$ and $n = 4$. It is known that under some conditions, three dimensional locally homogeneous Lorentzian manifolds are locally symmetric. Moreover, three dimensional locally symmetric Lorentzian manifolds which are not of constant sectional curvature are Walker manifolds, if they are not locally isometric to a Lorentzian product of a real line and a surface of constant Gauss curvature, [1, 2]. Motivated by these results, we construct several non-trivial examples of (generalized) Ricci solitons endowed with the three and four dimensional Walker metrics, [3].

MSC 2010: 53C21, 53C50, 53C25

Keywords: Ricci soliton, generalized Ricci soliton, Walker manifold

Acknowledgement: This work is supported by GAP project TGA-2018-41211 of Istanbul Technical University.

References

- [1] G. Calvaruso, Homogeneous structures on three-dimensional Lorentzian manifolds. *J. Geom. Phys.* **57** (2007), 1279–1291.
- [2] M. Brozos-Vazquez, G. Calvaruso, E. Garcia-Rio and S. Gavino-Fernandez, Three dimensional Lorentzian homogeneous Ricci solitons. *Isr. J. Math.* **188** (2012), 385–403.
- [3] M. Brozos-Vazquez, E. Garcia-Rio, P. Gilkey, S. Nikčević, and R. Vazquez-Lorenzo, The Geometry of Walker manifolds. Synthesis Lectures on Mathematics and Statistics 5, *Morgan and Claypool Publ.*, 2009.

Multi-party key exchange protocol and man in the middle attack

TUBA GÜLEŞCE TATLI¹, TURGUT HANOYMAK², ÖMER KÜSMÜŞ³

^{1,2,3}Van Yüzyüncü Yıl University, Van, Turkey

emails: ¹tubagulescetatli@gmail.com; ²turguthanoymak@gmail.com; ³omerkusmus@yyu.edu.tr

Key exchange protocols are used to generate a shared secret key between parties who want to communicate each other securely over an insecure channel. In this study, we first briefly mention about Diffie-Hellman protocol, then we generalize this to a multi-party type key exchange protocol and finally we give how an adversary can attack to this system by using the method of man in the middle attack, illustrating some concrete examples.

MSC 2010: 94A60, 94A62, 68P25

Keywords: Diffie-Hellman key exchange protocol, man in the middle attack, public key cryptography

References

- [1] W. Diffie, M. E. Hellman, New directions in Cryptography. *The Institute of Electrical and Electronics Engineers Transactions on Information Theory* **22** (1976), 644-654.
- [2] C. K. Kumar, G. J. A. Jose, C. Sajeev and C. Suyambulingom, Safety measures against man-in-the-middle attack in key exchange. *Asia Research Publishing Network (ARPN) Journal of Engineering and Applied Sciences* **7** (2006), 243-6.
- [3] M. Ahmed, B. Sanjabi, D. Aldiaz, A. Rezaei, and H. Omotunde, Diffie-Hellman and Its Application in Security Protocols. *International Journal of Engineering Science and Innovative Technology (IJESIT)* **1** (2012), 69-73.
- [4] W. Stalling, Cryptography and Network Security Principal and Practise. *Pearson Publishing*, 2006.
- [5] N. Asaithombi, A study on asymmetric cryptographic algorithms. *International Journal of Computer Science and Mobile Applications* **3** (2015), 8-13.
- [6] K. P. Arya, M. S. Aswal and V. Kumar, Comparative study of asymmetric key cryptography algorithms. *International Journal of Computer Science and Communication Networks* **5** (2011), 17-21.

Simulation studies for credibility-based multi-objective programming problems with fuzzy parameters

HANDE GÜNAY AKDEMİR

Giresun University, Giresun, Türkiye

email: hande.akdemir@giresun.edu.tr

In this study, we discuss optimal decisions for hybrid models combining fuzzy chance-constraints and expected values of objective functions. Triangular, trapezoidal and non-linear fuzzy numbers are considered in problem parameters like demands and costs. Fortunately, the concept of fuzzy variables allows us to find risk-neutral decisions via expected values based on credibility measures. Finally, numerical simulations are presented to illustrate the efficiency.

MSC 2010: 90C29, 90C70, 03E72

Keywords: Multiple objective programming, fuzzy parameters, credibility measure, chance constraints, simulation

References

- [1] M. S. Pishvaei, S. A. Torabi and J. Razmi, Credibility-based fuzzy mathematical programming model for green logistics design under uncertainty. *Computers & Industrial Engineering* **62** (2012), no. 2, 624-632.
- [2] B. Liu, Toward fuzzy optimization without mathematical ambiguity. *Fuzzy optimization and decision making* **1** (2002), no. 1, 43-63.
- [3] X. Zhou, Y. Tu, J. Han, J. Xu and X. Ye, A Class of Level-2 Fuzzy Decision-Making Model with Expected Objectives and Chance Constraints: Application to Supply Chain Network Design. *International Journal of Information Technology Decision Making* **16** (2017), no. 4, 907-938.
- [4] Y. Cheng, J. Peng, Z. Zhou, X. Gu and W. Liu, A Hybrid DEA-Adaboost Model in Supplier Selection for Fuzzy Variable and Multiple Objectives. *IFAC-PapersOnLine* **50** (2017), no. 1, 12255-12260.

Regularization of inverse coefficient determination problem in a hyperbolic problem

HAKKI GÜNGÖR

Ufuk University, Ankara, Turkey

email: hakki.gungor@ufuk.edu.tr

In this study, an inverse tension determination problem in a wave equation is investigated. Due to ill-posedness, a regularization process is carried out. After proving the existence and uniqueness of the solution, characterization of the solution is presented. The outcomes have also been tested with numerical examples.

MSC 2010: 65M32, 35L05

Keywords: Inverse Problem, adjoint method, wave equation

References

- [1] O. A. Ladyzhenskaya, *Boundary Value Problems in Mathematical Physics. Springer-Verlag, New York*, 1985.
- [2] G. F. Kuliev, Problem of Control with Control Functions at the Senior Derivatives and in the Right Sides of the Equation with Functional Constraints. *Tr. Azerb. Mat. O-va* **2** (1996), 122–140.
- [3] M. Subaşı, A. Kaçır, A variational technique for optimal boundary control in a hyperbolic problem. *Applied Mathematics and Computation* **218** (2012), 6629–6636.
- [4] R. K. Tagiyev, Correctness and Regularization of a Class of Problems of Optimal Control of the Coefficients of Linear Hyperbolic Equation. *Azerb. Kirov Gos. Univ, Baku*, (1984), 98-105.
- [5] G. F. Kuliev, Problem of Optimal Control of the Coefficients for Hyperbolic Equations. *Izv. Vyssh. Uchebn. faved., Mat.* **3** (1985), 39–44.
- [6] R. K. Tagiyev, On the Optimal Control Problem by Coefficients of the Hyperbolic Equation. *Trans.NAS Azerbaijan, Isc. Math.–Mech.* **21** (2001), no. 4, 230–235.
- [7] X. Feng, B. Sutton, S. Leuhart, et au., Identification Problem for the Wake Equation with Neumann Data Input and Dirichlet Data Abserving. *Nonlinear Anal.* **52** (2003), no. 7, 1777–1795.
- [8] R. K. Tagiyev, On Optimal Control of the Hyperbolic Equateona Coefficients. *Automotion snd Remoti Contral*
- [9] H. W. Engl, M. Hanke, A. Neubauer, Regularization of Inverse Problems. *Kluwer Academic Publishers, The Netherlands*, 1996.
- [10] M. Goebel, On existence of optimal control. *Mathematische Nachrichten* **93** (1979), 67-73.

Approximation by summation-integral type operators involving Brenke polynomials

ŞULE YÜKSEL GÜNGÖR

Gazi University, Ankara, Turkey

email: sulegungor@gazi.edu.tr

In this study, we introduce a sequence of summation-integral type operators linking generalized Brenke-Szász type and general Szász basis functions. A local and direct approximation theorem by means of Ditzian-Totik modulus of smoothness are obtained. The rate of convergence in terms of the Lipschitz class and the Lipschitz type maximal function is investigated.

MSC 2010: 41A25, 41A35, 41A36

Keywords: Brenke polynomials, Szász operators, Lipschitz class, Lipschitz type maximal function, degree of approximation

References

- [1] Ç. Atakut and I. Büyükyazıcı, Approximation by Kantorovich-Szász type operators based on Brenke type polynomials. *Numer. Funct. Anal. Optim.* **37** (2016), no. 12, 1488-1502.
- [2] T. Garg, P. N. Agrawal and S. Arici, Rate of convergence by Kantorovich-Szász type operators based on Brenke type polynomials. *Journal of Inequalities and Applications* **156** (2017); doi:10.1186/s13660-017-1430-z.
- [3] M. E. H. Ismail, Classical and Quantum Orthogonal Polynomials in One Variable. *Cambridge University Press, Cambridge*, 2005.
- [4] M. Mursaleen and K. J. Ansari, On Chlodowsky variant of Szász operators by Brenke type polynomials. *Applied Math. Comput.* **271** (2015), 991-1003.
- [5] S. Varma, S. Sucu and G. Icoz, Generalization of Szász operators involving Brenke type polynomials. *Comput. Math. Appl.* **64** (2012), no. 2, 121-127.
- [6] S. Varma and F. Tasdelen, On a generalization of Szász-Durrmeyer operators with some orthogonal polynomials. *Stud. Univ. Babeş-Bolyai Math.* **58** (2013), no. 2, 225-232.

Some notes on the order-to-topology continuous operators

KAZEM HAGHNEJAD AZAR

University of Mohaghegh Ardabili, Ardabil, Iran

email: haghnejad@uma.ac.ir

Let E be a Riesz space and F be a vector topology with topology τ . An operator T from E into F is said to be order-to-topology continuous whenever $x_\alpha \xrightarrow{o} 0$ implies $Tx_\alpha \xrightarrow{\tau} 0$ for each $(x_\alpha)_\alpha \subset E$. For each sequence $(x_n) \subset E$, if $x_n \xrightarrow{o} 0$ implies $Tx_n \xrightarrow{\tau} 0$, then T is called σ -order-to-topology continuous operator. The collection of all order-to-topology continuous operators will be denoted by $L_{\sigma\tau}(E, F)$; the subscript $\sigma\tau$ is justified by the fact that the order-to-topology continuous operators, that is,

$$L_{\sigma\tau}(E, F) = \{T \in L(E, F) : T \text{ is order-to-topology continuous} \}.$$

Similarly, $L_{\sigma\tau}^\sigma(E, F)$ will be denote the collection of all σ -order-to-topology continuous operators, that is,

$$L_{\sigma\tau}^\sigma(E, F) = \{T \in L(E, F) : T \text{ is } \sigma - \text{order-to-topology continuous} \}.$$

For a normed space F , we write $L_{on}(E, F)$ and $L_{ow}(E, F)$ for collection of order-to-norm topology continuous operators and order-to-weak topology continuous operators, respectively. $L_{on}^\sigma(E, F)$ and $L_{ow}^\sigma(E, F)$ have similar definitions. Let E be a σ -Dedekind complete Riesz space and F be a normed Riesz space. If T is interval-bounded, then $T \in L_{on}^\sigma(E, F)$ if and only if T is order weakly compact. Let E be a Riesz space and let F a normed Riesz space with order unit. Then $L_{on}(E, F)$ is a band in $L_b(E, F)$. In this paper, we will study some properties of this new classification of operators. We will investigate the relationships between order-to-topology continuous operators with order continuous, order weakly compact and b -weakly compact operators, see [2, 3]

MSC 2010: 46B42, 47B60

Keywords: Riesz space, order-to-topology continuous, b -weakly compact operator.

References

- [1] C. D. Aliprantis and O. Burkinshaw, Positive Operators. *Springer, Berlin*, 2006.
- [2] C. D. Aliprantis and O. Burkinshaw, Locally Solid Riesz Spaces. *Springer, Berlin*, 1978.
- [3] S. Alpay and B. Altin, C. Tonyali, On property (b) of vector lattices. *Positivity* **7** (2003), 135–139.
- [4] S. Alpay and B. Altin, A note on b -weakly compact operators. *Positivity* **11** (2007), no. 4, 575–582.
- [5] P. Meyer-Nieberg, Banach Lattices. *Universitext. Springer, Berlin*, 1991.

On mathematical aspects of blockchain architecture

TURGUT HANOYMAK¹, ATILLA BEKTAŞ²

¹Van Yüzüncü Yıl University, Van, Turkey

²Middle East Technical University, Ankara, Turkey

emails: ¹turguthanoymak@gmail.com; ²bektasatilla@gmail.com

Blockchain technology is a distributed database that enables transfer of assets we value beyond the transfer of data made in many places such as multimedia, communication, web interface in today's internet world. In its broadest terms, blockchain is the delivery of central trust over the Internet, allowing the removal of a central server or trusted authority. In this study, we firstly give an overview of blockchain architecture and then present the mathematics behind blockchain such as finite fields, digital signature algorithms, hashing algorithms, public key cryptography, and so on. Furthermore, technical challenges and recent advances are briefly listed. We also summarize possible future trends for blockchain.

MSC 2010: 94A60, 94A62, 68P25

Keywords: Blockchain, bitcoin, digital signature, public key cryptography

References

- [1] S. Nakamoto, Bitcoin: A Peer-to-Peer Electronic Cash System. <https://bitcoin.org/bitcoin.pdf>, (Accessed on 20.07.2018), 2009.
- [2] A. Çarkacıoğlu, Kripto-para Bitcoin. *Research Report, Capital Markets Board of Turkey, Research Department*, 2016.
- [3] A. M. Antonopoulos, Mastering Bitcoin. *O'Reilly Media, Inc.*, 2014.
- [4] Prypto, Bitcoin for Dummies. *A Wiley Brand. John Wiley & Sons, Inc.*, 2016.
- [5] E. Rykwaldner, The Math Behind Bitcoin. <https://www.coindesk.com/math-behind-bitcoin>, (Accessed on 20.07.2018), October, 2014.

A new class of set-valued contractions and related results

HÜSEYİN IŞIK

Muş Alparslan University, Muş, Turkey

email: isikhuseyin76@gmail.com

The aim of this study to investigate the existence of solutions for nonlocal integral boundary value problem of Caputo type fractional differential inclusions. To achieve this goal, we take advantage of fixed point theorems for multivalued mappings satisfying a new class of contractive conditions in the setting of complete metric spaces. We derive new fixed point results which extend and improve the results in [1, 2, 3] and others by means of this new class of contractions. We also supply some examples to support the new theory.

MSC 2010: 34A08, 34A60, 47H10

Keywords: Multivalued maps, fixed points, fractional differential inclusions, nonlocal boundary conditions

References

- [1] M. Jleli, B. Samet, A new generalization of the Banach contraction principle. *J. Inequal. Appl.* **2014** (2014), no. 38; doi:10.1186/1029-242X-2014-38.
- [2] S. B. Nadler, Multi-valued contraction mappings. *Pacific J. Math.* **30** (1969), 475-488.
- [3] F. Vetro, A generalization of Nadler fixed point theorem. *Carpathian J. Math.* **31** (2015), no. 3, 403-410.

On numerical solution of an optimal control problem involving hyperbolic equation

SEDA İĞRET ARAZ

Siirt University, Siirt, Turkey

email: sedaaraz@siirt.edu.tr

In this paper, we presents a numerical algorithm for solving a class of optimal control problems with hyperbolic equation. We show that the optimal solution is exist and unique in a regular space. After obtaining adjoint problem and calculating derivative of the cost functional, numerical approximations are obtained via Gradient Method. Computational results demonstrate that the proposed method is able to generate good numerical approximations for optimal control problems.

MSC 2010: 49J20, 35L20, 49J50

Keywords: Optimal control, hyperbolic equations, Frechet differentiability

References

- [1] T. Yeloğlu and M. Subaşı, Simultaneous control of the source terms in a vibrational string problem. *Iranian Journal of Science & Technology, Transaction A* **34** (2010), No. A1.
- [2] R. K. Tagiyev, On optimal control of the hyperbolic equation coefficients. *Automation and Remote Control* (2012), 1145-1155.
- [3] G. M. Bahaa, Boundary control problem of infinite order distributed hyperbolic systems involving time lags. *Intelligent Control and Automation* **3** (2012), 211-221.
- [4] A. Hasanov, Simultaneous determination of the source terms in a linear hyperbolic problem from the final over determination: weak solution approach. *Journal of Applied Mathematics* (2009), 1-19.
- [5] O. A. Ladyzhenskaya, Boundary Value Problems in Mathematical Physics. *Springer-Verlag, New York*, 1985.
- [6] M. Goebel, On existence of optimal control. *Math. Nachr.* **93** (1979), 67-73.
- [7] K. Yosida, Functional Analysis. *Springer-Verlag, New York*, 1980.
- [8] F. P. Vasilyev, Ekstremal Problemlerin Çözüm Metotları. *Nauka*, 1981.

Conditional expectation operators on measurable function spaces

MOHAMMAD REZA JABBARZADEH

University of Tabriz, Tabriz, Iran

email: mjabbar@tabrizu.ac.ir

In this note, we discuss matrix theoretic characterizations for weighted conditional type operators in some operator classes on $L^2(\Sigma)$ such as, self-adjoint, normal, quasinormal and positive operator classes. Also, we prove some basic results on the Moore-Penrose inverse and the Aluthge transformation of these type operators.

MSC 2010: 47G30 , 47B20, 47B38

Keywords: Conditional expectation, Moore-Penrose inverse, Aluthge transformation

References

- [1] Y. Estaremi and M. R. Jabbarzadeh, Weighted Lambert type operators on L^p spaces. *Oper. Matrices* **7** (2013), 101-116.
- [2] Y. Estaremi and M. R. Jabbarzadeh, Weighted composition Lambert-type operators on L^p spaces. *Mediterr. J. Math.* **11** (2014), 955-964.
- [3] J. Herron, Weighted conditional expectation operators. *Oper. Matrices* **5** (2011), 107-118.
- [4] M. R. Jabbarzadeh and M. Jafari Bakhshkandi, Centered operators via Moore-Penrose inverse and Aluthge transformations. *Filomat* **31** (2017), 6441-6448.
- [5] M. R. Jabbarzadeh and M. Sohrabi Chegeni, Moore-Penrose inverse of conditional type operators. *Oper. Matrices* **11** (2017), 289-299.
- [6] M. M. Rao, Conditional measure and applications. *Marcel Dekker, New York*, 1993.

Inverse kinematics computation for a 6-DOF articulated robot arm using conformal geometric algebra

BAHAR KALKAN¹, ŞENAY BAYDAŞ², BÜLENT KARAKAŞ³

^{1,2,3}Van Yuzuncu Yil University, Van, Turkey

emails: ¹baharkalkan1@gmail.com; ²senay.baydas@gmail.com; ³bulentkarakas@gmail.com

Geometric algebra provides a powerful computational framework for geometric applications in many areas including robotics. In fact, geometric algebra enables us to express fundamental robotics physics in a language that is free from coordinates or indices. The geometric algebra framework gives many equations a degree of clarity that is definitively lost in matrix algebra or tensor algebra. Geometric algebra represents orthogonal transformations more efficiently than the orthogonal matrices by reducing the number of coefficients. The rotation property can be applied to all objects in geometric algebra while it can be applied only on vectors in the quaternion algebra. Conformal geometric algebra is geometric algebra expanded to five dimensions from three dimensions. In conformal geometric algebra, in addition to the three-dimensional space, two more basic vectors are added, representing the location of the origin and the infinity. \mathbb{R}^n is extended with unit vectors e_+ and e_- and $\mathbb{G}^{n+1,1}$ is constructed by defining $e_+^2 = 1$ and $e_-^2 = -1$. Define the origin $o = \frac{e_- - e_+}{2}$ and infinity $\infty = e_- + e_+$. These are null vectors:

$$o^2 = \infty^2 = 0.$$

In this algebra straight lines, planes, circles and spheres can be defined as vectors and rotations and translations can be defined by the rotor. These rotors can be applied to any object. In this paper, a new algorithm for the forward displacement analysis of an Articulated robot arm with six degrees of freedom mechanism based on geometric algebra (GA) will be presented. This paper describes a novel method for solving the inverse kinematics of this robot arm using conformal geometric algebra and proposes a geometric algebra (GA) based approach to carry out kinematics of given mechanism.

MSC 2010: 15A66, 53A17

Keywords: Geometric algebra (GA), inverse kinematics, robotics, conformal geometric algebra (CGA), Clifford algebra

On some vector valued multiplier spaces obtained by Zweir matrix method

RAMAZAN KAMA

Siirt University, Siirt, Turkey

emails: ramazankama@siirt.edu.tr

In this study, by using the Zweir matrix and a sequence of continuous linear operators, we introduce some vector valued multiplier spaces and summing operators associate with this spaces, respectively and study a series of some properties of them.

MSC 2010: 46B15, 40A05, 46B45

Keywords: Vector valued multiplier space, Zweir matrix, Summing operator

References

- [1] A. Aizpuru and F. J. Pérez-Fernández, Characterizations of series in Banach spaces. *Math. Univ. Comenian.* **58** (1999), no. 2, 337–344.
- [2] B. Altay and R. Kama, On Cesàro summability of vector valued multiplier spaces and operator valued series. *Positivity* **22** (2018), 575-586.
- [3] C. Bessaga and A. Pelczynski, On bases and unconditional convergence of series in Banach spaces. *Stud. Math.* **17** (1958), 151-164.
- [4] C. Swartz, Operator valued series and vector valued multiplier spaces. *Casp. J. Math. Sci.* **3** (2014), no. 2, 277-288.
- [5] F. Başar, Summability Theory and Its Applications. *Bentham Science Publishers, İstanbul*, 2012.
- [6] F. Başar and B. Altay, On the space of sequences of p -bounded variation and related matrix mappings, (English, Ukrainian summary) *Ukrain. Mat. Zh.* **55** (2003), no. 1, 108-118; reprinted in *Ukrainian Math. J.* **55** (2003), no. 1, 136-147.
- [7] M. Şengönül, *On the Zweier sequence space*, *Demonstratio Mathematica* **40** (2007), no. 1, 181-196.

Λ – matrix as a summability operator and completeness of certain normed spaces via weakly unconditionally Cauchy series

MAHMUT KARAKUŞ¹, TUNAY BİLGİN²

^{1,2} Van Yuzuncu Yil University, Van, Turkey

emails: ¹mkarakus@yyu.edu.tr; ²tbilgin@yyu.edu.tr

Let $\sum_i x_i$ be a series in a real normed space. Then the series is said to be weakly unconditionally Cauchy if the sequence $((\sum_{i=1}^n x_{\pi(i)})_n)$ is weakly Cauchy sequence, for every permutation π of \mathbb{N} , the set of positive integers X [1]. Some characterizations of weakly unconditionally Cauchy series are well-known facts. For example, the series $\sum_n x_n$ in a Banach space is weakly unconditionally Cauchy iff it is a c_0 - multiplier convergent series [2].

In this study we interest some new characterizations of weakly unconditionally Cauchy series by completeness of certain normed spaces which are obtained from summability matrix $\Lambda = (\lambda_{nk})$ of Móricz [3] defined by

$$\lambda_{nk} = \begin{cases} \frac{\lambda_k - \lambda_{k-1}}{\lambda_n} & , \quad (1 \leq k \leq n) \\ 0 & , \quad (k > n) \end{cases} . \quad (1)$$

Here, $\lambda = (\lambda_k)$ is a strictly increasing sequence of positive reals tending to infinity, i.e., $0 < \lambda_1 < \lambda_2 < \dots$ and $\lim_k \lambda_k = \infty$.

MSC 2010: 46B15, 46A45, 40A05

Keywords: Weakly unconditionally Cauchy series, matrix summability, completeness of normed spaces

References

- [1] C. Swartz, Multiplier Convergent Series. *World Scientific Publishing, Singapore*, 2009.
- [2] J. Diestel, Sequences and Series in Banach spaces. *Springer-Verlag, New York*, 1984.
- [3] F. Móricz, On Λ – strong convergence of numerical sequences and fourier series. *Acta Math. Hung.* **54** (1989), no. 3-4, 319-327.
- [4] B. Altay and R. Kama, On Cesàro summability of vector valued multiplier spaces and operator valued series. *Positivity* **22** (2018), no. 2, 575-586.
- [5] A. Aizpuru, A. Gutiérrez-Dávila and A. Sala, Unconditionally Cauchy series and Cesàro summability *J. Math. Anal. Appl.* **324** (2006) 39-48.
- [6] F. Başar, Summability Theory and Its Applications. *Bentham Science Publishers, İstanbul*, 2012.

The maximal function in Sobolev spaces

YASİN KAYA

Dicle University, Diyarbakır, Turkey

email: ykaya@dicle.edu.tr

The theory of Hardy-Littlewood maximal function and Sobolev spaces one of the most important topic in analysis

In this presentation, I give an overview of the development of the maximal function in Sobolev spaces and show a result in this frame.

MSC 2010: 46E35, 47G10, 26A42

Keywords: Maximal function, Sobolev spaces, sublinearity

A novel Lyapunov type inequality for quasilinear impulsive systems

ZEYNEP KAYAR

Van Yuzuncu Yil University, Van, Turkey

email: zeynepkayar@yyu.edu.tr

We establish Lyapunov-type inequality and for Dirichlet problem associated with the quasilinear impulsive system involving the (p, q) -Laplacian operator. This inequality is used to obtain disconjugacy criterion and to find lower bounds for eigenvalues associated to related eigenvalue problems. Our results not only improve the recent related results and that of [1, 2] but also generalize them to the impulsive case.

MSC 2010: 34A37, 34C10, 34A34

Keywords: Lyapunov type inequality, impulse, quasilinear systems

References

- [1] M. Jleli and B. Samet, On Lyapunov-type inequalities for (p, q) -Laplacian systems. *Journal of Inequalities and Applications* **2017** (2017), no. 100, 1-9.
- [2] Z. Kayar, Lyapunov type inequalities and their applications for linear and nonlinear systems under impulse effect. *Ph.D Thesis, METU, Ankara*, 2014.

Total dominator coloring of a graph

ADEL P. KAZEMI

University of Mohaghegh Ardabili, Ardabil, Iran

email: adelpkazemi@yahoo.com

A total dominator coloring of a graph G is a proper coloring of G in which each vertex of the graph is adjacent to every vertex of some color class. The total dominator chromatic number of a graph is the minimum number of color classes in a total dominator coloring of it. Here, we talk on the total dominator coloring of a graph by giving some tight bounds for the total dominator chromatic number of a graph, a tree, join of two graphs and Nordhaus-Gaddum-like relations.

MSC 2010: 05C15, 05C69

Keywords: Total dominator coloring, Total dominator chromatic number, total domination number, central graph, Nordhaus-Gaddum relation

References

- [1] N. Ghanbari and S. Alikhani, More on the total dominator chromatic number of a graph. *ArXiv:1705.10231*.
- [2] M. A. Henning, Total dominator colorings and total domination in graphs. *Graphs and Combinatorics* **31** (2015), 953–974.
- [3] P. Jalilolghadr, A. P. Kazemi and A. Khodkar, Total dominator coloring of the circulant graphs $C_n(a, b)$. *manuscript*.
- [4] A. P. Kazemi, Total dominator chromatic number of a graph. *Transactions on Combinatorics* **4** (2015), no. 2, 57–68.
- [5] A. P. Kazemi, Total dominator coloring in product graphs. *Utilitas Mathematica* **94** (2014), 329–345.
- [6] A. P. Kazemi, Total dominator chromatic number of Mycielskian graphs. *Utilitas Mathematica* **103** (2017), 129–137.
- [7] A. P. Kazemi and F. Kazemnejad, Total dominator coloring of central graphs. *submitted*.

Invariant submanifolds of statistical Kenmotsu manifolds and their curvatures

MOHAMMAD BAGHER KAZEMI BALGESHIR

University of Zanjan, Zanjan, Iran

email: mbkazemi@znu.ac.ir

A Riemannian manifold (\bar{M}, g) with an affine and torsion free connection $\bar{\nabla}$ satisfying

$$\bar{\nabla}_V g(U, W) = \bar{\nabla}_U g(V, W) \quad \forall U, V, W \in \mathcal{T}(\bar{M}), \quad (1)$$

is called a statistical manifold [1, 2]. Moreover, there is an affine connection $\bar{\nabla}^*$ on \bar{M} which is called the dual connection of $\bar{\nabla}$ with respect to the g , such that

$$Ug(V, W) = g(\bar{\nabla}_U V, W) + g(V, \bar{\nabla}_U^* W). \quad (2)$$

In this paper, we study statistical manifolds which admit an almost contact and Kenmotsu structure [3]. We investigate the shape operator of invariant submanifolds of statistical Kenmotsu manifolds and prove it vanishes on these submanifolds if the the structure vector field is tangent to the submanifold.

MSC 2010: 53C15, 53C40, 60D05

Keywords: Kenmostu manifold, statistical structure, invariant submanifold

References

- [1] S. Amari and H. Nagaoka, Methods of information geometry. *Transl. Math. Monogr. 191.*, Amer. Math. Soc., 2000.
- [2] H. Furuhata, I. Hasegawa, Y. Okuyama, K. Sato and M. Shahid, Sasakian statistical manifolds. *J. Geom. Phys.* **117** (2017), 179–186.
- [3] K. Yano and M. Kon, Structures on manifolds. *World Scientific*, 1984.

Reduction of Navier-Stokes equation to a linear equation

WALEED S. KHEDR

Cairo, Egypt

email: waleedshawki@yahoo.com

In this article, we provide two theorems on pointwise coincidence between solutions of Navier-Stokes equation and solutions of standard linear second order parabolic equations with the same data. We show that the convection, the pressure, and the external forces (if applied) are governed by some sort of balance independent of the equation that governs the solution itself. In light of the well establishment of the theory of existence, regularity and uniqueness of linear second order parabolic equations, this result represents an important step to fully understand the qualitative properties of the solutions to Navier-Stokes equation. **Details:** The initial profile is \mathbf{v}_0 and the boundary conditions (in case of bounded domains) are denoted by \mathbf{v}^* . The model equation in hand is:

$$\begin{cases} \mathbf{v}_t + (\mathbf{v} \cdot \nabla)\mathbf{v} - \mu\Delta\mathbf{v} = -\nabla p + \mathbf{f}, & \nabla \cdot \mathbf{v} = 0 \quad \text{in } \Omega_t, \\ \mathbf{v}(\mathbf{x}, 0) = \mathbf{v}_0(\mathbf{x}) & \text{in } \Omega_0, \\ \mathbf{v}(\mathbf{x}, t) = \mathbf{v}^*(\mathbf{x}_{n-1}, t) & \text{on } \partial\Omega_t, \end{cases} \quad (1)$$

where Ω_t and $\partial\Omega_t$ denotes the fixed boundary if the domain is bounded. The solution \mathbf{v} is the vector field representing the velocity of the flow in each direction, and its rotation $\omega = \nabla \times \mathbf{v}$ is the vorticity. Note that $\nabla \cdot \omega = 0$ in $\bar{\Omega}_t$ by compatibility. The solution is investigated in light of the most common classical definition of weak solutions in order to ensure the possibility of generalizing the subsequent results to the widest classes of possible solutions. In particular, we follow the definition of weak solutions introduced first by Leray [1, 2, 3]. Further, the investigation in this article establishes rigorously the previously investigated results in [4, 5].

MSC 2010: 76D03, 76D05, 76M30, 76R10

Keywords: Fluid Mechanics, Navier-Stokes equation, convection

Acknowledgement: This is an extended abstract of the article submitted to arXiv on 24th of January 2018 with submission number arXiv:submit/2142016.

References

- [1] J. Leray, Étude de diverses équations intégrales non linéaires et de quelques problèmes que pose l'hydrodynamique. *J. Math. Pures Appl.* **12** (1933), 1-82.
- [2] J. Leray, Essai sur les mouvements plans d'un liquide visqueux que limitent des parois. *J. Math. Pures Appl.* **13** (1934), 341-418.
- [3] J. Leray, Essai sur le mouvement d'un liquide visqueux emplissant l'espace. *Acta Math.* **63** (1934), 193-248.
- [4] W. S. Khedr, Classical fundamental unique solution for the incompressible Navier-Stokes equation in R^N . *Jamp* **5** (2017), 939-952.
- [5] W. S. Khedr, Nonconvection and uniqueness in Navier-Stokes equation. *arXiv:1706.02552* (2017).

A note on the annihilator of certain local cohomology modules

AHMAD KHOJALI

University of Mohaghegh Ardabili, Ardabil, Iran

email: Khojali@uma.ac.ir

Let R be a commutative Noetherian ring of dimension d and \mathfrak{a} an ideal of R . It is easily seen that $\frac{R}{\text{Ann}_R(H_{\mathfrak{a}}^{d-1}(R))} \hookrightarrow \text{Hom}_R(H_{\mathfrak{a}}^{d-1}(R), H_{\mathfrak{a}}^{d-1}(R))$. Vanishing of $\text{Ann}_R(H_{\mathfrak{a}}^{d-1}(R))$ is interesting. In this direction, we consider the question whether non-vanishing of $H_{\mathfrak{a}}^{d-1}(R)$ is equivalent to the vanishing of its annihilator. In the case (R, \mathfrak{m}) is a regular local ring containing a field and $H_{\mathfrak{a}}^i(R) \neq 0$ for a given integer i , then in characteristic zero Lyubeznik [5] and in characteristic $p > 0$ Huneke and Koh [3] showed that $\text{Ann}_R H_{\mathfrak{a}}^i(R) = 0$. Here are some attempts to compute $\text{Ann}_R H_{\mathfrak{a}}^i(R)$. (1) If R is a local complete ring R with $H_{\mathfrak{a}}^i(R) = 0$ for every $i \neq \text{ht}(\mathfrak{a})$, then $\text{Ann}_R(H_{\mathfrak{a}}^{\text{ht}(\mathfrak{a})}(R)) = 0$ (see [2]). (2) If R is a complete Gorenstein local domain with some mild assumptions $\text{Ann}_R(H_{\mathfrak{a}}^i(R)) = 0$, where $i = \text{grade}(\mathfrak{a}, R)$ (see [6]). (3) If R is complete and local, then $\text{Ann}_R H_{\mathfrak{a}}^d(R) = \cap \mathfrak{q}$, that \mathfrak{q} are primary components of (0) with $\dim \frac{R}{\mathfrak{q}} = \dim R$ and $\text{rad}(\mathfrak{a} + \mathfrak{q}) = \mathfrak{m}$ (see [4]).

In this note we consider the annihilator of some certain local cohomology modules and some vanishing results of these modules will be considered. In particular, it is proved that: (1) If (R, \mathfrak{m}) be a local domain such that $H_{\mathfrak{a}}^d(R) = 0$ and $H_{\mathfrak{a}}^{d-1}(R)$ is not Artinian, then $\text{Ann}_R H_{\mathfrak{a}}^{d-1}(R) = 0$. (2) Let (R, \mathfrak{m}) be a Cohen-Macaulay local ring and \mathfrak{a} a 1-dimensional ideal. Then, $H_{\mathfrak{a}}^{d-1}(R)$ is not Artinian.

MSC 2010: 13D45

Keywords: Annihilator of local cohomology, non-Artinian local cohomology, Buchsbaum type modules

Acknowledgement: The author would like to thank the University of Mohaghegh Ardabili for their support.

References

- [1] M. Brodmann and R.Y. Sharp, An Algebraic Introduction with Geometric Applications. Second edition. *Cambridge University Press, Cambridge*, 2013.
- [2] M. Hellus and J. Stückrad, On endomorphism rings of local cohomology modules. *Proc. of AMS* **136** (2008), no. 7, 2333–2341.
- [3] C. Huneke and J. Koh, Cofiniteness and vanishing of local cohomology modules. *Math. Proc. Camb. Phil. Soc.* **110** (1991), 421–429.
- [4] L. Lynch, Annihilators of top local cohomology. *Comm. Alg.* **40** (2012), 542–551.
- [5] G. Lyubeznik, Finiteness propositionerties of local cohomology modules (an application of D-modules to Commutative Algebra). *Invent. math.* **113** (1993), 41–55.
- [6] W. Mahmood and P. Schenzel, On invariants and endomorphism rings of certain local cohomology modules. *J. Algebra* **372** (2012), no. 2, 56–67.

Sufficient conditions for global asymptotic stability of neural networks with time-varying delays

ERDAL KORKMAZ¹, CEMİL TUNC²

¹ Mus Alparslan University, Mus, Turkey

²Van Yuzuncu Yil University, Van, Turkey

emails: ¹korkmazerdal36@hotmail.com ; ²cemtunc@yahoo.com;

In this paper, for global asymptotic stability of the equilibrium point of neural networks with delays is obtained some new sufficient conditions by using Lyapunov technique. The results obtained have shown to improve the previous results derived in the literature . The results are supported by a few examples.

MSC 2010: 34B18, 34B25

Keywords: Lyapunov functionals, global asymptotic stability, delay differential equations.

References

- [1] S. Arik, An analysis of global asymptotic stability of delayed cellular neural networks. *IEEE Trans. Neural Netw.* **13** (2002), 1239-1242.
- [2] S. Arik, On the global asymptotic stability of delayed cellular neural networks. *IEEE Trans. Circuits Syst. I* **47** (2000), 571-574.
- [3] Y. Guo, Global asymptotic stability analysis for integro-differential systems modeling neural networks with delays. *Zeitschrift fr angewandte Mathematik und Physik* **61** (2010), no. 6, 971-978.
- [4] Q. Zhang, X. Wei and J. Xu, Global asymptotic stability of cellular neural networks with infinite delay. *Neural Netw. World* **15** (2005), 579-589.

On the cubic nonlinear Shrodinger's equation with repulsive delta potential

ZELİHA KÖRPİNAR¹, FATİH COŞKUN²

^{1,2}Muş Alparslan University, Muş, Turkey

emails: ¹zelihakorpinar@gmail.com; ²fatihcoskun323@gmail.com

In this work, is introduced to obtain approximate solutions of the cubic nonlinear Shrödinger's equation (NLSE) with repulsive delta potential subject to certain initial conditions by using Residual power series method (RPSM). The consequens show that this method is efficient and convenient and can be applied to a large sort of problems. The approximate solutions are compared with the known exact solutions.

MSC 2010: 35L05, 58Z05

Keywords: Residual power series method, cubic nonlinear Shrödinger's equation, potential

References

- [1] J. H. He, Approximate analytical solution for seepage flow with fractional derivatives in porous media. *Comput. Meth. Appl. Mech. Eng.* **167** (1998), 57–68.
- [2] O. Abu Arqub, Series solution of fuzzy differential equations under strongly generalized differentiability. *Journal of Advanced Research in Applied Mathematics* **5** (2013), 31-52.
- [3] O. Abu Arqub, A. El-Ajou and S. Momani, Constructing and predicting solitary pattern solutions for nonlinear time-fractional dispersive partial differential equations. *Journal of Computational Physics* **293** (2015), 385-399.
- [4] D. Baleanu, M. İnc, A. I. Aliyu and A. Yusuf, Optical solitons, nonlinear self-adjointness and conservation laws fort he cubic nonlinear Shrödingers equation with repulsive delta potential. *Superlattices and Microstructures* **111** (2017), 546-555.

On numerical solutions for fractional (1+1)-dimensional Biswas-Milovic equation

ZELİHA KÖRPİNAR¹, MUSTAFA İNÇ²

¹Muş Alparslan University, Muş, Turkey

²Fırat University, Elazığ, Turkey

emails: ¹zelihakorpinar@gmail.com; ²minc@firat.edu.tr

In this work, numerical solutions are obtained for fractional (1+1)-dimensional Biswas-Milovic equation that defines the long-space optical communications by using the residual power series method (RPSM). The RPSM gets Maclaurin expansion of the solution. The solutions of present equation are computed in the shape of quickly convergent series with quickly calculable fundamentals by using mathematica software package. Explanation of the method is given graphical consequens and series solutions are made use of to represent our solution. The found consequens show that technique is a power and efficient method in conviction of solution for the fractional (1+1)-dimensional Biswas-Milovic equation.

MSC 2000: 35L05, 58Z05

Keywords: Residual power series method, (1+1)-dimensional Biswas-Milovic equation, Series solution.

References

- [1] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, Theory and Applications of Fractional Differential Equations. *Elsevier, Amsterdam*, 2006.
- [2] A. Biswas, M. Mirzazadeh, M. Savescu, D. Milovic, K. R. Khan, M. F. Mahmood and M. Belic, Singular solitons in optical metamaterials by ansatz method and simplest equation approach. *J. Modern Opt.* **61** (2014), 1550–1555.
- [3] Q. Zhou, Optical solitons in the parabolic law media with high-order dispersion. *Optik* **125** (2014), 5432–5435.
- [4] A. Biswas, H. Triki, Q. Zhoud, S. P. Moshokoa, M. Z. Ullah and M. Belic, Cubic-quartic optical solitons in Kerr and power law media. *Optik* **144** (2017), 357–362.

On inextensible flow with Schrödinger flow

ZELİHA KÖRPİNAR¹, TALAT KÖRPİNAR², SELÇUK BAŞ³, M. TALAT SARIAYDIN⁴

^{1,2,3}Muş Alparslan University, Muş, Turkey

⁴Selçuk University, Konya, Turkey

emails: ¹zelihakorpinar@gmail.com; ²talatkorpinar@gmail.com; ³selcukbas79@gmail.com ;
⁴talatsariaydin@gmail.com

In this work, we study inextensible flows with differential geometry properties of surfaces by using Bäcklund transformations of integrable geometric Schrödinger flow. We give some new solutions by using the Schrödinger flow. Moreover, we characterize some solutions of curvature and torsion.

MSC 2000: 53C41, 53A10

Keywords: Schrödinger flow, \mathbb{E}^3 , extended Riccati mapping method, Bäcklund transformations.

References

- [1] M. do Carmo, Differential Geometry of Curves and Surfaces. *Prentice-Hall, Englewood Cliffs*, 1976.
- [2] C. Qu, J. Han and J. Kang, Bäcklund transformations for integrable geometric curve flows. *Symmetry* **7** (2015), 1376-1394.
- [3] C. Rogers and W. K. Schief, Bäcklund and Darboux Transformations Geometry and Modern Applications in Soliton Theory. *Cambridge University Press, Cambridge, UK*, 2002.
- [4] B. O'Neill, Semi-Riemannian Geometry. *Academic Press, New York*, 1983.

Idempotent unit group in commutative group rings of direct products

OMER KUSMUS¹, I. HAKKI DENIZLER², NECAT GORENTAS³

^{1,2,3} Van Yuzuncu Yil University, Van, Turkey

emails: ¹kusmuso@gmail.com; ²ihdenizler@gmail.com; ³negorentas@yyu.edu.tr

Let $\mathcal{U}(RG)$ denotes the unit group of the group ring RG of a given group G over the ring R and also $\mathcal{V}(RG)$ shows the normalized unit subgroup in $\mathcal{U}(RG)$. Idempotent units in $\mathcal{V}(RG)$ is formally defined as

$$\langle \sum_{g \in G} e_g g : \forall g \in G, e_g = e(g) \in R, e_g^2 = e_g, \sum_{g \in G} e_g = 1, e_g e_h \stackrel{g \neq h}{=} 0 \rangle$$

and displayed by $Id(RG)$. In this study, since R is a commutative unitary ring, both G and H are abelian groups, we search for some necessary and sufficient conditions for

$$Id(R(G \times H)) = Id(RG) \times Id(RH)$$

MSC 2010: 16S34, 16U60, 20K10, 20K20, 20K21

Keywords: Idempotents, nilpotents, units, unit group, group rings

References

- [1] P. Danchev, G -nilpotent units of commutative group rings. *Comment. Math. Univ. Carolin.* **53** (2012), 179–187.
- [2] P. Danchev, Idempotent units of commutative group rings. *Communications in Algebra* **38** (2010), 4649–4654.
- [3] P. Danchev, Idempotent units in commutative group rings. *Kochi J. Math.* **4** (2009), 61–68.
- [4] P. Danchev, Idempotent-nilpotent units in commutative group rings. *Bull. Greek Math. Soc.* **56** (2009), 21–28.
- [5] P. Danchev, G -unipotent units in commutative group rings. *Ann. Sci. Math. Quebec* **33** (2009), 47–53.
- [6] C. P. Milies and S. K. Sehgal, An Introduction to Group Rings. *Kluwer Acad. Publishers, New York*, 2002.

Investigating a quadratic Bezier curve according to N-Bishop frame

HATİCE KUŞAK SAMANCI¹, MUHSİN İNCESU²

¹ Bitlis Eren University, Bitlis, Turkey

² Muş Alparslan University, Muş, Turkey

emails: ¹hkusak@beu.edu.tr; ²m.incesu@alparslan.edu.tr

It is known that Bezier curve is one of the effective method for computer aided geometric design (CAGD). The second degree of general Bezier curve is called as a quadratic Bezier curve. Also, N-Bishop frame is a new Bishop frame introduced by O. Keskin and Y. Yaylı in [3]. In our work, a brief summary about a quadratic Bezier curve are firstly presented. Moreover, we research some geometric properties of the quadratic Bezier curve according to N-Bishop frame. Finally, we obtained the N-Bishop curvatures and derivative formulas for this curve.

MSC 2010: 53A04, 68U07

Keywords: Bezier Curves, CAGD, N-Bishop, curvature

References

- [1] D. Marsh, Applied Geometry for Computer Graphics and CAD. *Springer-Verlag, Berlin, 2nd edn.*, 2005.
- [2] G. Farin, A history of curves and surfaces. *Handbook of Computer Aided Geometric Design*, 2002.
- [3] O. Keskin, Y. Yayli, An application of N-Bishop frame to spherical images for direction curves. *International Journal of Geometric Methods in Modern Physics* **14** (2017), no. 11, 21 pp.; doi:10.1142/S0219887817501626.
- [4] R. L. Bishop, There is more than one way to frame a curve. *The American Mathematical Monthly* **82** (1975), no. 3, 246-251; doi:10.2307/2319846.

Review on fuzzy thermal image processing applications

FATİH KUTLU¹, ÖZKAN ATAN²

^{1,2}Van Yüzüncü Yıl University, Van, Turkey
emails: ¹fatihkutlu@yyu.edu.tr; ²oatan@yyu.edu.tr

The main purpose of this study is to improve the edge detection performance in thermal images with a new fuzzification method. In this method, contrary to other studies in the literature, the thermal image is fuzzified separately from the RGB channels and edge detection performance is improved with the aid of aggregation operators.

MSC 2010: 90C70, 94D05

Keywords: thermal image processing, fuzzy sets, edge detection

Acknowledgement: This work was supported by Research Fund of the Van Yüzüncü Yıl University. Project Number: FBA-2018-6531

References

- [1] S. J. Sangwine and R. E. Horne, The colour image processing handbook. *Springer Science & Business Media*, 2012
- [2] T. Chaira and A. K. Ray, Fuzzy image processing and applications with MATLAB. *CRC Press*, 2009.
- [3] M. EtehadTavakol, S. Sadri and E. Y. K. Ng, Application of K-and fuzzy c-means for color segmentation of thermal infrared breast images. *Journal of medical systems* **34** (2010), no. 1, 35-42.
- [4] K. Fukushima, H. Kawata, Y. Fujiwara, and H. Genno, Human sensory perception oriented image processing in a color copy system. *International Journal of Industrial Ergonomics* **15** (1995), no. 1, 63-74.
- [5] S. Araki, H. Nomura, and N. Wakami, Segmentation of thermal images using the fuzzy c-means algorithm. In: *Fuzzy Systems*, Second IEEE International Conference, (1993), 719-724.
- [6] N. M. Hien, N. T. Binh, N. Q. Viet, Edge detection based on Fuzzy C Means in medical image processing system. In: *System Science and Engineering (ICSSE)*, 2017 International Conference, (2017), 12-15.
- [7] M. Etehadtavakol, E. Y. K. Ng and N. Kaabouch, Automatic segmentation of thermal images of diabetic-at-risk feet using the snakes algorithm. *Infrared Physics & Technology* **86** (2017), 66-76.

On temporal intuitionistic fuzzy De Morgan triplets

FATİH KUTLU¹, FERİDE TUĞRUL², MEHMET ÇİTİL³

¹Van Yüzüncü Yıl University, Van, Turkey

^{2,3}Kahramanmaraş Sütçü İmam University, Kahramanmaraş, Turkey

emails: ¹fatihkutlu@yyu.edu.tr ²feridetugrul@gmail.com; ³citil@ksu.edu.tr

The aim of this study is to define negator, t-norm and t-conorms, which is the generalization of negation, conjunctions and disjunctions in the temporal intuitionistic fuzzy sets and to examine the De Morgan relations between these concepts.

MSC 2010: 94D05, 03E72

Keywords: De Morgan triplet, t-norm, t-conorm, negation, temporal intuitionistic fuzzy sets.

References

- [1] F. Kutlu, Ö. Atan and T. Bilgin, (2016). Distance Measure, Similarity Measure, Entropy and Inclusion Measure on Temporal Intuitionistic Fuzzy Sets. *Proceedings of IFSCOM* (2016), 130-148.
- [2] K. T. Atanassov, On Intuitionistic Fuzzy Sets Theory. *Springer, Berlin*, 2012.
- [3] L. Chen and C. Tu, Time-validating-based Atanassov's intuitionistic fuzzy decision-making. *IEEE Transactions on Fuzzy Systems* **23** (2015), no. 4, 743-756.
- [4] S. Yılmaz and G. Çuvalcıoğlu, On level operators for temporal intuitionistic fuzzy sets. *Notes on Intuitionistic Fuzzy Sets* **20** (2014), no. 2, 6-15.
- [5] G. Deschrijver, C. Cornelis, and E. E. Kerre, On the representation of intuitionistic fuzzy t-norms and t-conorms. *IEEE Transactions on Fuzzy Systems* **12** (2004), no. 1, 45-61.
- [6] L. A. Zadeh, Fuzzy sets. *Information and Control* **8** (1965), 338-353.
- [7] B. Bede, Mathematics of Fuzzy Sets and Fuzzy Logic, *Springer-Verlag Berlin Heidelberg*, 2013

An interplay between Riemann integrability and weaker forms of continuity

NISAR A. LONE

University of Kashmir, Srinagar, India

email: nisarsultan@gmail.com

It is folklore that a continuous function $f : [0, 1] \longleftarrow \mathbb{R}$ is Riemann integrable. The result holds true if the real space \mathbb{R} is replaced by a Banach space X . The result doesn't hold true if the function is assumed to be continuous with respect to a topology weaker than the norm topology. In this paper characterization of Banach spaces will be discussed in terms of the Riemann integrability of functions which are continuous in weaker topologies. The reverse part of the problem which results in introduction of the property of Lebesgue will also be discussed.

MSC 2010: 46G10, 46G12

Keywords: Riemann integral, weak continuity, Grothendieck space

References

- [1] A. Alexiewicz and W. Orlicz, Remarks on Riemann integration of vector-valued functions. *Studia Math.* **12** (1951), 125-132.
- [2] G. M. Cervantes, Riemann integrability versus weak continuity. *arXiv:1510.08801v1 [math.FA]* 2015.
- [3] Russel Gordon, Riemann integration in Banach spaces. *Rockey Mountain J. of Mathematics* **21** (1991), no. 3, 923-948.
- [4] C. Wang, On the weak property of Lebesgue of Banach spaces, j. Of Nanjing University. *Math. Biquarterly* **13** (1996), no. 2, 150-155.
- [5] C. Wang and Z. Yang, Some topological properties of Banach spaces and Riemann integration. *Rockey Mountain J. Of mathematics* **30** (2000), no. 1, 393-400.

Ehrlich-Aberth's type method with King's correction for the simultaneous approximation of polynomial zeros

ROSELAINNE NEVES MACHADO¹, LUIZ GUERREIRO LOPES²

¹Federal Institute of Rio Grande do Sul, Bento Gonçalves, RS, Brazil

²Faculty of Exact Sciences and Engineering, University of Madeira, Funchal, Madeira Is., Portugal

emails: ¹roselaine.machado@bento.ifrs.edu.br; ²lopes@uma.pt

There are many simultaneous iterative methods for approximating complex polynomial zeros, from more traditional numerical algorithms, such as the well-known third order Ehrlich-Aberth's method [1, 2], to the more recent ones. In this paper, we present a new combined method for the simultaneous determination of complex zeros of a polynomial, which uses the Ehrlich-Aberth iteration and a correction based on King's method [3]. Using King's correction, the order of convergence of the basic method is increased from 3 to 6. Numerical examples are given to illustrate the accuracy and computational efficiency of the proposed combined method for the simultaneous approximation of polynomial zeros.

MSC 2010: 30C10, 65H04, 65Y20

Keywords: Polynomial zeros, simultaneous iterative methods, combined iterative methods

Acknowledgement: This work was supported by the Federal Institute of Rio Grande do Sul (IFRS), Brazil.

References

- [1] O. Aberth, Iteration methods for finding all zeros of a polynomial simultaneously. *Math. Comp.* **27** (1973), no. 122, 339–344; doi:10.1090/S0025-5718-1973-0329236-7.
- [2] L. W. Ehrlich, A modified Newton method for polynomials. *Commun. ACM* **10** (1967), no. 2, 107–108; doi:10.1145/363067.363115.
- [3] R. F. King, A family of fourth order methods for nonlinear equations. *SIAM J. Numer. Anal.* **10** (1973), no. 5, 876–879; doi:10.1137/0710072.

Direct sum of neighborhoods in locally convex cones

MOHAMMAD REZA MOTALLEBI

University of Mohaghegh Ardabili, Ardabil, Iran

email: motallebi@uma.ac.ir

Using the direct sum of neighborhoods, we define the product cone topologies in locally convex cones. The polar of every product neighborhood may be written as the direct sum of its components which leads us to investigate the duality properties of product cones. In particular, we conclude that the weak topology on product cone is the locally convex product cone of its components.

MSC 2010: 20K25, 46A20, 46A03.

Keywords: Product and direct sum, duality, locally convex cone.

References

- [1] K. Keimel and W. Roth, Ordered cones and approximation. Lecture Notes in Mathematics **1517**. Springer Verlag, Heidelberg-Berlin-New York, 1992.
- [2] M. R. Motallebi, Locally convex inductive limit cones. *RACSAM* (2017); <https://doi.org/10.1007/s13398-017-0432-5>.
- [3] M. R. Motallebi, Locally convex projective limit cones. *Math. Slov.* **66** (2016), no. 6, 1387-1398.
- [4] M. R. Motallebi, On weak completeness of products and direct sums in locally convex cones. *Period. Math. Hung.* **75** (2017), no. 2, 322-329.
- [5] M. R. Motallebi and H Saiflu, Duality on locally convex cones. *J. Math. Anal. Appl.* **337** (2008), no. 2, 888-905.
- [6] W. Roth, Operator-valued measures and integrals for cone-valued functions. Lecture Notes in Mathematics **1964**. Springer Verlag, Heidelberg-Berlin-New York, 2009.

A note on superposition operators in Fibonacci sequence spaces $l_p(F)$

OĞUZ OĞUR

Giresun University, Giresun, Turkey

email: oguz.ogur@giresun.edu

Let X and Y be sequence spaces. A superposition operator P_g on X is a mapping from X to Y defined by $P_g(x) = (g(k, x_k))_{k=1}^{\infty}$, where $g : \mathbb{N} \times \mathbb{R} \rightarrow \mathbb{R}$ with $g(k, 0) = 0$ for all $k \in \mathbb{N}$. In this paper, we study on characterization of superposition operator P_g acts from $c_0(F)$ and $l_p(F)$ to $l_q(F)$, $1 \leq p, q \leq \infty$, where $F = (f_{nk})_{n,k=1}^{\infty}$ is Fibonacci matrix such as: $f_{nk} = \frac{f_k}{f_{n+2}-1}$ if $1 \leq k \leq n$ and $f_{nk} = 0$ if $k > n$. Also, we give the necessary and sufficient conditions for continuity of $P_g : l_p(F) \rightarrow l_q(F)$.

MSC 2010: 47H30, 46A45, 40A05, 46B45

Keywords: Superposition operator, Fibonacci sequence space, matrix transformation

References

- [1] T. S. Chew and P. Y. Lee, Orthogonally additive functionals on sequence spaces. *SEA Bull. Math.* **9** (1985), 81-85.
- [2] E. Kolk and A. Raidjoe, The Continuity Of Superposition Operators On Some Sequence Spaces Defined By Moduli. *Czechoslovak Mathematical Journal* **57** (2007), 777-792.
- [3] S. Petranuarat and Y. Kemprasit, Superposition operators on l_p and c_0 into l_q ($1 \leq p, q < \infty$). *Southeast Asian Bulletin of Mathematics* **21** (1997), 139-147.
- [4] R. Pluciennik, Continuity of superposition operators on w_0 and W_0 . *Comment. Math. Univ. Carolinae* **31** (1990), 529-542.
- [5] B. Sağır and N. Güngör, Continuity of superposition operators on the double sequence spaces. *Filomat* **31** (2015), no. 9, 2107-2118.
- [6] B. Sağır and N. Güngör, Locally boundedness and continuity of superposition operators on the double sequence spaces C_{r_0} . *J. Computational Analysis And Applications* **19** (2015), no. 2, 365-377.
- [7] E. E. Kara and M. İlkhan, Some properties of generalized Fibonacci sequence spaces. *Linear and Multilinear Algebra* **64** (2016), 2208-2223.
- [8] E. E. Kara, Some topological and geometrical properties of new Banach sequence spaces. *J. Inequal. Appl.* **2013** (2013), no. 38, 15 p.

On contact surgeries and a counterexample

SİNEM ONARAN

Hacettepe University, Ankara, Turkey

email: sonaran@hacettepe.edu.tr

Contact surgeries have long been an essential tool in the study of contact 3-manifolds. Contact surgeries are roughly defined as removing a neighborhood of a Legendrian knot and gluing a contact solid torus back for which we can extend the contact structure on its boundary to the inside. In this talk, we will focus on the behaviour of contact structures under contact $(+n)$ -surgeries along Legendrian knots where the surgery slope is measured with respect to the contact framing of the Legendrian knot. We give a counterexample to a conjecture by James Conway on overtwistedness of manifolds obtained by contact $(+n)$ -surgery.

MSC 2010: 53D10, 57M25, 57R65,

Keywords: Contact structure, contact surgery, Legendrian knot

Acknowledgement: This work is supported by the Scientific and Technological Research Council of Turkey, TÜBİTAK Project No:115F519.

Singular eigenvalue problems via Hilfer derivative

RAMAZAN OZARSLAN¹, ERDAL BAS², AHU ERCAN³

^{1,2,3}Firat University, Elazig, Turkey

emails: ¹ozarslanramazan@gmail.com; ²erdalbas@yahoo.com; ³ahuduman24@gmail.com

In this article singular eigenvalue problem is considered with Hilfer derivative. Self-adjointness of the operator is analyzed and some spectral properties are given. Let $\alpha \in (0, 1)$ and $\beta \in [0, 1]$. Singular eigenvalue problem with Hilfer derivative is defined as follows

$$\mathcal{L}_{\alpha[C]} y_{\lambda}(x) + \lambda w_{\alpha}(x) y_{\lambda}(x) = 0 \quad (1)$$

where $p(x) \neq 0$, $w_{\alpha}(x) > 0 \forall x \in (0, \pi]$, $w_{\alpha}(x)$ is weight function and p, q are real valued continuous functions in interval $(0, \pi]$ and $\frac{y_{\lambda}(x)}{x} \in C[0, \pi]$. The boundary conditions for the problem (1-3) are the following:

$$c_1 I_{0+}^{(1-\alpha)(1-\beta)} p(0) D_{\pi-}^{\alpha, \beta} y(0) + c_2 I_{\pi-}^{\beta(1-\alpha)} y(0) = 0, \quad (2)$$

$$d_1 I_{0+}^{(1-\alpha)(1-\beta)} p(\pi) D_{\pi-}^{\alpha, \beta} y(\pi) + d_2 I_{\pi-}^{\beta(1-\alpha)} y(\pi) = 0, \quad (3)$$

where $c_1^2 + c_2^2 \neq 0$ and $d_1^2 + d_2^2 \neq 0$.

MSC 2010: 26A33, 34A08, 35P05

Keywords: Fractional, Hilfer, singular, spectral.

References

- [1] R. Hilfer, Applications of Fractional Calculus in Physics. *World Scientific, Singapore*, 2000.
- [2] Z. Tomovski, R. Hilfer and H. M. Srivastava, Fractional and operational calculus with generalized fractional derivative operators and Mittag-Leffler type functions. *Integral Transforms and Special Functions* **21** (2010), no. 11, 797-814.
- [3] E. Bas and F. Metin, Fractional singular Sturm-Liouville operator for Coulomb potential. *Advances in Difference Equations* **2013** (2013), no. 300, 1-13.
- [4] Y. Xu and O. P. Agrawal, New fractional operators and application to fractional variational problem. *Computers & Mathematics with Applications*, In Press.
- [5] D. Baleanu and M. Og, On the existence interval for the initial value problem of fractional differential equation. *Hacettepe Journal of Math. Ist.* **4** (2011), no. 40, 581-587.

An efficient TVD-WAF scheme application for the 2D shallow water equations on unstructured meshes

NURAY ÖKTEM

Ankara Yıldırım Beyazıt University, Turkey

email: nbozkaya@gmail.com

The solutions of 2D shallow water equations (SWE) are often used to study free surface flows such as shallow lakes, dam-breaks, inundations, wide rivers, estuaries and coastal zones. Although the flow is 3D in nature, the simplified 1D or 2D shallow flow simulations may still cover a considerable amount of real flow characteristics and provide practical and quick numerical solutions in many engineering applications.

In this study a weighted averaged flux (WAF) method application [1] is presented for the solution of SWE with a new total variation diminishing (TVD) flux-limiter tool on unstructured meshes. The numerical solver is based on a 2nd order accurate (in both time and space) cell-centered finite volume formulation and it is especially constructed on an unstructured triangular mesh to be used for complex flow geometries as well. For the interface flux computations WAF alone provides 2nd order accuracy spatially but brings some numerical oscillations besides. In order to smooth these oscillations and ensure the stabilization of the interface flux solutions WAF is coupled with a new version of TVD for unstructured meshes. Shock wave development and propagation at interfaces is another difficulty in the resolution of SWE numerical models due to their hyperbolic nature. Thus in addition to TVD theory, an HLLC(Harten, Lax, van Leer-Contact wave, [2]) Riemann solver is required and so is implemented to support the shock capturing property and contact discontinuities recognition property of the coupled TVD-WAF algorithm. Moreover, a two step Runge-Kutta algorithm is applied to keep the 2nd order accuracy in time direction. Finally, a novel software written in Fortran programming language [3] for this numerical process is utilized and efficient numerical results are obtained for various benchmark problems. As a further application an open-channel junction flow is considered.

MSC 2010: 35Q35, 35L65, 65M08, 74J40

Keywords: Unstructured mesh, WAF, shallow flow, HLLC Riemann solver

References

- [1] E. F. Toro, Shock-capturing methods for free-surface shallow flows. *John Wiley & Sons, Cambridge University Press, Chichester*, 2001.
- [2] A. Harten, P.D. Lax, B. van Leer, On upstream differencing and Godunov type schemes for hyperbolic conservation laws. *SIAM Review* **25** (1983), no. 1, 35-61
- [3] N. Öktem, Computer code development for the numerical solution of two dimensional shallow flow equations on unstructured grid. *Sakarya University Journal of Science* **22** (2018), no. 2, 364-382

Blow up of solutions for a quasilinear Kirchhoff-type wave equations with degenerate damping terms

ERHAN PIŞKIN¹, FATMA EKİNCİ²

^{1,2}Dicle University, Diyarbakır, Turkey

emails: ¹episkin@dicle.edu.tr; ²ekincifatma2017@gmail.com

In this work, we analyze the influence of degenerate damping terms and source terms on the solutions of the quasilinear Kirchhoff-type wave equations. We will show the blow up of solutions in finite time with positive initial energy. This improves earlier results in the literature ([1], [2]).

MSC 2010: 35B44, 35L53

Keywords: Blow up, Kirchhoff-type wave equations, Degenerate damping terms

References

- [1] M. A. Rammaha and S. Sakuntasathien, Global existence and blow up of solutions to systems of nonlinear wave equations with degenerate damping and source terms. *Nonlinear Analysis* **72** (2010), 2658-2683.
- [2] E. Pişkin, Blow up of positive initial-energy solutions for coupled nonlinear wave equations with degenerate damping and source terms. *Boundary Value Problems* **2015** (2015), 1-11.

Semi-tensor bundle and the vertical lift of tensor fields

MURAT POLAT

Atatürk University, Erzurum, Turkey

email: murat_sel_22@hotmail.com

The present paper is devoted to some results concerning the vertical lift of tensor fields of type (p,q) from manifold M to its semi-tensor bundle tB of type (p,q) .

MSC 2010: 53A45, 55R10, 57R25

Keywords: Vector field, pull-back bundle, semi-tensor bundle.

References

- [1] T. V. Duc, Structure presque-transverse. *J. Diff. Geom.* **14** (1979), no. 2, 215-219.
- [2] H. Fattaev, The lifts of vector fields to the semitensor bundle of the type $(2, 0)$. *Journal of Qafqaz University* **25** (2009), no. 1, 136-140.
- [3] D. Husemoller, Fibre Bundles. *Springer, New York*, 1994.
- [4] V. Ivancevic and T. Ivancevic, Applied Differential Geometry, A Modern Introduction. *World Scientific, Singapore*, 2007.
- [5] H. B. Lawson and M. L. Michelsohn, Spin Geometry. *Princeton University Press, Princeton*, 1989.
- [6] A. Salimov, Tensor Operators and their Applications. *Nova Science Publ., New York*, 2013.
- [7] F. Yildirim and A. Salimov, Semi-cotangent bundle and problems of lifts. *Turk J Math* **38** (2014), no. 2, 325-339.

New Lie group of transformation for the non-Newtonian fluid flow narrating differential equations

KHALIL UR REHMAN¹, M.Y. MALIK²

^{1,2} Quaid-i-Azam University, Islamabad, Pakistan

¹ Air University, Islamabad, Pakistan

emails: ¹krehman@math.qau.edu.pk; ²drmymalik@qau.edu.pk

In this endeavour, a new Lie point of transformation for the fluid flow narrating differential equations are proposed. For this purpose a non-Newtonian fluid named tangent hyperbolic fluid is considered towards the flat surface in a magnetized flow field in the presence of both the heat and mass transfer characteristics. In addition, equation of concentration admits the role of chemically reactive species. A mathematical model in terms of the coupled PDE's is constructed. Lie group of analysis is performed to yield the Lie point of transformation for the tangent hyperbolic fluid flow narrating differential equations when both the heat and mass transfer individualities are taken into account. The resultant system of PDE's is reduced into the system of ODE's via obtained set of transformation. A self-coded computational scheme is executed and outcomes in this regard are reported by way of both the graphical and tabular structures.

MSC 2010: 22E70, 57S15, 65Z05

Keywords: Lie group of transformation, non-Newtonian fluid model, shooting method

Reproducing kernel method with Bernstein polynomials for fractional boundary value problems

MEHMET GIYAS SAKAR¹, ONUR SALDIR², FEVZİ ERDOĞAN³

^{1,2,3} Van Yuzuncu Yil University, Van, Turkey

emails: ¹giyassakar@hotmail.com; ²onursaldir@gmail.com; ³fevzier@gmail.com

In this article, a novel approach is introduced for numerical solution of linear and nonlinear boundary value problems with fractional order. Fractional derivative are taken in Caputo sense. This approach is based on reproducing kernel with Bernstein polynomials. So as to show the effect of the method, results are given as graphically and in tabulated forms.

MSC 2010: 46E22, 65Z05, 26A33

Keywords: Caputo derivative, reproducing kernel, Bernstein polynomials, boundary value problem

References

- [1] M. Cui and Y. Lin, Nonlinear numerical analysis in the reproducing kernel space. *Nova Science, New York*, 2009.
- [2] M. G. Sakar, Iterative reproducing kernel Hilbert spaces method for Riccati differential equation. *J Comput Appl Math* **309** (2017), 163–174; doi:10.1016/j.cam.2016.06.029.
- [3] M. Khaleghi, E. Babolian and S. Abbasbandy, Chebyshev reproducing kernel method: application to two-point boundary value problems. *Adv Differ Equ* **26** (2017), 1–19; doi:10.1186/s13662-017-1089-2.
- [4] M. G. Sakar, A. Akgül and D. Baleanu, On solutions of fractional Riccati differential equations. *Adv Differ. Equ.* **39** (2017), 1–10; doi:10.1186/s13662-017-1091-8.

A numerical approach for time-fractional Kawahara equation with reproducing kernel method

ONUR SALDIR¹, MEHMET GIYAS SAKAR², FEVZİ ERDOĞAN³

^{1,2,3}Van Yuzuncu Yil University, Van, Turkey

emails: ¹onursaldir@gmail.com; ²giyassakar@hotmail.com; ³ferdogan@yyu.edu.tr

We introduced a new approach based on reproducing kernel method for time fractional Kawahara equation. Approximate solution and convergence analysis of method are given. To show the power and effect of the method, an example is solved and results are given as tables and graphics. The results show that the method very convenient and efficient for Kawahara equation.

MSC 2010: 35R11, 46E22

Keywords: Reproducing kernel method, Kawahara equation, Caputo derivative, convergence

References

- [1] I. Podlubny, Fractional differential equations. *Academic Press, New York*, 1999.
- [2] M. G. Sakar, O. Saldır, A. Akgül, A novel technique for fractional Bagley-Torvik equation. *Proceedings of the National Academy of Sciences, India Section A: Physical Sciences*, 2018; <https://doi.org/10.1007/s40010-018-0488-4>.
- [3] M. Safavi, A. A. Khajehnasiri, Solutions of the modified Kawahara equation with time-and space-fractional derivatives. *Journal of Modern Methods in Numerical Mathematics* **7** (2016), no. 1, 10-18.
- [4] Y. Dereli, I. Dag, Numerical solutions of the Kawahara type equations using radial basis functions. *Numerical Methods for Partial Differential Equations* **28** (2012), no. 2, 542-553.

Rings without a middle class: past and recent

BÜLENT SARAÇ

Hacettepe University, Ankara, Turkey

email: bsarac@hacettepe.edu.tr

In 1964, Barbara Osofsky ([8]) proved her celebrated theorem which states that any ring R with identity is semisimple artinian if and only if every cyclic right (or left) R -module is injective. This happened to be the first step in search of characterizations of rings by homological properties of some certain type of their modules. As a part of this long standing research program, a new type of rings were introduced in 2011 (see [3]) and has been studied extensively by several authors including the speaker. In this talk, I would like to present some new ideas from torsion theory by which we can unify the developing theory of rings of the title and show how these ideas can be effective in exploring many significant properties of these rings. In the last part of my talk, I would also like to discuss the case, which is still a mystery, where the rings are not right noetherian and apply our new techniques to work out this mysterious case.

MSC 2010: 16D50, 16D70, 16P20

Keywords: Injective module, poor module, injectivity domain, QI-, PCI-, V-ring

Acknowledgement: This talk is mainly based on results obtained under the research project supported by TUBITAK with project number 117F084.

References

- [1] A. N. Alahmadi, M. Alkan and S. López-Permouth, Poor modules: the opposite of injectivity. *Glasg. Math. J.* **52** (2010), no. A, 7–17.
- [2] P. Aydoğdu and B. Saraç, On Artinian rings with restricted class of injectivity domains. *J. Algebra* **377** (2013), 49–65.
- [3] N. Er, S. López-Permouth and N. Sökmez, Rings whose modules have maximal or minimal injectivity domains. *J. Algebra* **330** (2011), 404–417.
- [4] N. Er, S. López-Permouth, N.K. Tung, Rings whose cyclic modules have restricted injectivity domains. *J. Algebra* **466** (2016), 208–228.
- [5] K. R. Goodearl, Singular Torsion and the Splitting Properties. *Memoirs of the American Mathematical Society*, vol. 124, AMS, 1972.
- [6] S. R. López-Permouth and J.E. Simental, Characterizing rings in terms of the extent of the injectivity and projectivity of their modules. *J. Algebra* **362** (2012), no. 1, 56–69.
- [7] G.O. Michler and O.E. Villamayor, On rings whose simple modules are injective. *J. Algebra* **25** (1973), 185–201.
- [8] B. Osofsky, Rings all of whose finitely generated modules are injective. *Pacific J. Math.* **14** (1964), 645–650.

Some applications about Möbius function

UMİT SARP¹, DAEYOUL KIM², SEBAHATTİN İKİKARDES³

^{1,3}Balikesir University, Balikesir, Turkey

²Chon-buk National University, 567 Baekje-daero, Republic of Korea

emails: ¹umitsarp@gmail.com; ²kdaeyeoul@jbnu.ac.kr; ³skardes@balikesir.edu.tr

In this paper, according to some numerical computational evidence, we investigate and prove certain relations and properties on Möbius function and some related functions.

MSC 2010: 11M36, 11F11, 11F30

Keywords: Möbius function, divisor function, perfect number.

Acknowledgement: This work was supported by Balikesir University Research, Grant No: 2017/20.

References

- [1] A. Bayad and D. Kim, Polygon numbers associated with the sum of odd divisors function. *to appear of Exp. Math.* **26** (2017), no. 3, 287-297.
- [2] L. E. Dickson, History of the theory of numbers. Vol. I: Divisibility and primality. *Chelsea Publishing Co., New York*, 1966.
- [3] P. Erdős, Some remarks on Euler's ϕ function and some related problems. *Bull. Amer. Math. Soc.* **51** (1945), 540-544.
- [4] D. Reiss, C. Jackson and B. Dallas, Stirling Numbers of the 1st Kind. *Western Washington University*, 2012.

Estimation of parameters of Gumbel distribution data

SANEM SEHRIBANOGLU

Van Yuzuncu Yil University, Van, Turkey

email: sanem@yyu.edu.tr

The Gumbel distribution is one of the particular states of generalized extreme value distribution (GEV). Probabilistic extreme value theory is an interesting and fascinating area with a great variety of applications. In probability theory and statistics, this distribution is used to model for the extremes (maximum or minimum) observations. The Gumbel distribution are frequently applied to forecast of natural events such as floods, air pollution, extreme sea levels, hydrology, meteorology, climatology, insurance, finance, geology and seismology.

In this paper, The parameters are estimated using maximum likelihood and Bayesian estimation procedure. In addition to these inferences are used The Newton-Raphson algorithm and Markov Chain Monte Carlo (MCMC) simulation method.

MSC 2010: :62H10, 62F15

Keywords: Generalized extreme value distribution, Gumbel distribution, MLE, Bayesian estimation

References

- [1] E. J. Gumbel, Bivariate exponential distributions. *Journal of the American Statistical Association* **55** (1960), no. 292, 698-707.
- [2] G. Gholami, Bayesian estimation of parameters in the exponentiated Gumbel distribution. *J. Statist. Res. Iran* **13** (2016), 181-195.
- [3] L. M. Lye, K. P. Hapuarachchi and S. Ryan, Bayes estimation of the extreme-value reliability function. *IEEE Transactions on Reliability* **42** (1993), no. 4, 641-644.
- [4] H. Demirhan, A generalized Gumbel distribution and its parameter estimation. *Communications in Statistics-Simulation and Computation* (2017), 1-20; doi:10.1080/03610918.2017.1361976.
- [5] S. F. Bagheri, M. Alizadeh and S. Nadarajah, Efficient estimation of the PDF and the CDF of the exponentiated Gumbel distribution. *Communications in Statistics - Simulation and Computation* **45** (2015), no. 1, 339-361; doi: 10.1080/03610918.2013.863922.
- [6] K. Cooray, Generalized Gumbel distribution. *Journal of Applied Statistics* **37** (2010) no. 1, 171-179.
- [7] E. Towler, B. Rajagopalan, E. Gilleland, R. S. Summers, D. Yates and R. W. Katz, Modeling hydrologic and water quality extremes in a changing climate: A statistical approach based on extreme value theory, *Water Resources Research* **46** (2010), W11504; doi: 10.1029/2009WR008876.

Some convergences in metric spaces

GUZİDE SENEL

Amasya University, Amasya, Turkey

email: g.senel@amasya.edu.tr

Porosity was defined by Denjoy in [1]. A detailed information about this type porosity was given by Thomson in [1]. The notion of porosity can also be used in metric space by replacing intervals with balls (see [3]) In this study, I will define a new type of convergence for metric valued sequences. Then I will give some properties of this new concept.

MSC 2010: 47A12, 15A60, 26C10,30C15

Keywords: Metric space, sequence, porosity

References

- [1] M. Altınok and M. Kucukaslan, Porosity convergence in metric spaces. *IFSCOM2017 Abstract No.5*, 2017.
- [2] A. Denjoy, Sur une propriete des series trigonometriques. *Verlag v.d. G. V. der Wie-en Natuur. Afd.* **29** (1920), 220-232.
- [3] B. S. Thomson, Real Functions, Lecture Notes in Mathematics, 1170, *Springer-Verlag, Berlin, Heidelberg, New York, Tokyo*, 1985.
- [4] L. Zajicek, Porosity and σ -porosity. *Real Analysis Exchange* **13** (1987-88), 314-350.

Existence of positive solutions for boundary value problems of nonlinear fractional differential equations

TUGBA SENLIK CERDIK¹, FULYA YORUK DEREN², NUKET AYKUT HAMAL³

¹Izmir, Turkey

^{2,3}Ege University, Izmir, Turkey

emails: ¹tubasenlik@gmail.com; ²fulya.yoruk@ege.edu.tr; ³nuket.aykut@ege.edu.tr

This study deals with the existence result of positive solution for the fractional boundary value problem. The arguments are based upon a fixed point theorem in a cone. Its application is also given.

MSC 2010: 34B10, 34B15, 34B18.

Keywords: Boundary value problem, positive solution, fractional differential equation.

References

- [1] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, Theory and Applications of Fractional Differential Equations. *Elsevier Science B.V, Amsterdam*, 2006.
- [2] I. Podlubny, Fractional Differential Equations. *Academic Press, San Diego*, 1999.
- [3] A. Guezane-Lakouda and R. Khaldi, Solvability of a fractional boundary value problem with fractional integral condition. *Nonlinear Analysis* **75** (2012), 2692-2700.

Elliptic curves containing sequences of consecutive cubes

GÖKHAN SOYDAN¹, GAMZE SAVAŞ ÇELİK²

^{1,2}Uludağ University, Bursa, Turkey

emails: ¹gsoydan@uludag.edu.tr; ²gamzesavascelik@gmail.com

Let us consider a rational elliptic curve given by a Weierstrass equation

$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6 \quad (1)$$

with $a_1, \dots, a_6 \in \mathbb{Q}$. We will say that the points $(x_i, y_i), i = 1, \dots, k$ on the curve (1) are in arithmetic progression of length k if the sequence x_1, x_2, \dots, x_k forms an arithmetic progression (AP for short). Firstly, Lee and Vélez, [5], found infinitely many curves of type $y^2 = x^3 + a$ containing $k = 4$ -length APs. Then many authors considered elliptic curves containing $k \geq 8$ -length APs (see [1], [2], [6]).

Recently, Kamel and Sadek, [4], considered sequences of rational points on elliptic curves given by the equation

$$y^2 = ax^3 + bx + c \quad (2)$$

over \mathbb{Q} whose x -coordinates form a sequence of “consecutive squares”.

In this work, we are interested in sequences of rational points on elliptic curves with the equation (2) whose x -coordinates form a sequence of “consecutive cubes”. We show that elliptic curves given by the equation (2) with 5-term sequences of rational points whose x -coordinates are elements of a sequence of consecutive cubes in \mathbb{Q} parametrized by an elliptic surface whose rank is positive. This implies the existence of infinitely many such elliptic curves. We also show that these five rational points in the sequence are linearly independent in the group of rational points of the elliptic curve they lie on. Especially, we introduce an infinite family of elliptic curves of rank ≥ 5 [3].

MSC 2010: 14G05, 11B83.

Keywords: Elliptic curves, rational points, sequences of consecutive cubes.

References

- [1] A. Bremner, On arithmetic progressions on elliptic curves. *Experiment Math.* **8** (1999), 409–413.
- [2] G. Campbell, A note on arithmetic progressions on elliptic curves. *J. Integer Seq.* **6** (2003), Article 03.1.3.
- [3] G. S. Çelik, G. Soydan, Elliptic curves containing sequences of consecutive cubes. *Rocky Mountain J. Math.* (2018), to appear.
- [4] M. Kamel, M. Sadek, On sequences of consecutive squares on elliptic curves. *Glasnik Math.* **52** (2017), 45–52.
- [5] J. B. Lee, W. Y. Vélez, Integral solutions in arithmetic progression for $y^2 = x^3 + k$. *Per. Math. Hung.* **25** (1992), 31–49.
- [6] M. Ulas, A note on arithmetic progressions on quartic elliptic curves. *J. Integer Seq.* **8** (2005), Article 05.3.1.

Jones polynomial for graphs of twist knots

ABDULGANİ ŞAHİN¹, BÜNYAMİN ŞAHİN²

¹Ağrı İbrahim Çeçen University, Ağrı, Turkey

²Bayburt University, Bayburt, Turkey

emails: ¹rukassah@gmail.com; ²bsahin@bayburt.edu.tr;

Graphs have made a great contribution to the development of algebraic topology. Along with this support, knot theory has taken an important place in low dimensional manifold topology. In 1984, Jones introduced a new polynomial for knots. The discovery of that polynomial opened a new era in knot theory. In a short time, this polynomial was defined by algebraic arguments and its combinatorial definition was made. The Jones polynomials of knot graphs and their applications were introduced by Murasugi. T. Uur and A. Kopuzlu found an algorithm for the Jones polynomials of torus knots $K(2, q)$ in 2006. In this paper, we compute the Jones polynomials for graphs of twist knots. We will consider signed graphs associated with each twist knot diagrams.

MSC 2000: 57M15, 57M25, 57M27

Keywords: Twist knots, knot graph, Jones polynomial

References

- [1] A. Kawauchi, A Survey of Knot Theory. *Birkhauser Verlag, Basel-Boston-Berlin*, 1996.
- [2] B. Bollobas, Modern Graph Theory *Springer Science + Business Media, Inc, New York*, 1998.
- [3] C. C. Adams, The Knot Book. *W. H. Freeman and Company, New York*, 1994.
- [4] I. Johnson, A. K. Henrich, A Interactive Introduction to Knot Theory. *Dover Publications, Inc., Mineola, New York*, 2017.
- [5] J. Hoste, P. D. Shanahan, Trace Fields of Twist Knots. *Journal of Knot Theory and Its Ramifications* **10(4)** (2001), 625–639; doi:10.1142/S0218216501001049.
- [6] K. Murasugi, Invariants of Graphs and Their Applications to Knot Theory. *In: Algebraic Topology Poznan* (1989), 83–97; doi:10.1007/BFb0084739. *Lecture Notes in Mathematics, vol 1474, Springer, Berlin, Heidelberg*, 1991.
- [7] K. Murasugi, Knot Theory and Its Applications. *Birkhauser Verlag, Boston*, 1996.
- [8] M. Henle, A Combinatorial Introduction to Topology. *Dover Publications, Inc., New York*, 1994.
- [9] T. Uğur, A. Kopuzlu, On Jones Polynomials of Graphs of Torus Knots $K(2, q)$. *International Mathematical Forum* **1**, no. **31** (2006), 1537–1541.

Kernel stable and uniquely generated modules

SERAP ŞAHİNKAYA¹, TRUONG CONG QUYNH²

¹Gebze Technical University, Kocaeli, TURKEY

² Danang University, DaNang City, VIETNAM

emails: ¹ssahinkaya@gtu.edu.tr; ²tcquynh@dce.udn.vn

Module theoretic notion of annihilator-stable rings which was defined recently by Nicholson [2] is defined and some characterizations of it studied. M is called kernel-stable module if every element $\alpha \in \text{End}(M)$ satisfies the following condition: If $\alpha(M) + \text{Ker}\beta = M$, $\beta \in \text{End}(M)$ then $(\alpha - \gamma)(m) \in \text{Ker}\beta$ for an automorphism of M and for all $m \in M$. For a pseudo-semi-projective module M , this notion is equivalent to uniquely generated module which was defined in [1].

MSC 2010: 16U60, 16U99 16E50, 16L30 19A13.

Keywords: Stable range, annihilator-stable rings, uniquely generated modules, von Neumann regular rings, unit-regular rings, matrix rings, pseudo-semi-projective module, kernel-stable modules

References

- [1] M. T. Kosan, T. C. Quynh, S. Sahinkaya, On rings with associated elements. *Comm. Algebra* **45** (2017), no. 7, 2747-2756.
- [2] W. K. Nicholson, Annihilator-stability and unique generation. *J. Pure and App. Algebra* **221** (2017), no. 10, 2557-2572.

Generalized class of boundary value problems with a constant retarded argument

ERDOĞAN ŞEN

Tekirdağ Namık Kemal University, Tekirdağ, Turkey

email: erdogan.math@gmail.com

In this study we shall find asymptotic formulas of eigenvalues and eigenfunctions for the eigenvalue problem $L := L(q; a, \lambda, r; h, H, d_j)$ ($j = 1, 2, 3$) which consists of Sturm-Liouville equation

$$-y''(x) + q(x)y(x - a) = \lambda^2 r(x)y(x) = 0$$

on $\Lambda = \cup \Lambda^\pm$ with boundary conditions

$$y'(0) - hy(0) = 0,$$

$$y'(T) + Hy(T) = 0$$

and transmission conditions

$$y(c+0) = d_1 y(c-0),$$

$$y'(c+0) = d_2 y'(c-0) + d_3 y(c-0)$$

where $r(x) = \frac{1}{r_1^2}$ for $x \in \Lambda^- = [0, c)$ and $r(x) = \frac{1}{r_2^2}$ for $x \in \Lambda^+ = (c, T]$; the real-valued function $q(x)$ is continuous in Λ and has a finite limit $q(c \pm 0) = \lim_{x \rightarrow c \pm 0} q(x)$, $x - a \geq 0$, if $x \in \Lambda^-$; $x - a \geq c$, if $x \in \Lambda^+$; λ is a real spectral parameter; a, r_i ($i = 1, 2$), h, H, d_j ($j = 1, 2, 3$) are arbitrary real numbers such that $r_1 r_2 d_1 d_2 \neq 0$ and $d_1 r_2 = d_2 r_1$.

MSC 2010: 34K10, 34L20, 35R10

Keywords: Differential equation with retarded argument, transmission conditions, asymptotics of eigenvalues and eigenfunctions

Dual pole indicatrix curve and surface

SÜLEYMAN ŞENYURT¹, ABDUSSAMET ÇALIŞKAN²

^{1,2} Ordu University, Ordu, Turkey

emails: ¹senyurtsuleyman@hotmail.com; ²abdussamet65@gmail.com;

In this paper, the vectorial moment of the unit Darboux vector, which consists of the motion of the Frenet vectors on any curve, is reexpressed in the form of Frenet vectors. According to the new version of C^* vector, the parametric equation of the ruled surface corresponding to the unit dual pole indicatrix curve is given. The integral invariants of the closed ruled surface are rederived and illustrated by presenting with examples.

MSC 2010: 14H45, 14H50, 53A04

Keywords: Darboux vector, ruled surface, dual pole indicatrix curve, vectorial moment

References

- [1] A. Abdel-Baky Rashad, An explicit characterization of dual spherical curve. *Commun. Fac. Sci. Univ. Ank. Series A* **51** (2002), no. 2, 1-9.
- [2] A. Abdel-Baky Rashad, On the Blaschke approach of ruled surface, *Tamkang Journal of Math.* **34** (2003), no. 2, 107-116.
- [3] H. H. Hacısalihoğlu, On the pitch of a closed ruled surface. *Mech. Mach. Theory* **7** (1972), 291-305.
- [4] İ. A. Güven, S. Kaya and H. H. Hacısalihoğlu, On closed ruled surfaces concerned with dual Frenet and Bishop frames. *Journal of Dynamical Systems and Geometric Theories* **9** (2011), no. 1, 67-74.
- [5] O. Gürsoy , The dual angle of pitch of a closed ruled surface. *Mech. Mach. Theory* **25** (1990), no. 2, 131-140.
- [6] O. Gürsoy, On the integral invariants of a closed ruled surface. *Journal of Geometry* **39** (1990), 89-91.
- [7] W. Fenchel, On the Differential Geometry of Closed Space Curves, *Bull. Amer. Math. Soc.*, **57** (1951), no. 1-2, 44-54.
- [8] Y. Tuncer , Vectorial moments of curves in Euclidean 3- space. *International Journal of Geometric Methods in Modern Physics* **14** (2017), no. 2, 1750020(16p).
- [9] Y. Yaylı and S. Saraçoğlu, Some notes on dual spherical curves. *Journal of Informatics and Mathematical Sciences* **3** (2011), no. 2, 177–189.

Curves and ruled surfaces according to alternative frame in dual space

SÜLEYMAN ŞENYURT¹, ABDUSSAMET ÇALIŞKAN²

^{1,2} Ordu University, Ordu, Turkey

emails: ¹senyurtsuleyman@hotmail.com; ²abdussamet65@gmail.com;

In this paper, the vectorial moments of the alternative vectors are expressed in terms of alternative frame. According to the new versions of these vectorial moments, the parametric equations of the closed ruled surfaces corresponding to the (\widehat{N}) , (\widehat{C}) , (\widehat{W}) dual curves are given. The integral invariants of the these surfaces are computed and illustrated by presenting with examples.

MSC 2010: 14H45, 14H50, 53A04

Keywords: Alternative frame, closed ruled surface, vectorial moment, distribution parameter, Gauss curvature, dual angle of pitch, viviani's curve.

References

- [1] M. P. Do Carmo, Differential geometry of curves and surfaces. *Prentice-Hall Englewood Cliffs, NJ MATH Google Scholar*, 1976.
- [2] W. Fenchel, On the Differential Geometry of Closed Space Curves. *Bull. Amer. Math. Soc.* **57** (1951), 44-54.
- [3] İ. A. Güven, S. Kaya and H. H. Hacısalihoğlu, On closed ruled surfaces concerned with dual Frenet and Bishop frames. *J. Dyn. Syst. Geom. Theor.* **9** (2011), no. 1, 67-74.
- [4] O. Gürsoy, On the integral invariants of a closed ruled surface. *J. Geom.* **39** (1990), no. 1, 80-91.
- [5] H. H. Hacısalihoğlu, On the pitch of a closed ruled surface. *Mech. Mach. Theory* **7** (1972), no. 3, 291-305.
- [6] O. Kaya and M. Önder, New Partner Curves in The Euclidean 3-Space E^3 . *Int. J. Geom.* **6** (2017), no. 2, 41-50.
- [7] A. A. B. Rashad, An explicit characterization of dual spherical curve. *Commun. Fac. Sci. Univ. Ank. Series A.* **51** (2002), no. 2, 43-49.
- [8] Y. Tunçer, Vectorial moments of curves in Euclidean 3-space. *International Journal of Geometric Methods in Modern Physics* **14** (2017), no. 2, 1750020.
- [9] B. Uzunoglu, İ. Gök and Y. Yaylı, A new approach on curves of constant precession. *Appl. Math. Comput.* **275** (2016), 317-232.
- [10] Y. Yaylı and S. Saraçoğlu, Ruled surfaces and dual spherical curves. *Acta Univ. Apulensis* **30** (2012), 337-354.

Schrödinger equation with potential vanishing exponentially fast

TANFER TANRIVERDI

Harran University, Şanlıurfa, Turkey

e-mail: ttanriverdi@harran.edu.tr

Explicit solutions of differential equation

$$y'' + (\lambda + 20 \operatorname{sech}^2 x) y = 0y$$

and its eigenvalues are obtained by calculating complex residues. Eigenfunction expansions for this differential equation are also explored [1, 2, 3, 4, 5, 6, 7, 8].

MSC 2010: 35A24, 35B05, 35B24, 32A27

Keywords: Sturm-Liouville, Schrödinger equation, complex residues, explicit solutions

Acknowledgement: The Author thanks to HÜBAK (Scientific Research Council of Harran University)

References

- [1] E. C. Titchmarsh, Eigenfunction Expansions Associated with Second-Order Differential Equations Part I. *Oxford University Press, Oxford*, 1962.
- [2] T. Tanriverdi, Boundary-Value Problems in ODE, Ph.D. thesis. *University of Pittsburgh, Pittsburgh*, 2001.
- [3] T. Tanriverdi and J. B. McLeod, Generalization of the eigenvalues by contour integrals. *Appl. Math. Comput.* **189** (2007), no. 2, 1765–1773; <https://doi.org/10.1016/j.amc.2006.12.055>.
- [4] T. Tanriverdi and J. B. McLeod, The Analysis of Contour Integrals. *Abstr. Appl. Anal.* **2008** (2008), Article ID 765920; <http://dx.doi.org/10.1155/2008/765920>.
- [5] T. Tanriverdi, Contour integrals associated differential equations. *Math. Comput. Modelling* **49** (2009), no. 34, 453–462; <https://doi.org/10.1016/j.mcm.2008.05.051>.
- [6] T. Tanriverdi, Differential equations with contour integrals. *Integral Transforms Spec. Funct.* **20** (2009), no. 2, 119–125; <https://doi.org/10.1080/10652460802499927>.
- [7] W. N. Everitt, A catalogue of Sturm-Liouville differential equations. *Birkhäuser Verlag, Basel*, 2005.
- [8] T. Tanriverdi, Evaluating Sine and Cosine Type Integrals. *IJASM* **5** (2018), no. 2, 11–13.

Blow up of solutions for a stochastic Klein-Gordon equation

HATICE TASKESEN

Van Yuzuncu Yil University, Van, Turkey

email: haticetaskesen@yyu.edu.tr

The Klein-Gordon equation is the first relativistic equation in quantum mechanics for the wave function of a particle with zero spin. It occurs in the study of various problems of mathematical physics such as general relativity, plasma physics, nonlinear optics, radiation theory, fluid mechanics, and was investigated in many papers [1], [2], [3], [4]

In the present work, we investigate the effect of stochastic terms on the explosion of solutions for a stochastic Klein-Gordon equation. By using a differential inequality and an energy inequality we prove that solutions of the problem blow-up in a finite time.

MSC 2010: 60H15, 35B44, 35Q40

Keywords: Stochastic Klein-Gordon equation, blow-up, energy inequality

References

- [1] Y. Luo, Y. Yang, S. Ahmed, T. Yu, M. Zhang, L. Wang, H. Xu, Global existence and blow up of the solution for nonlinear Klein-Gordon equation with general power-type nonlinearities at three initial energy levels. *Applied Numerical Mathematics* In press.
- [2] A. I. Komech and E. A. Kopylova, Weighted energy decay for 3D Klein-Gordon equation. *J. Differential Equations* **248** (2010), no. 3, 501-520.
- [3] M. Danielewski and L. Sapa, Nonlinear Klein-Gordon equation in Cauchy-Navier elastic solid. *Cherkasy University Bulletin:Physical and Mathematical Sciences* **1** (2017), 22-28.
- [4] M. Dimova, N. Kolkovska and N. Kutev, Revised concavity method and application to Klein-Gordon equation. *Filomat* **30** (2016), no. 3, 831-839.

On the blow-up of solutions for a stochastic Camassa-Holm equation

HATICE TASKESSEN¹, MOHANAD ALALOUSH²

^{1,2}Van Yuzuncu Yil University, Van, Turkey

emails: ¹haticetaskesen@yyu.edu.tr; ²alaloush.mohanad@gmail.com

In fluid dynamics, the Camassa-Holm (CH) equation is an integrable, bi-Hamiltonian model, which is proposed to explain the one-way propagation of shallow water waves on a flat bottom [1], [2]. However, it is difficult to include all the external effects in the deterministic equation, for example, the bottom of fluid may not be so flat or there will be some environmental noises. Therefore, a stochastic term must be added to the equation [3], [4]. In this work, the stochastic modified Camassa-Holm (SCH) equation is considered. Conditions that guarantee blow up of solutions in finite time are given.

MSC 2010: 60H15, 35B44, 35Q53

Keywords: Stochastic modified Camassa-Holm equation, blow-up, integrable equations

References

- [1] R. Camassa, D. D. Holm, An integrable shallow water equation with peaked solitons. *Phys. Rev. Lett.* **71** (1993), 1661–1664.
- [2] A. Constantin, J. Escher, Well-posedness, global existence and blow-up phenomena for a periodic quasi-linear hyperbolic equation. *Comm. Pure Appl. Math.* **51** (1998), 475-504.
- [3] Y. Chen, H. Gao, Well-posedness and large deviations of the stochastic modified Camassa-Holm equation. *Potential Anal.* **45** (2016), no. 2, 331-354.
- [4] H. Tang, On the Camassa–Holm equation with multiplicative noise. *SIAM J. Math. Anal.* **50** (2018), no. 1, 1322–1366.

Caristi type related fixed point theorems in two metric spaces

MUSTAFA TELCİ

Trakya University, Edirne, Turkey

email: mtelci@trakya.edu.tr

Let (X, d) and (Y, ρ) be metric spaces, let T be a mapping of X into Y and let S be a mapping of Y into X . It is proved that if there exists $x_0 \in X$ such that (X, d) and (Y, ρ) are related complete and

$$\max\{d(x, STx), \rho(y, TSy)\} \leq \varphi(x) - \varphi(STx) + \psi(y) - \psi(TSy)$$

for all x in $R_X(x_0)$ and y in $R_Y(x_0)$, where $\varphi : X \rightarrow [0, \infty)$ and $\psi : Y \rightarrow [0, \infty)$, then

1. $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} (ST)^n x_0 = z$ and $\lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} T(ST)^{n-1} x_0 = w$ exist.
2. $Sw = z$ and $Tz = w$ if and only if $F : X \times Y \rightarrow [0, \infty)$, $F(x, y) = d(x, Sy)$ and $G : X \times Y \rightarrow [0, \infty)$, $G(x, y) = \rho(y, Tx)$ are w.l.s.c. at (z, w) .

Further, if (2) holds, then $STz = z$ and $TSw = w$.

Some related fixed point results in two metric spaces are also derived by considering this result.

MSC 2010: 47H10, 54H25

Keywords: Fixed point, related complete, weak lower semi-continuous

References

- [1] A. Bollenbacher and T. L. Hicks, A fixed point theorem revisited. *Proc. Amer. Math. Soc.* **102** (1988), 898-900.
- [2] J. Caristi, Fixed point theorems for mappings satisfying inwardness conditions. *Trans. Amer. Math. Soc.* **215** (1976), 241-251.
- [3] B. Fisher, Fixed points on two metric spaces. *Glasnik Matematički* **16** (1981), no. 36, 333-337.
- [4] V. Popa and M. Telci, Fixed Point theorems for mappings implicit relations on two metric spaces. *Mathematica Balkanica New Series* **20** (2006), no. 2, 143-152.
- [5] M. Telci, Fixed points on two complete and compact metric spaces. *Applied Mathematics and Mechanics (English Edition)* **22** (2001), no. 5, 564-568.

Krasnoselskii fixed point theorem for singlevalued operators and multivalued operators

CESİM TEMEL

Van Yuzuncu Yil University, Van, Turkey

email: cesimtemel@yyu.edu.tr

In this study, we give some results of Krasnoselskii fixed point theorem for singlevalued operators and multivalued operators under weak topology in Banach spaces. In particular, we present the solutions of nonlinear operator equation

$$u = L(u) + S(u), \quad u \in U,$$

where L and S are weakly sequentially continuous operators. We also establish the existence of the solutions of inclusions of the form

$$u \in L(u) + S(u), \quad u \in U,$$

where L is based on the generalized D -Lipschitzian, $I - L$ may not be injective and S has weakly sequentially closed graph.

MSC 2010: 34K13, 47H04, 47H10

Keywords: Fixed point theorem, Krasnoselskii fixed point theorem, weakly sequentially continuous operator, multivalued operator

References

- [1] O. Arino, S. Gautier and J. P. Pento, A fixed point theorem for sequentially continuous mapping with application to ordinary differential equations. *Functional Ekvac.* **27** (1984), no. 3, 273-279.
- [2] C. Avramescu and C. Vladimirescu, A fixed points theorem of Krasnoselskii type in a space of continuous functions. *Fixed Point Theory* **5** (2004), 1-11.
- [3] B. C. Dage, Some generalization of multi-valued version of Schauder's fixed point theorem and applications. *Cubo* **12** (2010), 139-151.
- [4] Y. C. Liu and Z. X. Li, Krasnoselskii type fixed point theorems and applications. *Proc. Amer. Math. Soc.* **136** (2008), no. 4, 1213-1220.

Crossed modules of group-groupoids and double group-groupoids

SEDAT TEMEL¹, TUNCAR SAHAN², OSMAN MUCUK³

¹Recep Tayyip Erdogan University, Turkey

²Aksaray University, Turkey

³Erciyes University, Turkey

emails: ¹sedat.temel@erdogan.edu.tr; ²tuncarsahan@aksaray.edu.tr; ³mucuk@erciyes.edu.tr

The notion of a crossed module which is initially introduced by Whitehead in [4] is related to the second relative homotopy groups for topological spaces. The categorical equivalence of crossed modules and group-groupoids which are also known in literature as *2-groups* was proved by Brown and Spencer in [2]. This categorical equivalence has also been extended by Porter in [3, Section 3] to a more general algebraic categories called categories of groups with operations. Double groupoids which can be thought as a groupoid objects in the category of groupoids are very useful for the proof of 2-dimensional Seifert-van-Kampen Theorem.

In this study we define a double group-groupoid to be a group object in the category of double categories and prove that these types of double groupoids are categorically equivalent to the crossed modules of group-groupoids and crossed square of groups.

MSC 2010: 20L05, 22A22, 18D35.

Keywords: Double category, crossed module, group-groupoid

References

- [1] S. Temel, T. Sahan and O.Mucuk, Crossed modules, double group-groupoids and crossed squares. Available as <https://arxiv.org/abs/1802.03978>.
- [2] R. Brown and C. B. Spencer, G-groupoids, crossed modules and the fundamental groupoid of a topological group. *Proc. Konn. Ned. Akad.* **64** (1976), no. 79, 296-302.
- [3] T. Porter, Extensions, crossed modules and internal categories in categories of groups with operations. *Proc. Edinb. Math. Soc.* **30** (1987), 373-381.
- [4] J. H. C Whitehead, Note on a previous paper entitled "On adding relations to homotopy group". *Ann. of Math.* **47** (1946), 806-810.

An optimality condition for non-smooth convex problems via $*$ -subgradient

ALI HAKAN TOR

Abdullah Gul University, Kayseri, Turkey

email: hakan.tor@agu.edu.tr

There are several optimality conditions for non-smooth unconstrained convex optimization problems via several types of subgradients. While some of them are being used by some researcher, all of them are needed according to the type of problem. In other words, this variety arises from the structure of optimization problems. In this work, a new optimality condition for the non-smooth convex optimization problem is given by using $*$ -subgradient. $*$ -subgradient is a new concept based on logarithmic differentiation. Actually, however logarithmic differential is valid for smooth function, $*$ -subgradient can be used a non-smooth function, not necessarily differentiable.

MSC 2010: 49K99, 46N10, 47N10

Keywords: Optimality conditions, non-smooth convex problem, convex analysis

On the orbit surface of two parameter motion

FATİH TUĞRUL¹, ŞENAY BAYDAŞ², BÜLENT KARAKAŞ³

^{1,2,3}Van Yuzuncu Yil University, Van, Turkey

emails: ¹fatihugrul@yyu.edu.tr; ²senay.baydas@gmail.com; ³bulentkarakas@gmail.com

A displacement has that rotation axis is tangent, binormal, normal, Darboux vector, etc of a curve and translation vector with the knowledge of the curve is special displacement with two parameters. Image of a point under a displacement with two parameters is a surface. In this study, kinematics structures and properties of two parameters displacement are examined.

MSC 2010: 16H05, 70E15, 11E88

Keywords: Clifford algebra, 2 parameter motion, orbit surface

References

- [1] S. Aydintan, 2-Parametric Motions Along a Curve And Orbit Surface. *Gazi University, Ankara*, 1993.
- [2] E. Bayro-Corrochano, Motor algebra aproach for computing the kinematics of robot manipulators. *Journal of Robotic System* **17** (2000), no. 9, 495-516.
- [3] O. Bottema and B. Roth, Theotical Kinematics. *North Holland Publ. Com.*, 1979.
- [4] M. K. Karacan and Y. Yayli, On the instantaneous screw axes of two parameter motions. *Facta Universitatis, Series, Mechanics, Automatic, Control and Robotics* **6** (2007), no. 1, 81-88.
- [5] P. Lounest, Clifford algebras and spinors. *Bulg. J.Phys.* **38** (2011), 3-28.
- [6] J. Vince, Geometric Algebra:An Algebraic System for Computer Games and Animation. *Springer-Verlag London Limited*, 2009.

Instability in nonlinear functional differential equations of higher order

CEMİL TUNÇ

Van Yuzuncu Yil University, Van, Turkey

email: cemtunc@yahoo.com

The author gives sufficient conditions for instability of solutions of some nonlinear delay differential equations of higher order. The technic of the proofs is based on the construction of suitable Lyapunov functionals. Some examples are given to illustrate the results obtained. The results of this paper improve some recent results can found in the literature.

MSC 2010: 39B82, 34K06

Keywords: differential equation, stability, higher order

References

- [1] Ezeilo, J. O. C., Instability theorems for certain fifth-order differential equations. *Math. Proc. Cambridge Philos. Soc.*, no. 2, 84 (1978), 343350.
- [2] Li, Wen-jian; Duan, Kui-chen, Instability theorems for some nonlinear differential systems of fifth order. *J. Xinjiang Univ. Natur. Sci.* 17 (2000), no. 3, 15.
- [3] Li, W. J.; Yu, Y. H., Instability theorems for some fourth-order and fifth-order differential equations. *J. Xinjiang Univ. Natur. Sci.*, no. 2, 7 (1990), 710.
- [4] Sadek, A. I., Instability results for certain systems of fourth and fifth order differential equations. *Appl. Math. Comput.* 145 (2003), no. 23, 541549.
- [5] Tejumola, H. O., Integral conditions of existence and non-existence of periodic solutions of some sixth and fifth order ordinary differential equations. *J. Nigerian Math. Soc.* 31 (2012), 2333.
- [6] Tunç, C., Further results on the instability of solutions of certain nonlinear vector differential equations of fifth order. *Appl. Math. Inf. Sci.* 2 (2008), no. 1, 5160.
- [7] Tunç, C., Instability of solutions for certain nonlinear vector differential equations of fourth order. *Nelnn Koliv.* 12 (2009), no. 1, 120129; translation in *Nonlinear Oscil. (N. Y.)* 12 (2009), no. 1, 123132.
- [8] Tunç, C., An instability theorem for a certain fifth-order delay differential equation. *Filomat* 25 (2011), no. 3, 145151.
- [9] Tunç, C., On the instability of solutions of some fifth order nonlinear delay differential equations. *Appl. Math. Inf. Sci.* 5 (2011), no. 1, 112121.
- [10] Tunç, C., Instability of non-linear functional differential equations of fifth order. *ITB J. Sci.* 44A (2012), no. 3, 239249.
- [11] Tunç, C., Instability for nonlinear differential equations of fifth order subject to delay. *Nonlinear Dyn. Syst. Theory* 12 (2012), no. 2, 207214

A note on certain qualitative properties of solutions in Volterra integro-differential equations

OSMAN TUNÇ

Van Yuzuncu Yil University, Van, Turkey

email: osmantunc89@hotmail.com

Volterra integral and integro differential equations with and without delay are essential tools in sciences and many fields. Because of this reality, in the recent years the qualitative behaviors of solutions of differential, integral and integro differential equations have extensively been discussed and are still being investigated by numerous authors. In this work, we consider three nonlinear Volterra integro-differential equations. We investigated boundedness and exponential stability of the solutions by Lypanuov functional. The results improve some the results that can found in literature.

MSC 2010: 45D05, 45M10, 45Jxx

Keywords: Volterra integro-differential equation, first order, exponential stability, boundedness, Lyapunov function

References

- [1] M. Advar and Y. N. Raffoul, Inequalities and exponential stability and instability in finite delay Volterra integro-differential equations. *Rend. Circ. Mat. Palermo* **61** (2012), no. 3, 321-330.
- [2] L. C. Becker, Principal matrix solutions and variation of parameters for a Volterra integro-differential equation and its adjoint. *Electron. J. Qual. Theory Differ. Equ.* (2006), no. 14, 1-22.
- [3] L. C. Becker, Function bounds for solutions of Volterra equations and exponential asymptotic stability. *Nonlinear Anal.* **67** (2007), no. 2, 382-397.
- [4] L. C. Becker, Uniformly continuous L^1 solutions of Volterra equations and global asymptotic stability. *Cubo* **11** (2009), no. 3, 1-24.
- [5] T. A. Burton, Boundedness and periodicity in integral and integro-differential equations. *Differential Equations Dynam. Systems* **1** (1993), no. 2, 161-172.

Galois theory and palindromic polynomials

FATMA TUTAR¹, ŞENAY BAYDAŞ², BÜLENT KARAKAŞ³

^{1,2,3} Van Yuzuncu Yil University, Van, Turkey

emails: ¹ftutar94@gmail.com; ²senay.baydas@gmail.com; ³bulentkarakas@gmail.com

In this paper, Galois theory, field extension, palindromic polynomial, Galois groups are given. We introduce relation between Galois group and the roots of palindromic polynomials.

MSC 2010: 11R32, 12F10, 08A40

Keywords: Field extension, Galois group, Palindromic polynomials

References

- [1] A. O. Asar and A. Arıkan, Sayılar Teorisi. *Gazi Kitapevi, Ankara*, 2011.
- [2] J. B. Fraleigh, A First Course in Abstract Algebra. *Pearson Education, New York*, 2006.
- [3] P. Lindstorm, Galois Theory of Palindromic Polynomials (Master Thesis). *Universty of Oslo, Oslo, Norway*, 2015.
- [4] I. Stewart, Galois Theory. *Chapman and Hall Ltd, London*, 1945.
- [5] H. Sun, Y. Wang and H. X. Zhang, Polynomials with palindromic and unimodal coefficients. *Acta Mathematica Sinica* **31** (2015), no. 4, 565–575.

Canonical finite Blaschke products and decomposibility

SUMEYRA UCAR¹, NİHAL YILMAZ ÖZGÜR²

^{1,2}Balıkesir University, Balıkesir, Turkey

emails: ¹sumeyraucar@balikesir.edu.tr; ²nihal@balikesir.edu.tr

A canonical Blaschke product of degree n is a function of the following form:

$$B(z) = z \prod_{k=1}^{n-1} \frac{z - a_k}{1 - \overline{a_k}z},$$

where a_k are in the unit disc for $1 \leq k \leq n - 1$. In this study, we discuss a canonical finite Blaschke product B can be written as $B = B \circ M$ where M is a Möbius transformation different from identity using the nonzero zeros of B . Also, we investigate when such Blaschke products is composition of Blaschke products of lower degree.

MSC 2010: 30J10, 30D05, 51M15

Keywords: Finite Blaschke products, unit disc, composition of finite Blaschke products

References

- [1] R. L. Craighead and F. W. Carroll, A decomposition of finite Blaschke products. *Complex Variables Theory Appl.* **26** (1995), no. 4, 333-341.
- [2] U. Daepf, P. Gorkin and K. Voss, Poncelet's theorem, Sendov's conjecture, and Blaschke products. *J. Math. Anal. Appl.* **365** (2010), no. 1, 93-102.
- [3] L. R. Ford, Automorphic functions. *Chelsea Publishing Co., New York*, 1951.
- [4] U. Daepf, P. Gorkin, A. Shaffer, B. Sokolowsky and K. Voss, Decomposing finite Blaschke products. *J. Math. Anal. Appl.* **426** (2015), no. 2, 1201-1216.
- [5] A. L. Horwitz, L. A. Rubel, A Uniqueness Theorem for Monic Blaschke Products. *Proc. Amer. Math. Soc.* **96** (1986), no. 1, 180-182.
- [6] N. Yılmaz Özgür, Finite Blaschke products and circles that pass through the origin. *Bull. Math. Anal. Appl.* **3** (2011), no. 3, 64-72.
- [7] N. Yılmaz Özgür, Some Geometric Properties of Finite Blaschke Products. In: *Riemannian geometry and applications-Proceedings RIGA, Bucharest*, 2011.
- [8] N. Yılmaz Özgür, S. Uçar, On some geometric properties of finite Blaschke products. *Int. Electron. J. Geom.* **8** (2015), no. 2, 97-105.

Applicability of regression analysis on the oxygen enriched combustion of Kutahya-Tuncbilek lignite

OZLEM UGUZ¹, ALI DEMIRCI², HANZADE HAYKIRI ACMA³, SERDAR YAMAN⁴

¹Marmara University, Istanbul, Turkey

^{2,3,4}Istanbul Technical University, Istanbul, Turkey

emails: ¹ozlem.uguz@marmara.edu.tr; ²demircial@itu.edu.tr; ³hanzade@itu.edu.tr;
⁴yamans@itu.edu.tr

Coal is the primary fossil fuel for energy generation. However, combustion of coal causes emission of many pollutant gases and particles to the atmosphere. Thus, clean coal technologies have been developed to minimize the hazards of burning coals in thermal power plants. Among these technologies, oxygen enriched combustion is one of the most commonly applied ones to low quality coals (lignites) of Turkey. In oxygen enriched combustion, lignites are burnt under the atmospheres of elevated oxygen concentrations. Oxygen enriched combustion increases available heat, improves ignition characteristics, reduces exhaust gas volume and increases energy efficiency [1, 2]. In this study, ground Kutahya-Tuncbilek lignite samples were burnt in a horizontal tube furnace at 200°C, 450°C and 800°C under the atmospheres having the shares of 21%O₂ + 79%N₂, 30%O₂ + 70%N₂, 40%O₂ + 60%N₂, 50%O₂ + 50%N₂. After combustion tests, amount of carbon percentages in the burnout lignite sample was obtained by elemental analysis. Dependency of the carbon amount % on the furnace temperature and the oxygen share inside the furnace was investigated by applying 5 different regression models: $y = x_1 + x_2 + 1$ (Model1), $y = x_1^2 + x_2^2 + x_1 * x_2 + x_1 + x_2 + 1$ (Model 2), $y = x_1 * x_2 + x_1 + 1$ (Model 3), $y = x_1 * x_2 + x_2 + 1$ (Model 4), $y = x_1^2 + x_2^2 + x_1 + x_2 + 1$ (Model 5). Here x_1 , x_2 and y stood for the furnace temperature, oxygen share inside the furnace and carbon amount %, respectively. Predicted equations, RMSE values, p-values, R^2 values were obtained for each regression model. It was seen that Model2. It was seen that Model2 gave the best fit to the data with the values of $R^2 = 0.985$, $RMSE = 0.037$ and p-value= $2.06 * 10^{-5}$. Thus, the required oxygen/nitrogen ratio and the furnace temperature could be calculated to burn Kutahya-Tuncbilek lignite in the most effective way without the necessity of experimental work.

MSC 2010: 62J05, 80A25, 00A69

Keywords: Regression analysis, coal, oxygen enriched combustion, energy

References

- [1] J. Deng, Q. Li, Y. Xiao, and H. Wen, The effect of oxygen concentration on the non-isothermal combustion of coal. *Thermochim Acta.* **653** (2017), 106-115; doi: 10.1016/j.tca.2017.04.009.
- [2] [2] S. S. Daood, W. Nimmo, Deep staged, oxygen enriched combustion of coal. *Fuel* **101** (2012), 187-196; doi: 10.1016/j.fuel.2011.02.007.

Applicability of regression analysis on the oxygen enriched combustion of Adiyaman-Golbasi lignite

OZLEM UGUZ¹, ALI DEMIRCI², HANZADE HAYKIRI ACMA³, SERDAR YAMAN⁴

¹Marmara University, Istanbul, Turkey

^{2,3,4}Istanbul Technical University, Istanbul, Turkey

emails: ¹ozlem.uguz@marmara.edu.tr; ²demircial@itu.edu.tr; ³hanzade@itu.edu.tr;
⁴yamans@itu.edu.tr

Coal is predicted to cover about almost one third of the global energy needs in the year of 2021. Because of the high share of coal in energy industry, clean coal technologies have been developed by scientists. Oxygen enriched combustion is one of these technologies as it reduces emissions and increases efficiency [1, 2]. In this study, oxygen enriched combustion was applied to ground low quality Adiyaman- Golbasi lignite at temperatures of 200°C, 400°C and 600°C in the horizontal tube furnace, under the atmospheres with the shares of 21%O₂+79%N₂, 30%O₂+70%N₂, 40%O₂+60%N₂, 50%O₂ + 50%N₂. The carbon amounts % of the burnt samples were obtained by elemental analysis. Regression analysis was performed to the data to verify the carbon amount % left in the burnout sample when the combustion temperature and the oxygen/nitrogen ratio were given. Tested models for regression analysis were as follows: $y = x_1 + x_2 + 1$ (Model1), $y = x_1^2 + x_2^2 + x_1 * x_2 + x_1 + x_2 + 1$ (Model 2), $y = x_1 * x_2 + x_1 + 1$ (Model 3), $y = x_1 * x_2 + x_2 + 1$ (Model 4), $y = x_1^2 + x_2^2 + x_1 + x_2 + 1$ (Model 5). In these equations, x_1 , x_2 and y represented the furnace temperature, oxygen share inside the furnace and carbon amount %, respectively. As a result of the analysis it was found that Model2 gave the most accurate results, since it gave the values of $R^2 = 0.981$, $RMSE = 0.0465$ and p-value= $4.59 * 10^{-5}$. Thus, by applying Model2, required values of combustion temperature and oxygen/nitrogen ratio can be predicted to burn all of the carbon content in Adiyaman- Golbasi lignite.

MSC 2010: 62J05, 80A25, 00A69

Keywords: Regression analysis, coal, oxygen enriched combustion, energy

References

- [1] D. Menage, R. Lemaire, and P. Seers, Experimental study and chemical reactor network modeling of the high heating rate devolatilization and oxidation of pulverized bituminous coals under air, oxygen-enriched combustion (OEC) and oxy-fuel combustion (OFC). *Fuel Process Technol.* **177** (2018), 179-193; doi: 10.1016/j.fuproc.2018.04.025.
- [2] H. P. Kruczek, M. Ostrycharczyk, M. Czerep, M. Baranowski, J. Zgora, Examinations of the process of hard coal and biomass blend combustion in OEA (Oxygen enriched atmosphere). *Energy* **92** (2015), 40-46; 10.1016/j.energy.2015.05.112.

On uniformly pr -ideals in commutative rings

RABIA NAGEHAN UREGEN

Erzincan Binali Yildirim University, Erzincan, Turkey.

email: rabia.uregen@erzincan.edu.tr

Let R be a commutative ring with nonzero identity and I be a proper ideal of R . Then I is said to be a uniformly pr -ideal if there exists $N \in \mathbb{N}$ such that $ab \in I$ with $ann(a) = 0$ implies $b^N \in I$. In that case the smallest integer $N \in \mathbb{N}$ is called the order of I and denoted by $ord_R(I) = N$. Among many results in this presentation, it is given some characterizations and properties of this new classes of ideals similar to prime ideals. Furthermore, it is investigated that the relations between uniformly pr -ideals and some classical ideals such as r -ideals, uniformly primary ideals and strongly primary ideals.

MSC 2010: 13A15, 13E05

Keywords: r -ideal, pr -ideal, primary ideal, uniformly primary ideal

References

- [1] J. A. Cox and A. J. Hetzel, Uniformly primary ideals. *Journal of Pure and Applied Algebra* **212** (2008), no. 1, 1-8.
- [2] R. Mohamadian, r -ideals in commutative rings. *Turkish Journal of Mathematics* **39** (2015), no. 5, 733-749.
- [3] S. Koc and U. Tekir, r -Submodules and sr -Submodules, *Turkish Journal of Mathematics* **42** (2018), no. 4, 1863-1876.

On 2-absorbing ideals

REZVAN VARMAZYAR

Islamic Azad University, Khoy, Iran

email: varmazyar@iaukhoy.ac.ir

Let R be a commutative ring with identity. A proper ideal A of R is called a 2-absorbing ideal if $xyz \in A$ for $x, y, z \in R$ implies that $xy \in A$ or $xz \in A$ or $yz \in A$. In this article, some properties of 2-absorbing ideals are given.

MSC 2010: 13C99

Keywords: Prime ideal; n -absorbing ideal; 2-absorbing ideal.

References

- [1] D. F. Anderson and A. Badawi, On n -absorbing ideals of commutative rings. *Comm. Algebra* **39** (2011), 1646-1672.
- [2] A. Badawi, On 2-absorbing ideals of commutative rings. *Bull. Austral. Math. Soc.* **75** (2007), 417-429.

Studying the kinetic parameters and mechanism of the thermal decomposition (dehydration, dehydroxylation and decarbonylation) of some clays using TG traces

ZEKİ YALÇINKAYA¹, ŞENOL KUBİLAY², ALİ SAVRAN³, NECLA ÇALIŞKAN⁴

^{1,2,3,4} Van Yuzuncu Yil University, Van, Turkey

emails: ¹zeki@yyu.edu.tr; ²senolkubilay@yyu.edu.tr; ³alisavran@yyu.edu.tr;
⁴ncaliskan@hotmail.com

Two methods, Reich-Stivala (R-S) and double-log (D-L), were employed to distinguish one mechanism from among 15 theoretical possibilities for heterogeneous solid-state reactions using TG data [1-3]. As known, the rate of reaction of thermal decomposition of solids can be expressed by general equation

$$\frac{d\alpha}{dt} = Ae^{-E/RT} f(\alpha) = Ae^{-E/RT} (1 - \alpha)^n \quad (1)$$

Where $f(\alpha)$ is a function which depends on the reaction mechanism, A=frequency factor, T=temperature (K), α =conversion, n= reaction order. From eq (1), following eq (2) can readily be obtained (the R-S expression).

$$\log\left[\frac{g(\alpha_1)}{\alpha_2} \frac{T_1 T_2}{(T_1 - T_2)}\right] = \frac{E}{2,303R} \quad (2)$$

From eq (2), the double-log expression can be derived

$$\frac{\log\left[\frac{T_R^2}{g(\alpha_R)} \left(\frac{g(\alpha_1)}{T_1^2}\right)\right] T_1 (T_R - T_2)}{\log\left[\frac{T_R^2}{g(\alpha_R)} \left(\frac{g(\alpha_2)}{T_2^2}\right)\right] T_2 (T_R - T_1)} = Z_1 \quad (3)$$

Where T_R , α_R denote an arbitrary reference temperature and its corresponding conversion, respectively. Thermal gravimetric (TG) curves of clays (Bardaki, Tilkitepe from the Van region, Turkey) were determined. Deformations are defined here as changes of the clay by dehydration, dehydroxylation, and decarbonylation. Equations (2 and 3) were employed to determine the mechanisms of thermal decomposition of the clays. Activation energies related to the deformations calculated from the TG curves and the reaction order are also determined. All the calculations were made with EXCEL and recorded as EXCEL macro, then were used.

MSC 2010: 80A50, 00A69

Keywords: Thermogravimetry, kinetics, mechanism, Excel Macro

References

- [1] L. Reich and S. S. Stivala, Kinetics parameters from thermogravimetric curves. *Thermochimica Acta* **24** (1978), 9-16.
- [2] L. Reich and S. S. Stivala, Kinetic parameters from DTA curves. *Thermochim Acta* **25** (1978), 367-371.
- [3] L. Reich and S. S. Stivala, Correlation coefficients mechanism using TG data. *Thermochim Acta* **34** (1979), 287-292.

Clay transition (dehydration, dehydroxylation and decarboxylation) kinetics by DTA

ZEKİ YALÇINKAYA¹, ALİ SAVRAN², ŞENOL KUBİLAY³, NECLA ÇALIŞKAN⁴

^{1,2,3,4}Van Yüzüncü Yıl University, Van, Turkey

emails: ¹zeki@yyu.edu.tr; ²alisavran@yyu.edu.tr; ³senolkubilay@yyu.edu.tr,
³ncaliskan@hotmail.com

The effects of the kinetics of reactions of the type solid solid + gas on the corresponding differential thermal analysis traces are studied. When a reaction occurs in DTA, the change in the thermal properties of the sample is indicated by a deflection, or peak [1]. The information so obtained is used to analyze the differential thermal patterns of clays. Two samples of the clays were chosen from Bardakçı, Tilkitepe, the Van region, Turkey. The peaks are defined here as changes of the clay by dehydration, dehydroxylation, and decarboxylation.

Reich [2] have developed a method to obtain an approximate expression for the energy of activation and reaction order. We used the method (Eq 1) for determining the kinetic parameters from the shape of the differential thermal analysis peak.

$$\left(\frac{T_1}{T_2}\right)^2 \left[\frac{\Delta T_1}{\Delta T_2}\right] = F(T) = \left(\frac{\bar{a}_{T,1}}{\bar{a}_{T,2}}\right)^n \left[\frac{1 - \left(\frac{\bar{a}_{T,1}}{A_T}\right)^{1-n}}{1 - \left(\frac{\bar{a}_{T,2}}{A_T}\right)^{1-n}}\right], n \neq 1 \quad (1)$$

Where, T =temperature (K), ΔT =height of the DTA curve from the baseline; n =order of reaction, A_T =total thermogram area and $\bar{a} = \int_T^{\infty} \Delta T dT$.

One of previously reported algorithm [3] for the estimation of kinetic parameters, activation energy (E) and reaction order (n), from DTA traces assuming an n -type mechanism, after transformed the algorithm to EXCEL VBA macro, were used.

MSC 2010: 80A50, 00A69

Keywords: Differential thermal analysis, kinetics, mechanism, Excel Macro

References

- [1] H. E. Kissinger, Variation of peak temperature with heating rate in differential thermal analysis. *Journal of Research of the National Bureau of Standards* **57** (1956), no. 4, 217-221.
- [2] L. Reich, Kinetic parameters in polypropylene degradation from DTA traces. *J. Appl. Polymer Sci.* **10** (1966), 465-472.
- [3] L. Reich and S. S. Stivala, Computer-determination kinetics parameters (e and n) from DTA curves. *Thermochimica Acta* **84** (1985), 385-390.

On the weighted pseudo almost periodic solutions of nonlinear functional Nicholson's blowflies equations

RAMAZAN YAZGAN¹, CEMİL TUNÇ²

^{1,2} Van Yuzuncu Yil University, Van, Turkey

emails: ¹ryazgan503@gmail.com; ²cemtunc@yahoo.com

In this paper, we prove the existence and the global exponential stability of the unique weighted pseudo almost-periodic solution of a class generalized Nicholson's blowflies model with a linear harvesting term and mixed delays. Some sufficient conditions are given for the existence and the global exponential stability of equation considered by using point theorem and differential inequality techniques. The results of this paper complement the previously known ones. Finally, an illustrative example is given to demonstrate the effectiveness of our results.

MSC 2010: 34K14,34K20,92D25

Keywords: Weighted pseudo almost periodic solution, Nicholson's Blowflies model, Fixed point.

References

- [1] T. Diagana, Existence of weighted pseudo almost periodic solutions to some non-autonomous differential equations. *Int. J. Evol. Equ.* **2** (2008), 397–410.
- [2] T. Diagana, Weighted pseudo-almost periodic solutions to some differential equations. *Nonlinear Anal.* **8** (2008), 2250-2260.
- [3] B. Liu and C. Tunc, Pseudo almost periodic solutions for a class of nonlinear Duffing system with a deviating argument. *J. Appl. Math. Comput.* **49** (2015), 233-242.
- [4] L. Zhao and Y. Li, Global exponential stability of weighted pseudo-almost periodic solutions of neutral type high-order Hopfield neural networks with distributed delays. *Abstr. Appl. Anal.* **2014** (2014), 17 pages.
- [5] R. Yazgan, and C. Tunç, On the weighted pseudo almost periodic solutions of nonlinear functional Duffing equation. *App. Math. and Inform. Sci.*, **11** (2017), no. 6, 1609-1614.

A new generalization of Szász operators and its approximation properties

SERDAL YAZICI¹, BAYRAM ÇEKİM²

^{1,2}Gazi University, Ankara, Turkey

e-mails: ¹serdal.yazici@gazi.edu.tr ; ²bayramcekim@gazi.edu.tr

In this presentation, we attempt to introduce a new generalization of the Szász operators including Hermite polynomials with two variable and shed light on their approximation features with help of the classical modulus of continuity, Peetre's ω -K functional, the class of Lipschitz functions, second modulus of continuity, Voronovskaya type asymptotic formula for these operators.

MSC 2010: 41A25, 41A35

Keywords: Hermite polynomials, modulus of continuity, Lipschitz continuity, approximation theory, Voronovskaya type asymptotic formula.

References

- [1] Z. Ditzian, V. Totik, Moduli of Smoothness. *Springer-Verlag, New York*, 1987.
- [2] R. A. DeVore, G.G. Lorentz, Constructive Approximation. *Springer-Verlag, Berlin*, 1993.
- [3] A. D.Gadzhiev, The convergence problem for sequence of positive linear operators on unbounded sets and theorem analogues to that of P.P. Korovkin. *Sov.Math.Dokl.* **15** (1974), no. 5, 1436–1453.
- [4] P. P. Korovkin, On convergence of linear positive operators in the space of continuous functions (Russian). *Doklady Akad. Nauk. SSSR (NS)* **90** (1953), 961–964.
- [5] G. Krech, A note on some positive linear operators associated with the Hermite polynomials. *Carpathian J. Math.* **32** (2016), no. 1, 71–77.
- [6] D. D. Stancu, Approximation of function by a new class of polynomial operators. *Rev.Rourn.Math. Pures et Appl.* **13** (1968), no. 8, 1173–1194.

Characterization of regular morphisms in terms of abelian categories

TÜLAY YILDIRIM

Gebze Technical University, Kocaeli, Turkey

email: tyildirim@gtu.edu.tr

In this talk, we focus on the regular morphisms in abelian categories which asserts that a morphism $f : M \rightarrow N$ in an abelian category \mathbb{C} is called regular if there exists a morphism $g : N \rightarrow M$ such that $f = fgf$. In the present paper, we establish some further results involving regular morphisms in abelian categories which extend known properties for modules and prove a property on regular compositions of morphisms. We generalize from modules to abelian categories the concept of consecutive pair of morphisms, recently introduced by Facchini and Leroy [1]. Inspired by Ara, Goodearl, OMeara and Pardo [3], equivalent morphisms in abelian categories are taking into the consideration. Moreover, we consider one-sided unit regular morphisms in categories, based on Ehrlich's concept of one-sided unit regular element in a ring [2]. In this talk, after a brief introduction to the subject is discussed, a recent result with a joint work S. Crivei, M. Tamer Koşan and Tülay Yıldırım [4] among these lines will be presented.

MSC 2010: 34B05, 34A08

Keywords: Abelian category, regular morphism, consecutive pair of morphisms, equivalent morphisms, unit regular morphism.

Acknowledgement: We would like to thank professors Alberto Facchini and Grigore Calugareanu for insightful comments and suggestions on an earlier version of the paper. Also, M. Tamer Koşan and Tülay Yıldırım are supported by TUBITAK (117F070).

References

- [1] A. Facchini and A. Leroy, Elementary matrices and products of idempotents. *Linear Multilinear Algebra* **64** (2016), 19161935.
- [2] G. Ehrlich, Units and one-sided units in regular rings. *Trans. Amer. Math. Soc.* **216** (1976), 8190.
- [3] P. Ara, K. R. Goodearl, K. C. OMeara and E. Pardo, Diagonalization of matrices over regular rings. *Linear Algebra Appl.* **265** (1997), 147163.
- [4] S. Crivei, Tamer Kusan and T. Yildirim, On regular morphisms in abelian categories. *preprint* 2018.

Statistical inference for the inverse Weibull distribution

ASUMAN YILMAZ¹, MAHMUT KARA²

^{1,2}Van Yuzuncu Yil University, Van, Turkey

emails: ¹asumanduva@gmail.com; ²mkara2581@gmail.com

The inverse Weibull distribution is known as type 2 extreme value or Frechet distribution. This distribution has been used to model, many real life applications such as degradation of mechanical components, engineering, hydrological and ecological.

$$f(x) = \frac{\alpha}{\beta} \left(\frac{x}{\beta}\right)^{-(\alpha+1)} e^{-(x/\beta)^{-\alpha}}, \quad x > 0, \quad \alpha, \beta > 0$$

and

$$F(x) = e^{-(x/\beta)^{-\alpha}}, \quad x > 0, \quad \alpha, \beta > 0,$$

respectively. In this study, parameter estimation of inverse Weibull distribution are given by using the methods such as maximum likelihood, L- moment, least square, weighted least square, approximate Bayesian estimates with Lindleys method. Furthermore the performances of the obtained estimators are compared with respect to their biases, log-likelihood values, Q-Q plots, the density plots, distribution plots, distribution plots, AIC, BIC and mean square errors through a simulation study. At the end of study, the procedure is illustrated based on real data.

MSC 2000: 60J10, 60J22, 11K06

Keywords: Inverse Weibull distribution, maximum likelihood method, Lindley method.

References

- [1] C. B. Guure and N. A. Ibrahim, Approximate Bayesian estimates of Weibull parameters with Lindleys method. *Sains Malaysiana* **43** (2011), no. 9, 1433-1437.
- [2] F. R. S. Gusmao and E. M. Ortega, The generalized inverse Weibull distribution. *Statistical Papers* **52** (2011), no. 3, 591-619.
- [3] A. O. M. Ahmed and N. Akma, Bayesian survival estimator for Weibull distribution with censored data. *Journal of Applied Sciencies* **11** (2011), no. 2, 393-396.

Existence of solutions for a system of coupled fractional boundary value problems

FULYA YORUK DEREN

Ege University, Izmir, Turkey

email: fulya.yoruk@ege.edu.tr

In this study, we are interested in the existence and multiplicity of positive solutions for a system of fractional differential equations subject to Riemann Stieltjes integral boundary conditions. Our analysis is based upon some theorems from fixed point theory. Finally, an example is given to illustrate our main result.

MSC 2010: 34B10, 34B18, 39A10.

Keywords: Multiple positive solution, fractional differential equation, fixed point theorem.

Acknowledgement: This work is supported by Ege University, Scientific Research Project (BAP), Project Number: 2015 FEN 069.

References

- [1] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, Theory and applications of fractional differential equations, in: North-Holland Mathematics Studies. *Elsevier Science B.V, Amsterdam*, 2006.
- [2] X. Su, Boundary value problem for a coupled system of nonlinear fractional differential equations. *Applied Mathematics Letters* **22** (2013), 985-1008.
- [3] J. Henderson, R. Luca, Positive solutions for a system of nonlocal fractional boundary value problems. *Fractional Calculus and Applied Analysis* **15** (2009), no. 4, 64-69.

Input distinguishability of linear dynamic control systems

AWAIS YOUNUS¹, HONGWEI LOU²

^{1,2} Fudan University, Shanghai, China

¹ Bahauddin Zakariya University, Multan, Pakistan

email: awais@bzu.edu.pk; hwlou@fudan.edu.cn

When observabilities of hybrid dynamic systems are considered, the distinguishability of subsystems takes a very important role. Necessary and sufficient conditions for distinguishability of linear dynamic systems are obtained. The main result state that the input distinguishability of dynamic control systems is equivalent to nontrivial zero dynamics of the systems.

MSC 2010:

Keywords: Distinguishability, hybrid dynamic systems, zero dynamics

Abstracts of posters

On Caputo and Riemann-Liouville fractional-order derivatives with fixed memory length

MOHAMMED SALAH ABDELOUAHAB¹, SAFA BOURAFA²

^{1,2}Mila University center, Mila, Algeria

email: ¹ medsalah3@yahoo.fr

Recently some researchers have demonstrated that the fractional-order derivative of a non-constant periodic function is not a periodic function with the same period [1, 3, 2, 4, 5]. As a consequence of this property the time-invariant fractional order systems does not have any non-constant periodic solution unless the lower terminal of the derivative is $\pm\infty$, which is not practical. This property limits the applicability of the fractional derivative and makes it unfavorable for periodic real phenomenon [6]. Therefore, enlarging the applicability of fractional systems to such periodic real phenomenon is an important research topic. In this work, we extend the modification of the Grünwald-Letnikov definition of fractional derivative introduced in [6] to the Caputo and Riemann-Liouville fractional-order derivatives. This modification consists of fixing the memory length and varying the lower terminal of the derivative. It is shown that the modified definition of fractional derivative preserves the periodicity.

MSC 2010: 34A08, 34A12, 37C27

Keywords: Fractional-order derivative, memory length, periodic solution

References

- [1] M. S. Tavazoei, A note on fractional-order derivatives of periodic functions. *Automatica* **46** (2010), 945-948.
- [2] M. S. Tavazoei and M. Haeri, A proof for non existence of periodic solutions in time invariant fractional order systems. *Automatica* **45** (2009), 1886-1890.
- [3] M. S. Tavazoei, M. Haeri, M. Attari, S. Bolouki and M. Siami, More details on analysis of fractional-order van der pol oscillator. *Journal of Vibration and Control* **15** (2009), no. 6, 803-819.
- [4] M. Yazdani and H. Salarieh, On the existence of periodic solutions in time-invariant fractional order systems. *Automatica* **47** (2011), 1834-1837.
- [5] E. Kaslik and S. Sivasundaram, Non-existence of periodic solutions in fractional-order dynamical systems and a remarkable difference between integer and fractional-order derivatives of periodic functions. *Nonlinear Analysis: Real World Applications* **13** (2012), 1489-1497.
- [6] M. S. Abdelouahab and N. Hamri, The Grünwald-Letnikov fractional-order derivative with fixed memory length. *Mediterr. J. Math.* **13** (2016), 557-572.

On approximation by generalized Bernstein-Durrmeyer operators

ECEM ACAR¹, AYDIN İZGİ²

^{1,2}Harran University, Şanlıurfa, Turkey

¹karakusecem@harran.edu.tr; ²a-izgi@harran.edu.tr

The purpose of this paper is to introduce the generalized Bernstein-Durrmeyer type operators and obtain some approximation properties of these operators studied in the space of continuous functions of two variables on a compact set. The rate of convergence of these operators are given by using the modulus of continuity. The degree of approximation for the Lipschitz class of functions and the Voronovskaja type asymptotic theorem are studied and some differential properties of these operators are proved. Furthermore, the convergence of the operators by illustrative graphics in Maple to certain functions for two dimensional cases are given.

MSC 2010: 41A10, 41A25, 41A36

Keywords: Approximation, Bernstein-Durrmeyer Operators, modulus of continuity, rate of convergence, Voronovskaja type theorem

References

- [1] S. Bernstein, Démonstration du théorème de Weierstrass, fondé sur le calcul des probabilités. *Commun. Soc. Math. Kharkov* **13** (1912), no. 1, 1-2, 1912.
- [2] P. P. Krovkin, On the convergence of linear positive operators in the space of continuous functions. *Dokl. Akad. Nauk* **90** (1953), 961-964.
- [3] F. L. Martinez, Some Properties of Two-Dimensional Bernstein Polynomials. *Journal of Approximation Theory* **59** (1989), 300-306.
- [4] A. İzgi, Approximation by a class of new type Bernstein polynomials of one two variables. *Global Journal of Pure and Applied Mathematics* **8** (2012), no. 5, 55-71.
- [5] D. Cárdenas-Morales and V. Gupta, Two families of Bernstein-Durrmeyer type operators. *Applied Mathematics and Computation* **248** (2014), 342-353.
- [6] U. Abel, V. Gupta and R. N. Mohapatra, Local approximation by a variant of Bernstein-Durrmeyer operators. *Nonlinear Analysis* **68** (2008), 3372-3381.
- [7] V. Gupta, Different Durrmeyer Variants of Baskakov Operators. *Topics in Mathematical Analysis and Applications* **94** (2014), 419-446.
- [8] V. I. Volkov, On the convergence of sequences of linear positive operators in the space of continuous functions of two variables. *Dokl. Akad. Nauk SSSR* **115** (1957), 17-19.

A fuzzy methodology on surface representation of greenhouse gas estimation

FİLİZ KANBAY¹, NURTEN VARDAR²

^{1,2}Yildiz Technical University, Istanbul, Turkey

emails: ¹fkanbay@yildiz.edu.tr; ²vardar@yildiz.edu.tr;

In this study, the greenhouse gas emissions emitted from transit ships passing through the Bosphorus are predicted by using fuzzy inference system. The effect of the cruising speed and gross tonnage changes of ships are taken into consideration in a fuzzy model. The results are given as surfaces.

MSC 2010: 65D18, 68T27, 03B52

Keywords: Greenhouse gas, ship, fuzzy, surface

References

- [1] A. Kilic, Marmara Denizinde Gemilerden Kaynaklanan Egzoz Emisyonları. *BA FBE Dergisi* **11** (2009), no. 2, 124–134.
- [2] U. Kesgin, N. Vardar, A study on Exhaust gas Emissions from Ships in Turkish Straits. *Atmospheric Environment* **35** (2001) 1863–1870.
- [3] C. Trozzi, R. Vaccoro, Methodologies for Estimating Air Pollutant Emissions From Ships. *Techne Report MEET. (Methodologies for Estimating Air Pollutant Emissions from Transport) RF98*, 1998.
- [4] C. Deniz, D. Yalcin, Estimating Shipping Emissions in the Region of the Sea of Marmara. *Turkey Science of the Total Environment* **390** (2008).

Asymptotic aspect of some functional equations

MOHAMMAD BAGHER MOGHIMI

University of Mohaghegh Ardabili, Ardabil, Iran

email: moghimi@uma.ac.ir

In this note we study the asymptotic stability behavior of some functional equations such as Jensen $2f(\frac{x+y}{2}) = f(x) + f(y)$, quadratic $f(x+y) + f(x-y) = 2f(x) + 2f(y)$ and Drygas $f(x+y) + f(x-y) = 2f(x) + f(y) + f(-y)$ functional equations. We show that if a functional equation of these type functional equations holds approximately for large arguments with an upper bound, then it also holds approximately everywhere with a new upper bound which is a constant multiple of its bound.

MSC 2010: 39B82, 39B62

Keywords: Functional equation, Jensen equation, quadratic equation, Drygas equation, asymptotic stability

References

- [1] A. Bahyrycz, Zs. Pales and M. Piszczek, Asymptotic stability of the Cauchy and Jensen functional equations. *Acta Math. Hungar.* **150** (2016), 131–141.
- [2] B. Khosravi, M. B. Moghimi and A. Najati, Asymptotic aspect of Drygas, quadratic and Jensen functional equations in metric Abelian groups. *Acta Math. Hungar.* 1-18; <https://doi.org/10.1007/s10474-018-0807-x>.
- [3] M. Piszczek and J. Szczawinska, Stability of the Drygas functional equation on restricted domain, *Results. Math.* **68** (2015), 11–24.

Existence of positive solutions for second order impulsive boundary value problems on the half-line

ILKAY YASLAN KARACA¹, AYCAN SINANOĞLU ARISOY²

^{1,2}Ege University, Izmir, Turkey

emails: ¹ilkay.karaca@ege.edu.tr; ²aycansinanoglu@gmail.com;

This paper deals with the existence of positive solutions for a second-order impulsive boundary value problem on the half-line. Our existence result is based on a fixed point theorem and a compactness argument.

MSC 2010: 34B18, 34B37, 34B40

Keywords: Impulsive boundary value problem, infinite interval, fixed point theorem, positive solutions

References

- [1] D. Guo, A class of second order impulsive integro-differential equations on unbounded domain in a Banach space. *Appl. Math. Comput.* **125** (2002), 59-77.
- [2] H. Chen and J. Sun, An application of variational method to second-order impulsive differential equations on the half line. *Appl. Math. Comput.* **217** (2010), 1863-1869.
- [3] D. Guo and V. Lakshmikantham, Nonlinear Problems in Abstract Cones. *Academic Press, Boston*, 1988.

Index

- Öktem, Nuray, 111
İğret Araz, Seda, 86
İkikardes, Sebahattin, 118
İnç, Mustafa, 99
İncesu, Muhsin, 102
İzgi, Aydın, 154
Çalışkan, Abdussamet, 126, 127
Çalışkan, Necla, 144, 145
Çekim, Bayram, 147
Çitil, Mehmet, 104
Şahin, Abdulgani, 123
Şahin, Bünyamin, 123
Şahinkaya, Serap, 124
Şen, Erdoğan, 125
Şenyurt, Süleyman, 126, 127

A'zami, Jafar, 55
Abdelouahab, Mohammed Salah, 153
Acıgoz, Mehmet, 71
Acar, Ecem, 154
Agarwal, Ravi, 27
Ahmadkhanlu, Asghar, 34
Akalan, Evrim, 35
Akbulut, İrem, 36
Akturk, Tolga, 37, 77
Akyüz, Hayriye Esra, 39
Akyol, Mehmet Akif, 38
Alaloush, Mohanad, 130
Aldemir, Mehmet Şerif, 43, 44, 63, 72
Alsoy-Akgün, Nagehan, 40
Altınok, Hıfı, 41, 42
Altıntan, Derya, 45
Altay Demirbağ, Sezgin, 52
Altun, Yener, 46, 47
Aracı, Serkan, 71
Aral, Nazlım Deniz, 48
Arslan, Derya, 49
Asil, Vedat, 68
Aslan, Reşat, 50
Atan, Özkan, 51, 103
Atasever, Merve, 52
Atici Endes, Emine, 53
Aydın, İlknur, 65
Aydın, İzgi, 50
Aydogmus, Ozgur, 54

Baş, Selçuk, 56, 68, 100
Baleanu, Dumitru, 25
Bas, Erdal, 110

Baydaş, Şenay, 57, 70, 88, 135, 138
Behravesht, Hoshang, 58
Bekar, Murat, 59
Bektaş, Atilla, 84
Bespalova, Elena, 60
Bilgin, Tunay, 90
Bourafa, Safa, 153
Bozkaya, Canan, 61
Bugay, Leyla, 62
Bulut, Hasan, 37

Cancan, Murat, 63, 72
Cengiz, Cansu, 64
Cetinkaya, F. Ayca, 65
Chehrazi, Akram, 66
Cheng, Yen-Jen, 26
Coşkun, Fatih, 98

Dörma, Vedat, 70
Demirci Akarsu, Emek, 67
Demirci, Ali, 140, 141
Demirkol, Rıdvan Cem, 56, 68
Deniz, Derya, 41, 42
Denizler, İsmail Hakkı, 69
Denizler, I. Hakkı, 101
Duran, Ugur, 71
Duzce, Serkan Ali, 64

Ediz, Süleyman, 63, 72
Ekinci, Fatma, 112
Ercan, Ahu, 110
Erdogan, Fevzi, 115, 116
Erdur, Sultan, 73
Erkan, Esra, 74
Ertem Akbaş, Elif, 75
Erturk, Vedat Suat, 34

Fan, Feng-Lei, 26

Gök, Mustafa, 75
Gözen, Melek, 76
Güleşce Tatlı, Tuba, 79
Güler, Sinem, 78
Günay Akdemir, Hande, 80
Güngör, Şule Yüksel, 82
Güngör, Hakkı, 81
Ghaffarzadeh, Mehdi, 58
Ghasemi, Mohsen, 58
Gorentas, Necat, 101
Gunaydın, Eda, 77

- Gurefe, Yusuf, 77
- Haghnejad Azar, Kazem, 83
- Hamal, Nuket Aykut, 121
- Hanoymak, Turgut, 66, 79, 84
- Haykırı Acma, Hanzade, 140, 141
- Hristova, Snezhana, 27
- Işık, Hüseyin, 85
- Jabbarzadeh, Mohammad Reza, 87
- Körpınar, Talat, 56, 68, 100
- Körpınar, Zeliha, 98–100
- Küsmüş, Ömer, 79
- Kalkan, Bahar, 88
- Kama, Ramazan, 89
- Kanbay, Filiz, 155
- Kara, Mahmut, 149
- Karakaş, Bülent, 57, 70, 88, 135, 138
- Karakuş, Mahmut, 90
- Kasap, Mithat, 41, 42
- Kaya, Yasin, 91
- Kayar, Zeynep, 92
- Kazemi Balgeshir, Mohammad Bagher, 94
- Kazemi, Adel P., 93
- Khajepour, Maryam, 55
- Khalili Golmankhaneh, Alireza, 28
- Khojali, Ahmad, 96
- Kim, Daeyoul, 118
- Koepl, Heinz, 45
- Korkmaz, Erdal, 97
- Kuşak Samancı, Hatice, 102
- Kubilay, Şenol, 144, 145
- Kusmus Omer, 101
- Kutlu, Fatih, 51, 103, 104
- Lakestani, Mehrdad, 29
- Lone, Nisar A., 105
- Lopes, Luiz Guerreiro, 106
- Lou, Hongwei, 151
- Machado, Roselaine Neves, 106
- Malik, M. Y., 114
- Manguri, Abdalla Khdir Abdalla, 43, 44
- Marubayashi, Hidetoshi, 35
- Moghimi, Mohammad Bagher, 156
- Motallebi, Mohammad Reza, 107
- Mucuk, Osman, 133
- O'Regan, Ronald, 27
- Oğur, Oğuz , 108
- Onaran, Sinem, 109
- Ozarslan, Ramazan, 110
- Pişkin, Erhan, 112
- Polat, Murat, 113
- Quynh, Truong Cong, 124
- Rehman, Khalil Ur, 114
- Sahan, Tuncar, 133
- Sakar, Mehmet Giyas, 115, 116
- Saldır, Onur, 115, 116
- Sarıaydın, M. Talat, 100
- Saraç, Bülent, 117
- Sarp, Umit, 118
- Savaş Çelik, Gamze, 122
- Savran, Ali, 144, 145
- Sehribanoglu, Sanem, 119
- Senel, Guzide, 120
- Senlik Cerdik, Tugba, 121
- Sevinç, Zeynep, 48
- Shahmorad, Sedaghat, 30
- Sherratt, Jonathan A., 53
- Sinanoglu Arısoy, Aycan, 157
- Slyn'ko, Vitalii I., 31
- Soydan, Gökhan, 122
- Talaei, Younes, 30
- Tanrıverdi, Tanfer, 128
- Taskesen, Hatice, 129, 130
- Telci, Mustafa, 131
- Temel, Cesim, 132
- Temel, Sedat, 133
- Tor, Ali Hakan, 134
- Tuğrul, Fatih, 135
- Tuğrul, Feride, 104
- Tunç, Cemil, 31, 36, 46, 47, 73, 76, 97, 136, 146
- Tunç, Osman, 137
- Tutar, Fatma, 138
- Ucar, Sumeyra, 139
- Ueda, Akira, 35
- Uguz, Ozlem, 140, 141
- Uregen, Rabia Nagehan, 142
- Vardar, Nurten, 155
- Varmazyar, Rezvan, 143
- Weng, Chih-Wen, 26
- Yüce, Salim, 74
- Yıldırım, Tülay, 148
- Yılmaz Özgür, Nihal, 139

Yılmaz, Asuman, 149

Yalçınkaya, Zeki, 144, 145

Yaman, Serdar, 140, 141

Yaremchenko, Nataliia, 60

Yaslan Karaca, Ilkay, 157

Yaylı, Yusuf, 59

Yayli, Yusuf, 32

Yazıcı, Serdal, 147

Yazgan, Ramazan, 146

Yenerođlu, Mustafa, 56

Yiđit, Abdullah, 47

Yoruk Deren, Fulya, 121, 150

Younus, Awais, 151